



Original Investigation | Neurology

Association of Short-term Change in Leukocyte Telomere Length With Cortical Thickness and Outcomes of Mental Training Among Healthy Adults

A Randomized Clinical Trial

Lara M. C. Puhlmann, MSc; Sofie L. Valk, PhD; Veronika Engert, PhD; Boris C. Bernhardt, PhD; Jue Lin, PhD; Elissa S. Epel, PhD; Pascal Vrtička, PhD; Tania Singer, PhD

Abstract

IMPORTANCE Telomere length is associated with the development of age-related diseases and structural differences in multiple brain regions. It remains unclear, however, whether change in telomere length is linked to brain structure change, and to what extent telomere length can be influenced through mental training.

OBJECTIVES To assess the dynamic associations between leukocyte telomere length (LTL) and cortical thickness (CT), and to determine whether LTL is affected by a longitudinal contemplative mental training intervention.

DESIGN, SETTING, AND PARTICIPANTS An open-label efficacy trial of three 3-month mental training modules with healthy, meditation-naïve adults was conducted. Data on LTL and CT were collected 4 times over 9 months between April 22, 2013, and March 31, 2015, as part of the ReSource Project. Data analysis was performed between September 23, 2016, and June 21, 2019. Of 1582 eligible individuals, 943 declined to participate; 362 were randomly selected for participation and assigned to training or retest control cohorts, with demographic characteristics matched. The retest control cohorts underwent all testing but no training. Intention-to-treat analysis was performed.

INTERVENTIONS Training cohort participants completed 3 modules cultivating interoception and attention (Presence), compassion (Affect), or perspective taking (Perspective).

MAIN OUTCOMES AND MEASURES Change in LTL and CT.

RESULTS Of the 362 individuals randomized, 30 participants dropped out before study initiation (initial sample, 332). Data were available for analysis of the training intervention in 298 participants ($n = 222$ training; $n = 76$ retest control) (175 women [58.7%]; mean [SD] age, 40.5 [9.3] years). The training modules had no effect on LTL. In 699 observations from all 298 participants, mean estimated changes in the relative ratios of telomere repeat copy number to single-copy gene (T/S) were for no training, 0.004 (95% CI, -0.010 to 0.018); Presence, -0.007 (95% CI, -0.025 to 0.011); Affect, -0.005 (95% CI, -0.019 to 0.010); and Perspective, -0.001 (95% CI, -0.017 to 0.016). Cortical thickness change data were analyzed in 167 observations from 67 retest control participants (37 women [55.2%], mean [SD] age, 39.6 [9.0] years). In this retest control cohort subsample, naturally occurring LTL change was related to CT change in the left precuneus extending to the posterior cingulate cortex (mean $t_{161} = 3.22$; $P < .001$; $r = 0.246$). At the individual participant level, leukocyte telomere shortening as well as lengthening were observed. Leukocyte telomere shortening was related to cortical thinning ($t_{77} = 2.38$; $P = .01$; $r = 0.262$), and leukocyte telomere

(continued)

Key Points

Question Is naturally occurring short-term change in leukocyte telomere length related to structural plasticity of the brain, and can telomere length be influenced through mental training?

Findings In this randomized clinical trial of 298 healthy adults, the mental training intervention did not affect leukocyte telomere length. Naturally occurring change in leukocyte telomere length over 3 consecutive 3-month intervals was significantly associated with cortical thickness change in the left precuneus extending to the posterior cingulate cortex.

Meaning This study provides the first evidence to date for an association between short-term change in leukocyte telomere length and brain structure, suggesting that these processes may be mechanistically linked; the mental training used did not influence leukocyte telomere length of healthy, middle-aged adults.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY License.

Abstract (continued)

lengthening was related to cortical thickening ($t_{77} = 2.42$; $P = .009$; $r = 0.266$). All analyses controlled for age, sex, and body mass index.

CONCLUSIONS AND RELEVANCE The findings of this trial indicate an association between short-term change in LTL and concomitant change in plasticity of the left precuneus extending to the posterior cingulate cortex. This result contributes to the evidence that LTL changes more dynamically on the individual level than previously thought. Further studies are needed to determine potential long-term implications of such change in relation to cellular aging and the development of neurodegenerative disorders. No effect of contemplative mental training was noted in what may be, to date, the longest intervention with healthy adults.

TRIAL REGISTRATION ClinicalTrials.gov identifier: [NCT01833104](https://clinicaltrials.gov/ct2/show/study/NCT01833104)

JAMA Network Open. 2019;2(9):e199687. doi:10.1001/jamanetworkopen.2019.9687

Introduction

The length of telomeres (ie, protective chromosomal caps) functions as a biomarker for an individual's health and aging trajectory. Shortened leukocyte telomere length (LTL) has, for example, been associated with greater susceptibility to age-related diseases, including mild cognitive impairment and Alzheimer disease.¹⁻⁶ A large-scale, cross-sectional investigation⁷ also identified correlations between shorter LTL and smaller volumes of several brain regions associated with the development of Alzheimer disease, including the hippocampus, amygdala, temporal lobe, precuneus, and posterior cingulate.^{8,9} Cross-sectional evidence therefore suggests that shorter telomeres are associated with neurodegenerative processes. Less is known, however, about how change in telomere length relates to brain structure.

Shorter telomeres have been associated with genetics¹⁰ and lifestyle factors, such as obesity,¹¹ but also with psychological and emotional strains, including early life adversity, chronic stress, rumination, and loneliness.¹²⁻¹⁷ Leukocyte telomere length has therefore been described as a psychobiomarker that reflects the combined result of physiological and psychological burdens on an individual's health and aging profile.¹⁸ Recent evidence suggests that LTL may change more quickly and dynamically than previously assumed. Leukocyte telomere lengthening, for example, has been observed after 6 months of physical training programs,^{19,20} as well as after an intensive 1-month mental training intervention.²¹ Because telomere lengthening implies a reversal of biological aging processes, these observations have received much attention. However, the biological plausibility of short-term telomere change, herein defined as less than 2 years, remains controversial, particularly for lengthening.²²⁻²⁴


Relating change in LTL with brain structure may provide insight into the biological implications of short-term LTL change. If short-term LTL change reflects biological processes that are generally meaningful for an individual's aging trajectory, such change is unlikely to happen in isolation. Rather, LTL change should be associated with changes in other aging- and health-related markers. Structural brain indices are biomarkers of individual differences in aging and health.²⁵ The first aim of the present study was therefore to investigate whether naturally occurring aging- or lifestyle-related change in LTL over 9 months was related to structural changes in the brain. Cortical thickness (CT) was selected as our measure of brain structure as a more anatomically specific modality than, for example, volumetric measures,^{26,27} and was accordingly expected to be more sensitive to structural changes, including aging-related gray-matter decline.^{27,28}

The second aim of this study was to assess whether training in different mental practices over the same 9-month period could systematically influence LTL, potentially buffering against aging-related shortening or facilitating lengthening. Mental training protocols, such as the mindfulness-based stress reduction program,²⁹ have been found to reduce several psychological strains that are associated with shorter telomeres, including rumination, loneliness, and stress.^{30,31} However, of the 9 studies^{21,32-39} that have investigated LTL in association with mindfulness or meditation, as discussed in a review,⁴⁰ only 2 found evidence for a change in LTL. One study observed LTL after a 1-month retreat (N = 28),²¹ and the other after a 5-year lifestyle intervention (N = 10).³² The remaining 7 studies used less-intensive or shorter interventions, which may in part explain the absence of LTL change.³³⁻³⁹ Evidence for an effect of mental training on LTL therefore appears preliminary and requires replication on a larger scale, which was possible in the present study.

The present investigation was conducted as part of the ReSource Project,⁴¹ a longitudinal mental training study that included training cohorts (TCs) and a retest control cohort (RCC). The RCC was used to address our first aim: to assess potential dynamic associations between LTL and CT. We expected associations with CT change in brain regions previously linked cross-sectionally to LTL.⁷ Nonetheless, we conducted analyses on a whole brain level to be able to detect potential aging- or lifestyle-related CT changes more broadly. The main effects of the ReSource Project training on CT have been reported elsewhere.⁴²

Our second aim focused on a nonoverlapping sample, namely, participants trained in 3 distinct types of mental practices (TCs). These practices were designed to cultivate attention, interoception, and focus on the present moment (Presence); socioaffective capacities, such as compassion (Affect); and sociocognitive skills, such as meta-cognition (Perspective) (Figure 1).^{43,44} All 3 training modules had the potential to influence LTL by buffering the experience of acute or chronic stress.^{31,45} The Presence module could additionally have influenced LTL by reducing rumination,⁴⁶ whereas the Affect and Perspective modules may have had an effect through reduced loneliness or social

Figure 1. Study Design of the ReSource Project



A, Key concepts and core exercises taught during the modules Presence (yellow), Affect (red), and Perspective (green). B, Timeline of the ReSource Project and training sequence per cohort. Retest control participants were recruited in 2 cohorts for logistic reasons but were analyzed jointly. We therefore refer to a single retest control cohort in the text. The displayed study timeline was adapted to most accurately reflect the time points of blood sampling. Test phases for other variables may differ slightly. Samples of retest control cohort I were collected after approximately 2 months of no training before T1 and T2; however, given the smaller sample size of this cohort compared with retest control cohort II, combined RCC sampling occurred approximately every 3 months on average; the same is true for magnetic resonance imaging scans. The full ReSource Project design also included follow-up assessments (T4), but these were not included in the present investigation. Adapted from Singer et al.^{43,44}


isolation.^{42,47} We thus predicted leukocyte telomere lengthening or relative maintenance in the TCs compared with the RCC, which, on average, was expected to show aging-related attrition or no change in LTL. In the case of significant training-induced differences in LTL change, we had planned to subsequently analyze how this systematically induced change relates to structural plasticity of regions associated with LTL in the first aim and exploratively at the whole-brain level.

Methods

Participants

In the context of the ReSource Project,⁴¹ 362 healthy adults were initially recruited and randomized to retest control or training cohorts (RCC, $n=100$; TCs, $n=262$) using bootstrapping without replacement. Following dropout before study commencement, the initial sample consisted of 332 participants (175 women; age, 20-55 years; mean [SD], 40.5 [9.3] years) distributed across the RCC and 3 different training cohorts (RCC, $n=90$; TC1, $n=80$; TC2, $n=81$; TC3, $n=81$) (Figure 2).^{44,48,49} The RCC participants underwent all testing but no training, allowing an estimate of retest effects

Figure 2. Study Flow Diagram



Combined numbers from 2 recruitment periods in 2012-2013 and 2013-2014 are shown. BMI indicates body mass index; CT, cortical thickness; fMRI, functional magnetic resonance imaging; MRI, magnetic resonance imaging; ΔLTL, change in leukocyte telomere length (uncorrected); RCC, retest control cohort; SCID, Structured Clinical Interview for DSM-IV Disorders (Axis I and Axis II)^{48,49}; and TC, training cohort. Adapted from Singer et al.⁴⁴

relevant to some measures of the ReSource Project that are not included herein. All participants were meditation-naïve and extensively screened for mental and physical illness through questionnaires and 2 clinical diagnostic interviews; further details have been reported elsewhere.⁵⁰ Data for the present investigation were collected between April 22, 2013, and March 31, 2015. Participants gave written informed consent, could withdraw from the study at any time, and were financially compensated. The study was conducted in compliance with the Declaration of Helsinki⁵¹ and approved by the research ethics committees of the University of Leipzig and the Humboldt University in Berlin, Germany. The trial protocol is available in [Supplement 1](#). This study followed the Consolidated Standards of Reporting Trials (CONSORT) reporting guideline for randomized clinical trials.

Intervention Design

Data were acquired from participants in the RCC 4 times in intervals of approximately 3 months. The TC participants were tested at similar intervals in which data acquisition followed training in 1 of 3 distinct modules termed Presence, Affect, and Perspective. Training cohorts 1 and 2 practiced the 3 modules in counterbalanced order to account for potential sequence effects. Training cohort 3 completed only the 3-month Affect training to isolate the specific effects of the Presence module from the Affect module (Figure 1B). Further details have been reported elsewhere.⁴³ Each module began with a 3-day silent retreat. For the remainder of the training period, participants attended weekly 2-hour group sessions and performed the 2 core practices of each module at home for approximately 30 minutes daily (Figure 1A). The Presence module aimed to cultivate present moment-focused attention and interoception through 2 widely used meditation techniques: Body Scan and Breathing Meditation. The Affect module was designed to cultivate socioaffective qualities, such as compassion, gratitude, and nonjudgmental acceptance of difficult emotions, using Loving-Kindness Meditation and Affect Dyads. The Affect Dyad is a partner exercise specifically aiming at socioaffective abilities.⁴⁷ The Perspective module targeted sociocognitive and metacognitive analytical abilities. The daily core practices of the Perspective module included Observing-Thoughts Meditation and Perspective dyads. As an analog to the Affect Dyad, the Perspective Dyad teaches cognitive perspective taking on the beliefs and thoughts of oneself and others in a partner format. The detailed trial protocol ([Supplement 1](#)) has also been published elsewhere.⁴⁴

Sampling of LTL

Blood was collected at 4 points (T0-T3) and frozen at -80°C until assay. Genomic DNA was extracted from whole blood (QIAamp DNA blood mini kit; Qiagen). Leukocyte telomere length was measured using quantitative polymerase chain reaction assay as previously described,⁵² and reported as T/S ratios (ie, the relative ratio of telomere repeat copy number to single-copy gene; interassay coefficient of variability, 2.3%). All DNA samples were run twice, each with triplicate wells. If the difference between the 2 runs was greater than 7%, samples were run a third time and the average of the 2 closest values was used for data analysis. DNA samples from the same participant were assayed as 1 batch, using the same reagent lots, and run on the same assay plate.

Magnetic Resonance Imaging Acquisition

Magnetic resonance imaging (MRI) acquisition was carried out on a 3-T scanner (Siemens Verio; Siemens) with a 32-channel head coil. T1-weighted images were acquired using a 3-dimensional magnetization-prepared rapid gradient-echo sequence (176 sagittal slices; repetition time, 2300 milliseconds; echo time, 2.98 milliseconds; inversion time, 900 milliseconds; flip angle, 7° ; field of view, $240 \times 256 \text{ mm}^2$; and matrix, $240 \times 256; 1 \times 1 \times 1 \text{ mm}^3$ voxels). Imaging hardware and console software (Syngo B17; Siemens Healthineers) were held constant throughout data collection.

Processing of Structural Data

The T1-weighted MRIs were processed using FreeSurfer, version 5.1.0, to generate cortical surface models for measurements of CT.⁵³ We chose the most general cross-sectional image processing

procedure to enable baseline data analysis for cross-sectional study goals of the ReSource Project (eg, as used by Valk et al⁵⁴) before the completion of data acquisition, which spanned more than 2 years. The exact processing steps of cortical reconstruction and volumetric segmentation are described in previous publications.⁵⁵⁻⁵⁷ Briefly, T1-weighted images were intensity normalized and skull stripped, followed by a tessellation of the gray matter/white matter cortical boundary and automated topologic correction. Surface deformation was performed along intensity gradients, placing borders of the inner (gray matter/white matter) and outer (gray matter/cerebrospinal fluid) cortical interfaces at the location where the greatest shift in intensity defined the transition to the other tissue class. Cortical thickness was calculated as the shortest distance from the gray matter/white matter boundary to the gray matter/cerebrospinal fluid boundary at each vertex on the tessellated surface and is reported in millimeters.⁵⁸

Preprocessing of Dependent Variables

Change in LTL was calculated as difference scores between each participant's measurements from a set of consecutive testing times (ie, T1 minus T0, T2 minus T1, and T3 minus T2). Specifically, we calculated LTL change as the Verhulst *D* value⁵⁹ (herein termed *DLTL*) to correct for regression to the mean instead of the common practice of controlling for baseline LTL because it has recently been suggested that the latter approach inflates type I error rates.⁶⁰

For statistical analysis of CT, we generated participant-specific CT change maps (Δ CT) by subtracting vertexwise thickness maps of subsequent measurement times. Thickness data at each vertex were normalized before change calculation by regressing out the effects of global thickness to emphasize relative region-specific change patterns.

All difference scores diverging more than 3 SDs from the sample mean difference score were defined as outliers. There were no outliers in the Δ CT data analyzed in the present study. Seven *DLTL* outliers were winsorized to the respective upper or lower boundaries of 3 SD. In addition, MRI data were excluded if they did not pass quality control by 2 independent expert raters (S.L.V. and B.C.B.) owing to excessive movement or artifacts in the T1-weighted MRI images (eTable 5 and eTable 6 in Supplement 2). Five scans were excluded because of low image quality.

Statistical Analysis

Data analysis was conducted between September 23, 2016, and June 21, 2019. All longitudinal analyses were performed using multivariate linear mixed models, which are robust to unbalanced and incomplete data in longitudinal designs,⁶¹ allowing the inclusion of all eligible data points for a given analysis. Body mass index, age, and sex were selected as covariates owing to their established association with LTL^{11,62,63} and brain structure.⁶⁴ We also included the variable time point to control for potential effects of the time of measurement on Δ CT. Detailed model descriptions can be found in eAppendix 1 in Supplement 2. All analyses were conducted with an α threshold of .05 or less.

The whole-brain, linear mixed-model analysis of CT data was performed using SurfStat for Matlab,⁶⁵ first with the predictors *DLTL* and time only, and subsequently with the added variables of age, body mass index, and sex, to control for their potential influence. Statistical results were corrected for multiple comparisons following random field theory by means of a conservative cluster-determining threshold of $P < .005$ and familywise error corrections of $P < .01$ (2-tailed) for 20 mm full width at half maximum, smoothed surface-based thickness data, which correspond to the recently recommended threshold for the analysis of surface-based anatomic CT data.⁶⁶ All follow-up analyses were conducted on cluster extent Δ CT in the region identified to be significantly associated with *DLTL* (eAppendix 1 in Supplement 2 provides further details). Effect sizes were calculated as effect size correlations r through the following formula⁶⁷:

$$r = \frac{t}{\sqrt{(t^2 + df)}}$$

Linear mixed models for the analysis of the training intervention were fit using the function lmer of the package lme4⁶⁸ for the statistics software R, version 3.5.1⁶⁹ (eAppendix 1 in Supplement 2). The effect of any term of interest on DLTL was evaluated by comparing the fit of a full model with a reduced model lacking only the term of interest, by means of likelihood ratio tests.⁷⁰

Results

Participants

Sample sizes and reasons for missing cases of all variables relevant to the current investigation are described in Figure 2 as well as eTable 5 and eTable 6 in Supplement 2. Of the 362 individuals randomly assigned to TCs or RCC, 30 participants dropped out before study initiation (initial sample, 332). For the analysis of the mental training intervention, data were available in 298 participants (175 women [58.7%]; mean [SD] age, 40.5 [9.3] years; 222 assigned to TCs, 76 to the RCC). For the analysis of ΔCT, all required covariates and MRI data were available in 67 RCC participants (37 women [55.2%]; mean [SD] age, 40.5 [9.3] years), providing 167 observations in total (eTable 5 and eTable 6 in Supplement 2). Mean (SD) LTL in T/S ratios at baseline was 1.016 (0.16) and individual-level DLTL ranged from −0.259 to 0.251. Further demographic characteristics are reported in the Table.

LTL Change and CT

The first whole-brain analysis identified a significant association between DLTL and ΔCT in the left precuneus extending to the posterior cingulate cortex (PCC) (mean values of cluster extend test: $t_{164} = 3.21$; $P < .001$; $r = 0.243$; 59% overlap with precuneus, 26% overlap with PCC [Automated Anatomical Labeling atlas]) (Figure 3A). In repeating the analysis with the added variables age, body mass index, and sex, the same region was identified (mean $t_{161} = 3.22$; $P < .001$; $r = 0.246$; 61% overlap with precuneus, 24% overlap with PCC).

In subsequent follow-up cluster-level tests, the association between DLTL and ΔCT in the identified region was also significant when analyzed with an additional random slope term, as well as when controlling for change in leukocyte cell type counts (eAppendix 3 and eTable 1 in Supplement 2). Further follow-up analyses assessed the consistency of the observed phenomenon by testing whether the association between DLTL and ΔCT was reliable across the 3 intervals of change. Linear models showed that, when analyzed separately, change in LTL and CT were significantly associated at 2 of the 3 intervals (T0-T1: $t_{54} = 2.60$; $P = .006$; $r = 0.334$; T1-T2:

Table. Demographic Characteristics at Baseline and per Module^a

Characteristic	Baseline	No Training	No Training MRI	Presence ^b	Affect ^c	Perspective ^b
Observations, No.	285	198	167	145	216	140
Women, No. (%)	166 (58.2)	118 (60.0)	89 (53.0)	85 (58.6)	124 (57.4)	79 (56.4)
Participants, No.	285	76	67	145	216	140
Women, No. (%)	166 (58.2)	46 (60.5)	37 (55.2)	85 (58.6)	125 (57.9)	79 (56.4)
Age, mean (SD), y	40.4 (9.3)	39.5 (9.3)	39.6 (9.0)	41.2 (9.3)	40.8 (9.3)	40.9 (9.6)
BMI, mean (SD)	23.5 (3.2)	23.9 (3.0)	23.9 (2.9)	23.5 (3.3)	23.4 (3.2)	23.6 (3.3)
LTL in T/S ratio, mean (SD)	1.016 (0.16)	1.043 (0.16)	1.040 (0.17)	1.002 (0.16)	0.996 (0.16)	1.011 (0.17)
Smokers, No. (%)	34 (11.9)	8 (10.5)	6 (9.0)	21 (14.5)	28 (13.0)	20 (14.3)
Practice, mean (SD), No./wk						
Meditation	NA	NA	NA	4.76 (1.17)	3.84 (1.31)	3.62 (1.23)
Dyad	NA	NA	NA	NA	3.80 (0.70)	3.51 (0.71)

Abbreviations: BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); LTL, leukocyte telomere length; MRI, magnetic resonance imaging; NA, not applicable; T/S ratio, relative ratio of telomere repeat copy number to single-copy gene.

^a Summary statistics include all participants for whom LTL change data and covariate measures were available. An extensive description of baseline demographic characteristics was reported by Singer et al.⁷¹ eTable 4 in Supplement 2 details LTL

difference scores per time point and module, and eAppendix 2 in Supplement 2 provides descriptive analyses.

^b Combined number of cohorts 1 and 2.

^c Combined number of training cohorts 1 to 3, resulting in a higher number than Presence and Perspective training.


$t_{48} = 3.75$; $P < .001$; $r = 0.476$; T2-T3: $t_{50} = 1.28$; $P = .10$; $r = 0.178$) (Figure 3B). In an exploratory analysis of change from baseline to 9 months, however, we did not find a significant association ($t_{56} = 0.46$; $P = .31$; $r = 0.06$) (eFigure 1 in Supplement 2).

Leukocyte telomere shortening and lengthening manifested themselves in a similar number of observations (Figure 3A, scatterplot). Examining these separately in follow-up analyses revealed that LTL change was positively associated with Δ CT regardless of direction, that is, leukocyte telomere shortening with a greater tendency for cortical thinning ($t_{77} = 2.38$; $P = .01$; $r = 0.262$), and lengthening with a greater tendency for cortical thickening ($t_{77} = 2.42$; $P = .009$; $r = 0.266$) (Figure 3C). Corresponding to the observed association of DLTL with Δ CT in the left hemisphere, exploratory analyses identified a contralateral subthreshold association with Δ CT in the right precuneus/PCC (eAppendix 4 and eFigure 3 in Supplement 2).

Mental Training Intervention

Likelihood ratio tests showed no significant main effect of the training module by time interaction ($\chi^2 = 3.26_3$; $P = .35$) or module alone on ($\chi^2 = 2.20_3$; $P = .53$) DLTL (Figure 4A and B). We could therefore not pursue the hypothesis that systematic training-induced differences in LTL change would be associated with changes in brain structure. Estimated mean change in T/S ratios in model

Figure 3. Association Between Leukocyte Telomere Length Change (DLTL) and Cortical Thickness Change (Δ CT)



A, DLTL was positively associated with Δ CT in the left precuneus/posterior cingulate cortex (PCC). Automated Anatomical Labeling atlas: 61% (59%) overlap with precuneus (primary region); 24% [26%] overlap with PCC (secondary region); percentages in brackets indicate results from analysis without age, body mass index, and sex. B, Association between DLTL and Δ CT at each change interval. C, Association of leukocyte telomere lengthening and shortening with Δ CT. For visual display through scatterplots, Δ CT in the cingulate/PCC region was averaged and plotted against DLTL. Each dot represents 1 observation rather than 1 participant. Up to 3 measures of DLTL were available from the same participant, each from a different change interval and controlled for by the linear mixed-model analysis. Displayed regression lines in panels B and C were derived from linear models fit independently for each subsample (B: 1 sample per time interval; C: separate samples for leukocyte telomere lengthening/shortening). Shaded areas represent 95% CIs. CDT indicates cluster-determining threshold; FWE, familywise error correction.

2, our main model of interest, were, for no training, 0.004 (95% CI, -0.010 to 0.018); Presence, -0.007 (95% CI, -0.025 to 0.011); Affect, -0.005 (95% CI, -0.019 to 0.010); and for Perspective, -0.001 (95% CI, -0.017 to 0.016).


As a follow-up analysis, we explored whether the combined 9-month training period affected DLTL; this was not the case ($F_{1,196} = 1.88$; $P = .17$) (Figure 4C; eAppendix 1 in Supplement 2). Additional exploratory analyses showed that the effect of training remained nonsignificant even when considering practice frequency or participant personality or when focusing on within-participant differences (eAppendix 4, eTable 2, and eTable 3 in Supplement 2). Checking the residuals of the above models through visual inspection and diagnostic tests did not reveal any evidence for influential cases, systematic error, or multicollinearity (eAppendix 5 in Supplement 2).

Discussion

With the present investigation, we sought to improve our understanding of the neural and psychological processes associated with short-term change in LTL through 2 steps. We first related DLTL to structural brain changes over 3 consecutive 3-month intervals in an RCC. Second, we assessed the effect of 3 distinct, 3-month mental training modules on DLTL.

Repeated measures of DLTL in RCC participants were associated with ΔCT in the left precuneus/PCC. When analyzed separately, the association was significant over 2 of the 3 consecutive, 3-month

Figure 4. Raw and Estimated Change in Leukocyte Telomere Length (DLTL) per Training Module



intervals. Precuneus and PCC were previously shown to be linked to LTL in a large, cross-sectional, population-based study.⁷ The longitudinal association between short-term changes in LTL and structural indices of the left precuneus/PCC noted in our study suggests a dynamic association between the 2 phenomena.

To our knowledge, there is only 1 other study associating change in LTL and brain structure. Leukocyte telomere shortening and loss of structural integrity were found to correlate in older adults at a single, 2.9-year follow-up measurement.⁷² In contrast, we observed similar rates of leukocyte telomere shortening and lengthening. Shortening was related to a greater tendency for cortical thinning and lengthening was related to a greater tendency for thickening, even when analyzed separately. The identified association was therefore not predominantly driven by aging-related decline. In an exploratory analysis, Δ LTL from baseline to the 9-month follow-up measurement showed no association with Δ CT, and we cannot say with certainty why this occurred. Our results may, however, indicate that short-term LTL change does not follow a continuous trajectory but rather represents transient change.

Telomere fluctuations that average out over time have previously been observed,⁷³ but there is currently no model for the potential underlying biological processes. One mechanism that could be responsible for the associated changes in LTL and CT may be the action of the cellular enzyme telomerase. Telomerase counteracts telomere shortening by replenishing telomeric DNA during cell division.^{22,74} Increases in telomerase activity have been associated with short-term telomere lengthening, most notably following 6-month physical training protocols,^{20,75} and may also have mediated the lengthening observed in this study. Likewise, cortical thickening could be facilitated by an increased telomerase-associated proliferative capacity of critical support cells in the brain capable of undergoing mitosis in the adult, resulting in greater tissue volume, as previously proposed.⁷ Heightened telomerase activity may therefore lead to associated increases in LTL and CT, assuming that central and peripheral telomerase activity are related. In line with this reasoning, telomerase activity was previously found to correlate positively with hippocampal volume, although not with LTL, in a small sample of patients with major depressive disorder.⁷⁶ While it is unclear which neurobiological mechanisms exactly underlie cortical thickening in the age group evaluated herein, these authors described the potential gliogenesis-enhancing and neuroprotective effects of telomerase in more detail. Concomitant telomere shortening and cortical thinning, if not reflective of aging-related decline, could similarly result from accelerated telomere attrition and decreased cellular proliferation under lowered telomerase activity. In the present study, we could not assess whether such fluctuations in telomerase activity occurred and, if so, whether they result from endogenous processes or external influences, such as the lifestyle factors stress or physical exercise.^{20,77} The same lifestyle variables have also been directly associated with differences in telomere length^{15,20} and brain structure,^{78,79} and therefore provide an explanatory framework for the observed concomitant change that may not involve mediation through telomerase activity.

The present study identified significantly less widespread association between change in LTL and brain structure than was reported by King and colleagues.⁷ Likely reasons for this divergence include King and colleagues'⁷ region of interest-based, cross-sectional study design compared with our whole-brain longitudinal analysis approach, as well as their older and considerably larger sample ($N = 1960$), which sensitized their analyses to associations with very small effect sizes of R^2 less than 1%. Owing to our comparatively smaller participant sample, the present study was not equipped to detect similarly sized effects. Nonetheless, a specific association between LTL change and thickness change in the left precuneus/PCC was identified. The precuneus/PCC region is a prominent metabolic hub and central node of the default mode network.⁸⁰ The observed change in CT may thus, for example, reflect a particularly strong susceptibility to telomerase-related synaptic plasticity grounded in this region's unique metabolic and connectional properties. Moreover, it is notable that precuneus/PCC structure and function are impaired in age-related neurodegenerative disorders, such as Alzheimer disease.^{8,9,81} In cross-sectional studies, shorter telomeres have been associated with similar age-related conditions.³⁻⁶ The specific coupling of short-term LTL change with Δ CT in the

precuneus/PCC could be a window to cellular processes implicated in the development of age-related diseases. The long-term dynamics of this association should be investigated in future research using targeted, longitudinal population studies tracking aging-related changes in telomere length in combination with multimodal MRI.

Contrary to our predictions, we found no effect of contemplative mental training on change in LTL. This result contrasts with the findings of 2 preceding interventions that identified telomere lengthening in relation to meditation.²¹ The first study found telomere lengthening in men with low-risk prostate cancer following a relatively broad, 5-year lifestyle intervention.³² The differing participant population and intervention design may account for the diverging outcomes. The other study found telomere lengthening in healthy, middle-aged participants after a short, but intense, 1-month residential training retreat, which may have unique benefits.³¹ In contrast, our results provide what we believe to be the largest body of evidence that longitudinal contemplative mental training does not systematically lengthen leukocyte telomeres of healthy adults. Because we detected no mean attrition in the RCC over the entire 9-month study period, we cannot make any conclusions regarding potential relative maintenance of telomere length through mental training.

Effects of the ReSource Project training modules on CT alone have been reported.⁴² Because we found evidence for an association between CT and LTL on the individual participant level, but no effect of training on LTL, it appears that the mechanism underlying the herein observed association is independent of the capacities trained during the intervention. Other work showed that the ReSource Project intervention reduced physiological responses to acute stress and perceived social connectedness.^{45,47} Our present findings therefore provide indirect evidence that alleviating these psychophysiological strains does not lead to short-term telomere lengthening in healthy individuals.

Limitations

The present study has limitations. The effect of measurement error on short-term LTL change, particularly lengthening, is controversial.^{23,24} In the present sample, given our interassay coefficient of variability of 2.3% and mean LTL T/S ratio of approximately 1.0, measurement error should, on average, be 0.023. On an individual participant level, observed change ranged from -0.259 to 0.251 and is therefore unlikely predominantly attributable to measurement error, although lengthening or shortening classifications of particularly small change values could be distorted. Regarding the association with CT, measurement error should increase the risk of type II rather than type I error. The herein detected robust associations at multiple intervals of change are therefore unlikely attributable to measurement error. Rather, stronger associations may be found if measurement error was minimized. A further limitation is that, by evaluating structural change using CT, we were unable to examine potential associations with subcortical or allocortical regions, such as the hippocampus, which has been associated with LTL in cross-sectional studies.⁸²

Conclusions

Our findings contribute to the evidence that LTL changes more dynamically on the individual level than previously thought and indicate that short-term LTL change is associated with structural brain alteration. Further studies will need to identify the long-term implications of such changes in relation to cellular aging and the development of neurodegenerative disorders, as well as how to activate protective processes that influence LTL. In contrast to our hypotheses and some earlier reports, LTL of healthy adults was not influenced through contemplative mental training over 9 months.

ARTICLE INFORMATION

Accepted for Publication: June 25, 2019.

Published: September 25, 2019. doi:[10.1001/jamanetworkopen.2019.9687](https://doi.org/10.1001/jamanetworkopen.2019.9687)

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2019 Puhlmann LMC et al. *JAMA Network Open*.

Corresponding Author: Lara M. C. Puhlmann, MSc, Research Group “Social Stress and Family Health,” Max Planck Institute for Human Cognitive and Brain Sciences, Stephanstrasse, 1a, 04103 Leipzig, Germany (puhlmann@cbs.mpg.de).

Author Affiliations: Research Group, “Social Stress and Family Health,” Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany (Puhlmann, Engert, Vrtička); Institute of Systems Neuroscience, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany (Valk); Institute of Neuroscience and Medicine (INM-7: Brain & Behaviour), Research Centre Jülich, Jülich, Germany (Valk); Montreal Neurological Institute, McGill University, Montreal, Québec, Canada (Bernhardt); Department of Biochemistry and Biophysics, University of California, San Francisco (Lin); Department of Psychiatry, University of California, San Francisco (Epel); Social Neuroscience Lab, Max Planck Society, Berlin, Germany (Singer).

Author Contributions: Drs Vrtička and Singer share senior authorship. Ms Puhlmann had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Puhlmann, Valk, Engert, Epel, Singer.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Puhlmann, Engert, Vrtička.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Puhlmann, Valk, Vrtička.

Obtained funding: Singer.

Administrative, technical, or material support: Lin, Singer.

Supervision: Valk, Engert, Epel, Vrtička, Singer.

Conflict of Interest Disclosures: Dr Lin is a paid consultant to Telomere Diagnostics Inc, formerly Telome Health, and owns stock in the company; the company had no role in this research. Dr Singer reported receiving grants from the European Research Council during the conduct of the study. No other disclosures were reported.

Funding/Support: Dr Singer, as principal investigator, received funding for the ReSource Project from the European Research Council under the European Community's Seventh Framework Program (FP7/2007-2013/ERC grant agreement 205557) and from the Max Planck Society.

Role of the Funder/Sponsor: The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Sharing Statement: See [Supplement 3](#).

Additional Contributions: We thank the members of the former Department of Social Neuroscience involved in the ReSource Project over many years, in particular all ReSource Project teachers who taught the intervention program. Astrid Ackermann, Christina Bochow, PhD, Matthias Bolz, and Sandra Zurborg, PhD, managed the large-scale longitudinal study; Hannes Niederhausen, Henrik Grunert, and Torsten Kästner provided technical support; Sylvia Tydeks and Elisabeth Murzik recruited participants; and Manuela Hofmann, Sylvie Neubert, and Nicole Pampus helped with recruitment and data collection in the ReSource Project in general. All of these contributors were under a regular contract in the Department of Social Neuroscience at the Max Planck Institute of Human Cognitive and Brain Sciences, and no additional compensation was received by any outside funding sponsor.

Additional Information: This study forms part of the ReSource Project, headed by Tania Singer. Data for this project were collected between 2013 and 2016 at the former Department of Social Neuroscience at the Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany.

REFERENCES

1. Willeit P, Willeit J, Kloss-Brandstätter A, Kronenberg F, Kiechl S. Fifteen-year follow-up of association between telomere length and incident cancer and cancer mortality. *JAMA*. 2011;306(1):42-44. doi:10.1001/jama.2011.901
2. Sanders JL, Fitzpatrick AL, Boudreau RM, et al. Leukocyte telomere length is associated with noninvasively measured age-related disease: The Cardiovascular Health Study. *J Gerontol A Biol Sci Med Sci*. 2012;67(4):409-416. doi:10.1093/gerona/glr173
3. Martin-Ruiz C, Dickinson HO, Keys B, Rowan E, Kenny RA, Von Zglinicki T. Telomere length predicts poststroke mortality, dementia, and cognitive decline. *Ann Neurol*. 2006;60(2):174-180. doi:10.1002/ana.20869
4. Hochstrasser T, Marksteiner J, Humpel C. Telomere length is age-dependent and reduced in monocytes of Alzheimer patients. *Exp Gerontol*. 2012;47(2):160-163. doi:10.1016/j.exger.2011.11.012

5. Panossian LA, Porter VR, Valenzuela HF, et al. Telomere shortening in T cells correlates with Alzheimer's disease status. *Neurobiol Aging*. 2003;24(1):77-84. doi:[10.1016/S0197-4580\(02\)00043-X](https://doi.org/10.1016/S0197-4580(02)00043-X)
6. Thomas P, O'Callaghan NJ, Fenech M. Telomere length in white blood cells, buccal cells and brain tissue and its variation with ageing and Alzheimer's disease. *Mech Ageing Dev*. 2008;129(4):183-190. doi:[10.1016/j.mad.2007.12.004](https://doi.org/10.1016/j.mad.2007.12.004)
7. King KS, Kozlitina J, Rosenberg RN, Peshock RM, McColl RW, Garcia CK. Effect of leukocyte telomere length on total and regional brain volumes in a large population-based cohort. *JAMA Neurol*. 2014;71(10):1247-1254. doi:[10.1001/jamaneurol.2014.1926](https://doi.org/10.1001/jamaneurol.2014.1926)
8. Buckner RL, Snyder AZ, Shannon BJ, et al. Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J Neurosci*. 2005;25(34):7709-7717. doi:[10.1523/JNEUROSCI.2177-05.2005](https://doi.org/10.1523/JNEUROSCI.2177-05.2005)
9. Reiman EM, Jagust WJ. Brain imaging in the study of Alzheimer's disease. *Neuroimage*. 2012;61(2):505-516. doi:[10.1016/j.neuroimage.2011.11.075](https://doi.org/10.1016/j.neuroimage.2011.11.075)
10. Codd V, Nelson CP, Albrecht E, et al; CARDIoGRAM consortium. Identification of seven loci influencing mean telomere length and their association with disease. *Nat Genet*. 2013;45(4):422-427, e1-e2. doi:[10.1038/ng.2528](https://doi.org/10.1038/ng.2528)
11. Gielen M, Hageman GJ, Antoniou EE, et al; TELOMAAS group. Body mass index is negatively associated with telomere length: a collaborative cross-sectional meta-analysis of 87 observational studies. *Am J Clin Nutr*. 2018;108(3):453-475. doi:[10.1093/ajcn/nqy107](https://doi.org/10.1093/ajcn/nqy107)
12. Simon NM, Smoller JW, McNamara KL, et al. Telomere shortening and mood disorders: preliminary support for a chronic stress model of accelerated aging. *Biol Psychiatry*. 2006;60(5):432-435. doi:[10.1016/j.biopsych.2006.02.004](https://doi.org/10.1016/j.biopsych.2006.02.004)
13. Wilson SJ, Woody A, Padin AC, Lin J, Malarkey WB, Kiecolt-Glaser JK. Loneliness and telomere length: immune and parasympathetic function in associations with accelerated aging. *Ann Behav Med*. 2019;53(6):541-550.
14. Epel ES, Puterman E, Lin J, Blackburn E, Lazaro A, Mendes WB. Wandering minds and aging cells. *Clin Psychol Sci*. 2013;1(1):75-83. doi:[10.1177/2167702612460234](https://doi.org/10.1177/2167702612460234)
15. Pepper GV, Bateson M, Nettle D. Telomeres as integrative markers of exposure to stress and adversity: a systematic review and meta-analysis. *R Soc Open Sci*. 2018;5(8):180744. doi:[10.1098/rsos.180744](https://doi.org/10.1098/rsos.180744)
16. Deng W, Cheung ST, Tsao SW, Wang XM, Tiwari AFY. Telomerase activity and its association with psychological stress, mental disorders, lifestyle factors and interventions: a systematic review. *Psychoneuroendocrinology*. 2016;64:150-163. doi:[10.1016/j.psyneuen.2015.11.017](https://doi.org/10.1016/j.psyneuen.2015.11.017)
17. Humphreys KL, Esteves K, Zeanah CH, Fox NA, Nelson CA III, Drury SS. Accelerated telomere shortening: tracking the lasting impact of early institutional care at the cellular level. *Psychiatry Res*. 2016;246:95-100. doi:[10.1016/j.psychres.2016.09.023](https://doi.org/10.1016/j.psychres.2016.09.023)
18. Epel ES. Telomeres in a life-span perspective: a new "psychobiomarker"? *Curr Dir Psychol Sci*. 2009;18(1):6-10. doi:[10.1111/j.1467-8721.2009.01596.x](https://doi.org/10.1111/j.1467-8721.2009.01596.x)
19. Puterman E, Weiss J, Lin J, et al. Aerobic exercise lengthens telomeres and reduces stress in family caregivers: a randomized controlled trial—Curt Richter Award Paper 2018. *Psychoneuroendocrinology*. 2018;98:245-252. doi:[10.1016/j.psyneuen.2018.08.002](https://doi.org/10.1016/j.psyneuen.2018.08.002)
20. Werner CM, Hecksteden A, Morsch A, et al. Differential effects of endurance, interval, and resistance training on telomerase activity and telomere length in a randomized, controlled study. *Eur Heart J*. 2019;40(1):34-46. doi:[10.1093/eurheartj/ehy585](https://doi.org/10.1093/eurheartj/ehy585)
21. Conklin QA, King BG, Zanesco AP, et al. Insight meditation and telomere biology: The effects of intensive retreat and the moderating role of personality. *Brain Behav Immun*. 2018;70:233-245. doi:[10.1016/j.bbi.2018.03.003](https://doi.org/10.1016/j.bbi.2018.03.003)
22. Epel E. How "reversible" is telomeric aging? *Cancer Prev Res (Phila)*. 2012;5(10):1163-1168. doi:[10.1158/1940-6207.CAPR-12-0370](https://doi.org/10.1158/1940-6207.CAPR-12-0370)
23. Steenstrup T, Hjelmberg JV, Kark JD, Christensen K, Aviv A. The telomere lengthening conundrum—artifact or biology? *Nucleic Acids Res*. 2013;41(13):e131-e131. doi:[10.1093/nar/gkt370](https://doi.org/10.1093/nar/gkt370)
24. Bateson M, Nettle D. The telomere lengthening conundrum—it could be biology. *Aging Cell*. 2017;16(2):312-319. doi:[10.1111/acer.12555](https://doi.org/10.1111/acer.12555)
25. Cole JH, Ritchie SJ, Bastin ME, et al. Brain age predicts mortality. *Mol Psychiatry*. 2018;23(5):1385-1392. doi:[10.1038/mp.2017.62](https://doi.org/10.1038/mp.2017.62)

26. Winkler AM, Kochunov P, Blangero J, et al. Cortical thickness or grey matter volume? the importance of selecting the phenotype for imaging genetics studies. *Neuroimage*. 2010;53(3):1135-1146. doi:10.1016/j.neuroimage.2009.12.028
27. Lemaitre H, Goldman AL, Sambataro F, et al. Normal age-related brain morphometric changes: nonuniformity across cortical thickness, surface area and gray matter volume? *Neurobiol Aging*. 2012;33(3):617.e1-617.e9. doi:10.1016/j.neurobiolaging.2010.07.013
28. Hutton C, Draganski B, Ashburner J, Weiskopf N. A comparison between voxel-based cortical thickness and voxel-based morphometry in normal aging. *Neuroimage*. 2009;48(2):371-380. doi:10.1016/j.neuroimage.2009.06.043
29. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract*. 2003;10(2):144-156. doi:10.1093/clipsy.bpg016
30. Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits: a meta-analysis. *J Psychosom Res*. 2004;57(1):35-43. doi:10.1016/S0022-3999(03)00573-7
31. Khoury B, Sharma M, Rush SE, Fournier C. Mindfulness-based stress reduction for healthy individuals: a meta-analysis. *J Psychosom Res*. 2015;78(6):519-528. doi:10.1016/j.jpsychores.2015.03.009
32. Ornish D, Lin J, Chan JM, et al. Effect of comprehensive lifestyle changes on telomerase activity and telomere length in men with biopsy-proven low-risk prostate cancer: 5-year follow-up of a descriptive pilot study. *Lancet Oncol*. 2013;14(11):1112-1120. doi:10.1016/S1470-2045(13)70366-8
33. Lengacher CA, Reich RR, Kip KE, et al. Influence of mindfulness-based stress reduction (MBSR) on telomerase activity in women with breast cancer (BC). *Biol Res Nurs*. 2014;16(4):438-447. doi:10.1177/1099800413519495
34. Tolahunase M, Sagar R, Dada R. Impact of yoga and meditation on cellular aging in apparently healthy individuals: a prospective, open-label single-arm exploratory study. *Oxid Med Cell Longev*. 2017;2017:7928981. doi:10.1155/2017/2784153
35. Rima D, Shiv BK, Bhavna Ch, Shilpa B, Saima Kh. Oxidative stress induced damage to paternal genome and impact of meditation and yoga: can it reduce incidence of childhood cancer? *Asian Pac J Cancer Prev*. 2016;17(9):4517-4525.
36. Wang X, Sundquist K, Hedelius A, Palmér K, Memon AA, Sundquist J. Leukocyte telomere length and depression, anxiety and stress and adjustment disorders in primary health care patients. *BMC Psychiatry*. 2017;17(1):148. doi:10.1186/s12888-017-1308-0
37. Thimmapuram J, Pargament R, Sibliss K, Grim R, Risques R, Toorens E. Effect of heartfulness meditation on burnout, emotional wellness, and telomere length in health care professionals. *J Community Hosp Intern Med Perspect*. 2017;7(1):21-27. doi:10.1080/20009666.2016.1270806
38. Carlson LE, Beattie TL, Giese-Davis J, et al. Mindfulness-based cancer recovery and supportive-expressive therapy maintain telomere length relative to controls in distressed breast cancer survivors. *Cancer*. 2015;121(3):476-484.
39. Duraimani S, Schneider RH, Randall OS, et al. Effects of lifestyle modification on telomerase gene expression in hypertensive patients: a pilot trial of stress reduction and health education programs in African Americans. *PLoS One*. 2015;10(11):e0142689. doi:10.1371/journal.pone.0142689
40. Conklin QA, Crosswell AD, Saron CD, Epel ES. Meditation, stress processes, and telomere biology. *Curr Opin Psychol*. 2018;28:92-101. doi:10.1016/j.copsyc.2018.11.009
41. Singer T, Kok BE, Bornemann B, Zurborg S, Bolz M, Bochow CA. *The ReSource Project: Background, Design, Samples, and Measurements*. 2nd ed. Leipzig, Germany: Max Planck Institute for Human Cognitive and Brain Sciences; 2016.
42. Valk SL, Bernhardt BC, Trautwein FM, et al. Structural plasticity of the social brain: differential change after socio-affective and cognitive mental training. *Sci Adv*. 2017;3(10):e1700489. doi:10.1126/sciadv.1700489
43. Singer T, Kok BE, Bornemann B, Zurborg S, Bolz M, Bochow CA. Design, timeline, and training setting. In: *The ReSource Project: Background, Design, Samples, and Measurements*. 2nd ed. Leipzig, Germany: Max Planck Institute for Human Cognitive and Brain Sciences; 2016:33-35.
44. Singer T, Kok BE, Bornemann B, Zurborg S, Bolz M, Bochow CA. The Resource Protocol. In: *The ReSource Project: Background, Design, Samples, and Measurements*. 2nd ed. Leipzig, Germany: Max Planck Institute for Human Cognitive and Brain Sciences; 2016:25-30.
45. Engert V, Kok BE, Papassotiropoulos I, Chrousos GP, Singer T. Specific reduction in cortisol stress reactivity after social but not attention-based mental training. *Sci Adv*. 2017;3(10):e1700495. doi:10.1126/sciadv.1700495

46. Kok BE, Singer T. Phenomenological fingerprints of four meditations: differential state changes in affect, mind-wandering, meta-cognition, and interoception before and after daily practice across 9 months of training. *Mindfulness (N Y)*. 2017;8(1):218-231. doi:10.1007/s12671-016-0594-9
47. Kok BE, Singer T. Effects of contemplative dyads on engagement and perceived social connectedness over 9 months of mental training: a randomized clinical trial. *JAMA Psychiatry*. 2017;74(2):126-134. doi:10.1001/jamapsychiatry.2016.3360
48. Gibbon M, Spitzer RL, Williams JBW, Benjamin LS. *Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II), User's Guide*. Washington, DC: American Psychiatric Publishing; 1997.
49. First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I), Clinician Version, Administration Booklet*. Washington, DC: American Psychiatric Publishing; 2012.
50. Singer T, Kok BE, Bornemann B, Zurborg S, Bolz M, Bochow CA. Recruitment, sample description, and dropout. In: *The ReSource Project: Background, Design, Samples, and Measurements*. 2nd ed. Leipzig, Germany: Max Planck Institute for Human Cognitive and Brain Sciences; 2016:44-51.
51. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053
52. Lin J, Epel E, Cheon J, et al. Analyses and comparisons of telomerase activity and telomere length in human T and B cells: insights for epidemiology of telomere maintenance. *J Immunol Methods*. 2010;352(1-2):71-80. doi:10.1016/j.jim.2009.09.012
53. FreeSurfer. <http://surfer.nmr.mgh.harvard.edu>. Published 2013. Accessed April 21, 2015.
54. Valk SL, Bernhardt BC, Böckler A, Trautwein F-M, Kanske P, Singer T. Socio-cognitive phenotypes differentially modulate large-scale structural covariance networks. *Cereb Cortex*. 2017;27(2):1358-1368.
55. Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis—I: segmentation and surface reconstruction. *Neuroimage*. 1999;9(2):179-194. doi:10.1006/nimg.1998.0395
56. Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis—II: inflation, flattening, and a surface-based coordinate system. *Neuroimage*. 1999;9(2):195-207. doi:10.1006/nimg.1998.0396
57. Han X, Jovicich J, Salat D, et al. Reliability of MRI-derived measurements of human cerebral cortical thickness: the effects of field strength, scanner upgrade and manufacturer. *Neuroimage*. 2006;32(1):180-194. doi:10.1016/j.neuroimage.2006.02.051
58. Valk SL, Bernhardt BC, Böckler A, Kanske P, Singer T. Substrates of metacognition on perception and metacognition on higher-order cognition relate to different subsystems of the mentalizing network. *Hum Brain Mapp*. 2016;37(10):3388-3399. doi:10.1002/hbm.23247
59. Verhulst S, Aviv A, Benetos A, Berenson GS, Kark JD. Do leukocyte telomere length dynamics depend on baseline telomere length? an analysis that corrects for "regression to the mean". *Eur J Epidemiol*. 2013;28(11):859-866. doi:10.1007/s10654-013-9845-4
60. Bateson M, Eisenberg DTA, Nettle D. Controlling for baseline telomere length biases estimates of the rate of telomere attrition (preprint, version 2). Zenodo (preprint, version 2). <https://zenodo.org/record/2458376#.X5fsI3dFxPY>. Published December 20, 2018. Accessed February 18, 2019.
61. Pinheiro JC, Bates DM. *Mixed-Effects Models in S and S-PLUS*. New York, NY: Springer; 2000. doi:10.1007/978-1-4419-0318-1
62. Gardner M, Bann D, Wiley L, et al; Halcyon study team. Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol*. 2014;51(1):15-27. doi:10.1016/j.exger.2013.12.004
63. Mirabello L, Huang W-Y, Wong JYY, et al. The association between leukocyte telomere length and cigarette smoking, dietary and physical variables, and risk of prostate cancer. *Aging Cell*. 2009;8(4):405-413. doi:10.1111/j.1474-9726.2009.00485.x
64. Salat DH, Buckner RL, Snyder AZ, et al. Thinning of the cerebral cortex in aging. *Cereb Cortex*. 2004;14(7):721-730. doi:10.1093/cercor/bbh032
65. Worsley K, Taylor J, Carbonell F, et al. SurfStat: a Matlab toolbox for the statistical analysis of univariate and multivariate surface and volumetric data using linear mixed effects models and random field theory. *Neuroimage*. 2009;47:S102. doi:10.1016/S1053-8119(09)70882-1
66. Greve DN, Fischl B. False positive rates in surface-based anatomical analysis. *Neuroimage*. 2018;171(171):6-14. doi:10.1016/j.neuroimage.2017.12.072
67. Rosnow RL, Rosenthal R, Rubin DB. Contrasts and correlations in effect-size estimation. *Psychol Sci*. 2000;11(6):446-453. doi:10.1111/1467-9280.00287

68. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw*. 2015;67(1):1-48. doi:10.18637/jss.v067.i01
69. R Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. 2018. <https://www.R-project.org>. Accessed August 21, 2019.
70. Dobson AJ. *An Introduction to Generalized Linear Models*. Boca Raton, FL: Chapman & Hall/CRC; 2002.
71. Singer T, Kok BE, Bornemann B, Zurborg S, Bolz M, Bochow CA. Participants' descriptives at baseline (T0). In: *The ReSource Project: Background, Design, Samples, and Measurements*. 2nd ed. Leipzig, Germany: Max Planck Institute for Human Cognitive and Brain Sciences; 2016:112.
72. Staffaroni AM, Tosun D, Lin J, et al. Telomere attrition is associated with declines in medial temporal lobe volume and white matter microstructure in functionally independent older adults. *Neurobiol Aging*. 2018;69:68-75. doi:10.1016/j.neurobiolaging.2018.04.021
73. Svenson U, Nordfjäll K, Baird D, et al. Blood cell telomere length is a dynamic feature. *PLoS One*. 2011;6(6):e21485. doi:10.1371/journal.pone.0021485
74. Blackburn EH, Epel ES, Lin J. Human telomere biology: A contributory and interactive factor in aging, disease risks, and protection. *Science*. 2015;350(6265):1193-1198. doi:10.1126/science.aab3389
75. Melk A, Tegtbur U, Hilfiker-Kleiner D, et al. Improvement of biological age by physical activity. *Int J Cardiol*. 2014;176(3):1187-1189. doi:10.1016/j.ijcard.2014.07.236
76. Wolkowitz OM, Mellon SH, Lindqvist D, et al. PBMC telomerase activity, but not leukocyte telomere length, correlates with hippocampal volume in major depression. *Psychiatry Res*. 2015;232(1):58-64. doi:10.1016/j.psychres.2015.01.007
77. Epel ES, Blackburn EH, Lin J, et al. Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci U S A*. 2004;101(49):17312-17315. doi:10.1073/pnas.0407162101
78. McEwen BS. The brain is the central organ of stress and adaptation. *Neuroimage*. 2009;47(3):911-913. doi:10.1016/j.neuroimage.2009.05.071
79. Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci*. 2002;25(6):295-301. doi:10.1016/S0166-2236(02)02143-4
80. Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*. 2006;129(pt 3):564-583. doi:10.1093/brain/awl004
81. Palmqvist S, Schöll M, Strandberg O, et al. Earliest accumulation of β -amyloid occurs within the default-mode network and concurrently affects brain connectivity. *Nat Commun*. 2017;8(1):1214.
82. Nilsson G, Tamm S, Månsson KNT, Åkerstedt T, Lekander M. Leukocyte telomere length and hippocampus volume: a meta-analysis. *FlourRes*. 2015;4:1073. doi:10.12688/flourresearch.7198.1

SUPPLEMENT 1.

Trial Protocol

SUPPLEMENT 2.

eAppendix 1. Detailed Methodology

eAppendix 2. Descriptive Analyses

eAppendix 3. Random Slopes Model

eTable 1. Correlations Between Change in LTL and Leukocytic Cell Type Concentrations in the RCC

eAppendix 4. Exploratory Analyses

eAppendix 5. Residual Checks

eFigure 1. Change in Telomere Length and Cortical Thickness From Baseline to 9-Month Follow-up

eFigure 2. Relative Difference in Telomere Length per Training Module

eFigure 3. DLTL and Δ CT in Right Precuneus/PCC ROI

eTable 2. Mean (SD) of Baseline Personality Trait Scores

eTable 3. Results of Exploratory Analyses

eTable 4. Mean (SD) Unmodelled Change in LTL After Each Training Module

eTable 5. Available Samples and Reasons for Missing Values in the Raw Data

eTable 6. Available Samples and Reasons for Missing Values in Change Scores

eReferences

SUPPLEMENT 3.

Data Sharing Statement