

Oxytocin has ‘Tend-and-Defend’ Functionality in Group Conflict Across Social Vertebrates

Zegni Triki^{1,*}, Katie Daughters², and Carsten K.W. De Dreu^{3,4}

¹Department of Zoology, Stockholm University

²Department of Psychology, Essex University, United Kingdom

³Institute of Psychology, Leiden University, The Netherlands

⁴Center for Research in Experimental Economics and Political Decision Making, University of Amsterdam, The Netherlands

*Corresponding author at: zegni.triki@gmail.com

Abstract

Across vertebrate species, intergroup conflict confronts individuals with a tension between group interests best served by participation in conflict and personal interest best served by not participating. Here, we identify the neurohormone oxytocin as pivotal to the neurobiological regulation of this tension in distinctly different group-living vertebrates, including fish, birds, rodents, non-human primates, and humans. In the context of intergroup conflict, a review of emerging work on pro-sociality suggests that oxytocin and its fish and birds homologs, isotocin and mesotocin, respectively, can elicit participation in group conflict and aggression. This is because it amplifies (i) concern for the interests of genetically related or culturally similar ‘in-group’ others, and (ii) willingness to defend against outside intruders and enemy conspecifics. Across a range of social vertebrates, oxytocin can induce aggressive behaviour to ‘tend-and-defend’ the in-group during intergroup contests.

Keywords: Parochial Altruism; In-group; Out-group; Neuromodulation; Decision-making; Vertebrates

1. Introduction

Interactions between groups of conspecifics can be cooperative and benign but also hostile, for example, when groups compete for (access to) food, mating opportunities and territory [1,2]. Moreover, across species and all else equal, groups are more likely to be victorious when their members contribute to the collective aggression of rivalling other groups and prevent defeat when they contribute to the collective defence against enemy attacks [3,4]. And yet, joining conflict requires investing personal resources and increases the risk of injury. Participating in out-group aggression and in-group defence thus requires individuals to solve a tension between personal interests on the one hand and group interests on the other [4–7].


The tension between personal interests, served by withholding participating in conflict and group interests served by pro-actively contributing, is seen in several species across taxa [1,7]. Perhaps there are evolutionary preserved biological mechanisms that regulate individual participation in intergroup conflict. Here, we examine this possibility at the neurobiological level by focusing on the role of oxytocin (and its homologs isotocin and mesotocin [8,9]) in regulating key parameters underlying conflict participation. We uncover a remarkable cross-species commonality in how isotocin in social fish, mesotocin in gregarious birds, and oxytocin in group-living mammals biologically prepares for a ‘tend-and-defend’ response during intergroup conflict and not, or less so, for the aggressive subordination and exploitation of rivalling groups of conspecifics.

2. Evolution and Neurobiology of Oxytocin

Oxytocin is a nine-amino acid peptide (i.e., nonapeptide) synthesised primarily in the brain. It can act centrally as a neuromodulator and/or peripherally as a hormone [9]. Across taxa and species, the mammalian oxytocin has several homologs, such as ‘isotocin’ in bony fishes [10], ‘mesotocin’ in nonmammalian tetrapods (lungfish, amphibians, reptiles and birds) [11] and up to five structural variants of oxytocin recently sequenced in new world monkeys [12] (Fig. 1).

Oxytocin can thus be viewed as an ancient peptide widely preserved across taxa. It shares common ancestors with another nonapeptide, ‘vasopressin’, that can be traced back all the way to snails and insects (Fig. 1). However, not all insects have this nonapeptide gene ancestor, such as silkworms, fruit flies, mosquitos, spiders, and honeybees, suggesting a potential loss of such genes [13]. The vertebrates witnessed the emergence of oxytocin and its sister nonapeptide vasopressin (and their homologs) about 500 million years ago through gene duplication of a common ancestral gene, presumably in jawless fish [8]. Since then, the oxytocin structure has remained highly preserved, from bony fishes to mammals, where the structural differences between oxytocin and its homologs and the recently discovered mammalian variants occur in one or two amino acids (Fig. 1). These differences notwithstanding, oxytocin variants and homologs thus share important structural and functional elements. For readability, we use here the nomenclature ‘oxytocin’ for all mammalian oxytocins, isotocin and mesotocin [14].

Figure 1. Oxytocin amino acids sequences across taxa/species. A simplified phylogenetic tree shows oxytocin sequence in different species and taxa with the common ancestors of oxytocin and vasopressin in invertebrates. To differentiate oxytocins in mammals, the variant amino acid and its position are indicated as a prefix. The Leu⁸-Oxytocin is taken here as a reference to see which amino acid(s) differ from this structure, where non-matching amino acids are colour coded. NWM refers to new world monkeys (e.g., marmosets, spider monkeys, capuchin monkeys, etc.). Nonapeptides sequences are from [8,12]. Illustrations by Z. Triki.



In all vertebrates, oxytocin is synthesised mainly in the magnocellular and parvocellular hypothalamic neurons. From here, oxytocin can be released centrally or relayed to the posterior pituitary gland, where oxytocin is released into the bloodstream and eventually cleared out in other fluids such as saliva and urine (Side Box 1). In teleost and amphibians, the hypothalamic parvocellular and magnocellular neurons are located in the preoptic area and anterior hypothalamus. In other vertebrates, such as reptiles, birds and mammals, rather two separate nuclei, the paraventricular and supraoptic nuclei, harbour the oxytocin neurons (for further details, see [15]).

Upon its release from neuronal soma, axons, and dendrites, oxytocin exerts widespread effects in the brain via oxytocin specific G protein-coupled receptor [9]. Oxytocin binding on this receptor activates a set of signalling cascades that can quickly modulate the evolutionary ancient and structurally and functionally preserved social decision-making network in the vertebrate brain [16,17]. This network includes various brain nuclei known for their crucial roles in regulating social recognition, affiliation and parental behaviour, responses to social stressors, and aggression [2,17–19].

Side Box 1: Measuring and Manipulating Oxytocin in Social Vertebrates

The box summarises the most common techniques used in biology to manipulate and measure oxytocin levels across taxa. To study causal effects, several methods exist to manipulate oxytocin (left panel). Non-invasive intranasal administration of oxytocin or an antagonist is commonly used in (non-)human primates. Invasive techniques include injections and intracerebroventricular infusions and are more commonly used in small mammals, fish, and birds. Finally, causality is studied by comparing models with versus without in-tact oxytocin circuitry (i.e., knockdown (out) models). To examine correlations between naturally occurring oxytocin release and behaviour, central oxytocin is obtained from cerebrospinal fluid or directly from the brain, and peripheral oxytocin can be obtained from blood plasma, saliva, or urine (top right panel). Various assaying techniques exist to detect the presence of the nonapeptide in the sample (bottom right panel). For further details on the different techniques' accuracy and validity and the extent to which endogenous/exogenous oxytocin levels can be an informative tool in behavioural studies, please see [20–27].

Manipulating (Oxytocin Agonist/Antagonist)				Measuring			
Non-Invasive	<i>Nasal spray</i>	<i>Nebuliser</i>	<i>Nasal drops</i>	Central	<i>Cerebrospinal Fluid</i>	<i>Microdialysis</i>	<i>Brain Harvesting</i>
	Primates [28]	Primates [29]	Rodents [30] Birds [31]		Primates [32] Rodents [33]	Rodents [30]	Rodents [34] Birds [35] Fish [36]
Invasive	<i>Intravenous/ intra-Muscular/ intraperitoneal</i>	<i>Intracerebroventricular Infusion</i>		Peripheral	<i>Blood Plasma</i>	<i>Saliva</i>	<i>Urine</i>
	Primates [37] Rodents [30] Birds [35] Fish [38]	Primates [39] Rodents [40] Birds [41] Fish [42]			Primates [32] Carnivores [43] Rodents [30] Birds [44] Fish [45]	Primates [28] Birds [46]	Primates [47] Carnivores [48]
Genetic	<i>Knockout Models</i>	<i>Knock-down Models</i>	<i>Behavioural Phenotypes (selection experiments)</i>	Assay			<ul style="list-style-type: none"> • Enzyme-linked immunosorbent assay (ELISA) [28] • Radioimmunoassay (RIA) [24] • Liquid Chromatography-Mass Spectrometry (LC-MS) [52] • Immunohistochemistry (IHC) [53] • mRNA Quantification [54]
	Rodents [49]	Rodents [34] Birds [50]	Rodents [51]				

3. Oxytocin and Participation in Group Conflict


There is growing evidence that in a range of species, oxytocin plays a significant role in forming and solidifying social structures (e.g., [20] [Lemaine et al., this issue]). In particular, affiliation among conspecifics is often associated with higher oxytocin levels. For example, studies that use oxytocin levels from blood plasma, urine, or saliva as an informative tool on central oxytocin release, have recorded elevated oxytocin following affiliative touch [55–57] and cooperative exchange [58–60] in mammalian species such as (human) primates and dogs. Also, strongly bonded marmoset monkeys showed synchronised fluctuations of oxytocin over a six week period [61] (also see [62]). Similar positive effects of affiliation on oxytocin levels are found in gregarious birds [63], lizards [64], and fish [36]. Other work observed links between oxytocin levels in distinct brain regions on the one hand, and a range of social behaviours on the other, including suckling (in rats and sheep [65–67] and mating (e.g., in voles [68]).

At first blush, the mutually reinforcing relationship between affiliation and oxytocin may appear antagonistic to the possibility that oxytocin prepares individuals for participation in hostile group conflict with conspecifics. However, for group conflict to be won, or not lost, individuals within rivaling groups need to contribute to their group's fighting capacity at some personal cost (Fig. 2) (also see [2,5–7,69,70]). Making such costly contributions serves the group and can thus be seen as a form of pro-social behaviour towards one's in-group. Indeed, as we [2,71] and others (e.g., [1,5,7]) have argued and shown, in many group-living species an individual's conflict participation p_i is a function of a concern for in-group (henceforth α_i) and out-group interests (henceforth α_o), expected out-group threat (henceforth β), and compliance with group norms for participation (*viz.* reputation concerns; henceforth γ) (see also [De Dreu & Triki, this issue]). If we set each parameter to vary between -1 and 1 inclusive, participation likelihood increases when there is a positive concern for in-group interests ($\alpha_i > 0$), negative concern for out-group interests ($\alpha_o < 0$), perceived out-group threat ($\beta > 0$) or when the animal expects participation returns reputation benefits ($\gamma > 0$) [1,2,4] [De Dreu and Triki, this issue]. This means that participation can be expected when and because oxytocin increases (i) in-group concern α_i , and/or (ii) creates negative out-group concern ($\alpha_o < 0$), and/or (iii) increases perceived out-group threat β , and/or (iv) increases expectation of reciprocity and reputation benefits from participating (γ). In the remainder of this section, we examine the evidence for the role of oxytocin on each of these parameters underlying participation in conflict (also see [67,72–75]) (Side Box 2).

Figure 2 Intergroup Conflict as a Multilevel Contest Game of Strategy

(A) Individuals nested in two groups (Circle: $a_i...a_n$ and Square: $b_i...b_n$) can contribute personal resources (e.g., skills, time and energy) to their group's capacity for out-group attack A; red) and/or to protect against enemy attacks (in-group defence D; blue). Conflict participation is risky – the individual may get

injured – and resources contributed are ‘wasted.’ **(B)** Conflict participation increases the likelihood of victory with concomitant ‘spoils of war’ (a win/lose outcome; top panel), and of surviving out-group attacks (a stalemate outcome; bottom panel). Because (participating in) conflict is wasteful, even winning groups typically are less wealthy post-conflict.



3.1 Parochial Preferences (in-group interest $\alpha_I >$ out-group interest α_O)

Studies with human participants revealed that concerns for genetically related or culturally similar conspecifics (in-group) are typically stronger than for unrelated and unfamiliar (out-group) conspecifics [76] (Side Box 2). Oxytocin has a mechanistic role to play in such in-group biased preferences (i.e., $\alpha_I > \alpha_O$) [77,78]. For instance, in-group participants in a foraging game helped each other more often compared to out-group participants, a behaviour that was mediated by endogenous oxytocin (i.e., measured in saliva) [58]. Similarly, Chinese males had a frontocentral positive activity of larger amplitude in response to the pain expressions of in-group (Asian targets) but not out-group members (Caucasian targets), especially following intranasal administration of oxytocin rather than placebo [79] (also [80]).

Side Box 2: Inferring Conflict Participation Parameters from Vertebrate Decision-Making

Concern for in-group (α_I) and out-group (α_O) can be inferred from behavioural choices, neural activation in, e.g., mesolimbic reward circuitry and, in humans, self-reports. In humans, social concerns can be inferred from economic decision-making games such as the Dictator Game (DG), wherein participants donate x out of an endowment e to an anonymous recipient (with $0 \leq x \leq e$). Higher donations to in-group rather than out-group members reflect stronger concern for in-group (α_I) than out-group interests (α_O) [76,81]. Variants of such games have been used to infer social preferences in non-human primates [82]. In nonmammalian

vertebrates, such as social fish, social preferences are inferred from time spent in proximity of a conspecific [38] or from costly helping of a conspecific [83]. To infer *expectations of reciprocity* (viz. β_T), studies with humans used Trust Games. Participants can transfer x out of an endowment e to a recipient (with $0 \leq x \leq e$). The recipient then receives $3x$ and can return y to the participant (with $0 \leq y \leq 3x$). Greater transfers reflect expectations of reciprocity (or ‘trust’), and greater back-transfers reflect a willingness to reciprocate (or ‘trustworthiness’) [81]. Vice versa, expectations of competition can be inferred from partner choice, with rejecting partners who did not cooperate on earlier occasions as a measure of negative expectations (in humans[81]; in birds [84]; in fish [85]). Finally, *reputation concerns* have been inferred from third-party punishment games [82], where participants, after decision-making, express through punishment social disapproval of the others’ (non-cooperative) behaviour and/or induce a norm for cooperation on future trials [86]. Punishment and behavioural adjustments to (threat of) punishment are seen across social vertebrates, including chimpanzees [87] and social fish [88].

In humans, oxytocin seems to amplify α_I and neither increases nor decreases α_O – oxytocin makes humans like their in-group more and does not condition (dis)liking out-groups. This was shown, for example, when human participants indicated their liking for individuals from their own nationality (i.e., Dutch citizens) and individuals from a more or less rivalling nationality (e.g., Germans). Compared to placebo-treated individuals, those given intranasal oxytocin expressed a greater liking for in-group members (an increase in α_I) but did not increase or decrease their liking for out-group members (i.e., α_O was similar in oxytocin and placebo conditions) [77]. Recent work on wild chimpanzees suggests that these effects may generalize to other species, including voles [89,90], sheep [91] and chimpanzees [92]). In another series of experiments with human participants, individuals were organised in two groups of three and could contribute to club goods A and B out of a personal endowment. Whereas contributions to A and B equally benefitted the members of one’s own group, contributions to B (but not A) also imposed a cost on the out-group members. Intranasal oxytocin (versus placebo) increased contributions to club good A, reflecting an increase in α_I . However, oxytocin neither increased nor decreased contributions to club good B, suggesting oxytocin did not affect α_O [93,94] (also see [95,96]).

Although follow-up experiments in humans sometimes show that oxytocin can increase α_O (e.g., [97,98]), this effect is rarely as strong as the oxytocin-induced increase on α_I . This mirrors findings with non-human vertebrates. For example, marmosets treated with marmoset-specific Pro⁸-Oxytocin reduced pro-sociality toward strangers compared to those treated with saline or consensus-mammalian Leu⁸-Oxytocin [99] (also [100]). Chimpanzees had higher urinary oxytocin concentrations before and after hostile intergroup encounters, which predicted within-group affiliative behaviours [47]. Resident male mice exhibit higher attack bites against intruders of different strains (viz. out-group) than against intruders of their own strain. Yet compared to oxytocin receptor wild-type mice, oxytocin receptor-null residents exhibited

greater aggression toward intruders of their own strain, suggesting that oxytocin modulates α_I more than α_O [49]. In a monogamous zebra finch, affiliation towards one's partner requires the activation of oxytocin receptor [41], while oxytocin knockdown birds and those treated with an oxytocin antagonist experienced affiliation behaviour deficit [50,63] (for similar findings in pinyon jays, see [30]). Finally, work on the mutualistic cleaner fish and its various coral reef fish clients showed that cleaners injected with oxytocin break less often the already engaged cleaner-client social interaction to initiate a new interaction with a newly arrived client [101].

Together, there is growing evidence for the possibility that across social vertebrates, oxytocin appears to increase a positive concern for the interests of familiar conspecifics more than for the interests of genetically or culturally unfamiliar, out-group conspecifics: $\alpha_I > \alpha_O$. At least in humans, this parochial preference is also reflected in in-group-biased expectations of reciprocity derived from trust games (Side Box 1; [72,93]. In short, when individuals with elevated levels of oxytocin participate in conflict this is more likely due to an increase in α_I than because of a decrease in α_O .

3.2 Responding to Out-group Threat (β)

Nursing rats protect their offspring against intruders by aggressing them with fast attacks directed toward the intruder's neck or back region, lateral threats to force the intruder aside, and standing in an upright posture in front of the intruder, sometimes using the front legs to hold the intruder down [102]. Such 'maternal defence' rests on oxytocin, where oxytocin knockout rats and those treated with oxytocin antagonists abstain from aggressing intruders [19,102,103].

A suite of follow-up studies shows oxytocin-mediated aggression towards threatening outsiders is not confined to (female) rodents. For example, when groups of wild meerkats were given intravenous oxytocin (or placebo), individuals spent over twice as much time 'on guard'; a personally costly behaviour that helps to protect the group against an outside threat from predators and hostile conspecifics [104] (see [100] for similar results in marmoset monkeys). Likewise, estrildid finches that form year-round male-female pairs aggressively defend their territories from intruders. Yet, such aggressive defence is significantly reduced following the blockade of oxytocin receptors in the avian brain [35]. And in social fish such as cichlids and sticklebacks, the presence of an intruder incites higher oxytocin neuronal activity [53] (also see [105]), and higher brain oxytocin levels associate with an aggressive defence of nest and territory [36].

Experiments with human participants confirmed that oxytocin could elicit defensive aggression and suggest that such aggression is closely tied to rivalling out-group threats. For example, several studies showed that oxytocin increases competition against out-group members if, and only if, out-group hostility would hurt the individual and/or its in-group members [93,106] (also see [74]). Other studies using different experimental tasks produced similar results. For example, individuals given oxytocin more quickly (and less accurately) aggressed ethnically different rather than ethnically similar intruders [107].

Taken together, there is converging evidence across social vertebrates that oxytocin up-

regulates attention and aggressive responses toward predators and rivaling conspecifics. In addition to parochial preferences (Section 3.1), individuals with elevated levels of oxytocin may increase their conflict participation because of enhanced perception of out-group threat and increased readiness to protect and defend genetically related and culturally familiar conspecifics [108].

3.3 Reputation and Group Norms for Participation (γ)

Individuals within groups adapt behaviour to other group members' choices, including those of 'first-movers' and group leaders [Glowacki and McDermott, this issue; Smith, Kappeler et al., this issue]. Such behavioural alignment or 'compliance' enables the individual to benefit from the protection offered by the group and, in addition, facilitates the coordination of collective action towards some group goal [81,109]]. Behavioural alignment thus is functional toward both individual and group survival and prosperity both in general and in the context of intergroup conflict. Furthermore, groups are more likely to win intergroup contests when individual contributions are well-coordinated and aligned with leader initiatives [78,96,110] (also see [De Dreu & Triki, this issue]).

There is some evidence that oxytocin facilitates behavioural alignment and compliance with group norms. Humans, for instance, change their private views in the direction of their group members' opinions more when given oxytocin rather than placebo [111–114]. Likewise, oxytocin mediates interpersonal synchronisation at both the neural and behavioural levels in humans [115–119], marmoset monkeys [61], dogs [120], and social fish [121]. In one study with humans, individuals within groups aligned their contributions to group conflict better when given oxytocin than placebo. As a result, their groups won greater 'spoils of war' [96]. Oxytocin may, therefore, prepare the individual for conflict participation because it increases sensitivity to and compliance with leader initiatives and group norms for participation.

Because compliance can have adaptive functionality to the group, individuals are willing to enforce compliance in other group members [81,86]. For example, humans punish those who fail to contribute to group conflict, and such (threat of) punishment increases subsequent conflict participation [110]. At least in humans, there is some evidence that oxytocin prepares the individual for such norm enforcement. For instance, in one study, participants as neutral third parties punished group members who had exploited another person's trust more when given oxytocin rather than placebo [122] (also see [123–126]). In short, oxytocin facilitates interpersonal synchronisation and alignment across various social vertebrates at the neural, physiological, and behavioural levels. Possibly, and especially when collaborations require strong synchrony in space and time [127], individuals with elevated oxytocin may participate in group conflict because of amplified γ – the readiness to align with and follow other group members' initiatives.



4. Conclusions

Our review reveals converging evidence for the possibility that oxytocin has a 'tend-and-defend' functionality that prepares for active conflict participation through an increase in parochial in-group preferences (α) and perceived threat from out-groups (β). We observed little

to no evidence that oxytocin modulates (negative) concern for out-groups (α_0) and concomitant aggression aimed at exploiting and sub-ordinating outsiders (Fig. 3).

Figure 3 Oxytocin and conflict participation parameters across social vertebrates

Oxytocin creates parochial preferences ($\alpha_I > \alpha_0$) because it up-regulates α_I (concern for genetically and culturally related conspecifics) and less α_0 (concern for genetically and culturally unrelated conspecifics). Oxytocin also up-regulates β (the willingness to defend aggressively against intruders and groups of rivaling conspecifics). At least in primates, oxytocin increases γ (behavioural alignment with group norms for participation). Arrows indicate the direction of an effect of, or association with oxytocin. Empty cells indicate no or too little evidence is available. Illustrations by C. De Dreu

Social Vertebrate	$\alpha_I > \alpha_0$	β	γ	P_{part}
	↑	↑	↑	Primates > 0
	↑	↑		Non-Primate Mammals > 0
	↑	↑		Birds > 0
	↑	↑	↑	Fish > 0

Our conclusion comes with some limitations. First, we allowed for some degrees of freedom in interpreting animal behaviour as reflective of social preferences (α), threat-responding (β), and norm compliance (γ). Such ‘heuristic’ treatment ignores that both animal behaviour and hormones are often equifinal – different behaviours or hormones serving the same function – and multi-final – the same behaviour or hormone serving several functions [128–130]. Future experiments could try to isolate these parameters further and, in addition, examine possible interactions (e.g., social preferences up-regulate threat-responding). Second, not all parameters

in the conflict participation function have been covered across social vertebrates, and there are a range of context-dependencies that can complicate straightforward predictions. Conclusive evidence for oxytocin-induced reputation concerns and compliance with group norms, for example, appears limited to humans (Fig. 3). Third, our analysis collapsed across various measurements and manipulations of oxytocin, and some evidence is strictly correlational. For example, research with humans mostly relied on up-regulating oxytocin and has not examined how oxytocin antagonists reduce conflict participation. Also, research often either considered only females or males, while some effects might be sex-specific.

The converging evidence for ‘tend-and-defend’ functionality across social vertebrates should not be taken as if oxytocin is required for participation in group conflict to emerge. Some highly social species such as bees engage in lethal intergroup conflict [131] yet lack oxytocin homologs. Whereas social vertebrates may have co-opted the oxytocinergic circuitry to support a ‘tend-and-defend’ response during the intergroup conflict, other species may rely on different neuroendocrine systems to produce strategic engagement in intergroup conflict. In addition, in social vertebrates, other neurohormonal mechanisms may contribute to conflict participation. For example, oxytocin and vasopressin co-evolved, where vasopressin differs in two amino acids compared to oxytocin [14]. Yet, like oxytocin, vasopressin regulates affiliative behaviour and context-dependent aggressive behaviour (e.g., competition, territory defence) [38,132]. Furthermore, the sex steroid testosterone mediates aggressive behaviour, which can influence group conflict outcomes [133], and the stress hormone cortisol mediates the natural “fight-or-flight” response to threatening conspecifics [47]. Future work into the neurohormonal underpinnings of conflict participation is needed, particularly in how distinctly different neurotransmitters and hormones interact in producing prosocial behaviour towards genetically related and culturally similar conspecifics and aggression towards more or less rivaling outgroups.

5. Author Acknowledgments

This project has received funding from the Swiss National Science Foundation (grant n° P2NEP3_188240 and P400PB_199286) to Z.T. and from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (AdG agreement n° 785635) to C.K.W.D.D. Authors contributed equally to all aspects of the project. No conflict of interest was declared.

6. References

1. Radford AN, Majolo B, Aureli F. 2016 Within-group behavioural consequences of between-group conflict: a prospective review. *Proc. R. Soc. B Biol. Sci.* **283**, 20161567. (doi:10.1098/rspb.2016.1567)
2. De Dreu CKW, Gross J, Fariña A, Ma Y. 2020 Group cooperation, carrying-capacity stress, and intergroup conflict. *Trends Cogn. Sci.*
3. Wilson DS, Wilson EO. 2007 Rethinking the Theoretical Foundation of Sociobiology. *Q. Rev. Biol.* **82**, 327–348. (doi:10.1086/522809)

4. De Dreu CKW, Gross J. 2019 Revisiting the form and function of conflict: Neurobiological, psychological, and cultural mechanisms for attack and defense within and between groups. *Behav. Brain Sci.* **42**. (doi:10.1017/S0140525X18002170)
5. Bornstein G. 2003 Intergroup conflict: individual, group, and collective interests. *Personal. Soc. Psychol. Rev. Off. J. Soc. Personal. Soc. Psychol. Inc* **7**, 129–145. (doi:10.1207/S15327957PSPR0702_129-145)
6. Choi J-K, Bowles S. 2007 The Coevolution of Parochial Altruism and War. *Science* **318**, 636–640. (doi:10.1126/science.1144237)
7. Rusch H, Gavrillets S. 2020 The logic of animal intergroup conflict: A review. *J. Econ. Behav. Organ.* **178**, 1014–1030. (doi:10.1016/j.jebo.2017.05.004)
8. Stoop R. 2012 Neuromodulation by Oxytocin and Vasopressin. *Neuron* **76**, 142–159. (doi:10.1016/j.neuron.2012.09.025)
9. Jurek B, Neumann ID. 2018 The Oxytocin Receptor: From Intracellular Signaling to Behavior. *Physiol. Rev.* **98**, 1805–1908. (doi:10.1152/physrev.00031.2017)
10. Acher R, Chauvet J, Chauvet M, Crepy D. 1968 Molecular evolution of neurohypophysial hormones: Comparison of the active principles of three bony fishes. *Gen. Comp. Endocrinol.* **11**, 535–538.
11. Acher R, Chauvet J, Chauvet M. 1970 Phylogeny of the neurohypophysial hormones: The avian active peptides. *Eur. J. Biochem.* **17**, 509–513.
12. Ren D, Lu G, Moriyama H, Mustoe AC, Harrison EB, French JA. 2015 Genetic Diversity in Oxytocin Ligands and Receptors in New World Monkeys. *PLOS ONE* **10**, e0125775. (doi:10.1371/journal.pone.0125775)
13. Stafflinger E, Hansen KK, Hauser F, Schneider M, Cazzamali G, Williamson M, Grimmlikhuijzen CJP. 2008 Cloning and identification of an oxytocin/vasopressin-like receptor and its ligand from insects. *Proc. Natl. Acad. Sci.* **105**, 3262–3267. (doi:10.1073/pnas.0710897105)
14. Theofanopoulou C, Gedman G, Cahill JA, Boeckx C, Jarvis ED. 2021 Universal nomenclature for oxytocin–vasotocin ligand and receptor families. *Nature* **592**, 747–755. (doi:10.1038/s41586-020-03040-7)
15. Knobloch HS, Grinevich V. 2014 Evolution of oxytocin pathways in the brain of vertebrates. *Front. Behav. Neurosci.* **8**. (doi:10.3389/fnbeh.2014.00031)
16. O’Connell, Hofmann HA. 2011 The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *J. Comp. Neurol.* **519**, 3599–3639. (doi:10.1002/cne.22735)
17. Rilling JK, Young LJ. 2014 The biology of mammalian parenting and its effect on offspring social development. *Science* **345**, 771–776.

18. Goodson JL. 2005 The vertebrate social behavior network: Evolutionary themes and variations. *Horm. Behav.* **48**, 11–22. (doi:10.1016/j.yhbeh.2005.02.003)
19. de Moura Oliveira VE, Lukas M, Wolf HN, Durante E, Lorenz A, Mayer A-L, Bludau A, Bosch OJ, Grinevich V, Egger V. 2021 Oxytocin and vasopressin within the ventral and dorsal lateral septum modulate aggression in female rats. *Nat. Commun.* **12**, 1–15.
20. Crockford C, Deschner T, Ziegler T, Wittig R. 2014 Endogenous peripheral oxytocin measures can give insight into the dynamics of social relationships: a review. *Front. Behav. Neurosci.* **8**, 68. (doi:10.3389/fnbeh.2014.00068)
21. McCullough ME, Churchland PS, Mendez AJ. 2013 Problems with measuring peripheral oxytocin: Can the data on oxytocin and human behavior be trusted? *Neurosci. Biobehav. Rev.* **37**, 1485–1492. (doi:10.1016/j.neubiorev.2013.04.018)
22. Graustella AJ, MacLeod C. 2012 A critical review of the influence of oxytocin nasal spray on social cognition in humans: evidence and future directions. *Horm. Behav.* **61**, 410–418.
23. Engel S, Laufer S, Miller R, Niemeyer H, Knaevelsrud C, Schumacher S. 2019 Demographic, sampling-and assay-related confounders of endogenous oxytocin concentrations: A systematic review and meta-analysis. *Front. Neuroendocrinol.* **54**, 100775.
24. Szeto A, McCabe PM, Nation DA, Tabak BA, Rossetti MA, McCullough ME, Schneiderman N, Mendez AJ. 2011 Evaluation of Enzyme Immunoassay and Radioimmunoassay Methods for the Measurement of Plasma Oxytocin: *Psychosom. Med.* **73**, 393–400. (doi:10.1097/PSY.0b013e31821df0c2)
25. Valstad M, Alvares GA, Andreassen OA, Westlye LT, Quintana DS. 2016 The relationship between central and peripheral oxytocin concentrations: a systematic review and meta-analysis protocol. *Syst. Rev.* **5**, 1–7.
26. Churchland PS, Winkielman P. 2012 Modulating social behavior with oxytocin: how does it work? What does it mean? *Horm. Behav.* **61**, 392–399.
27. Evans SL, Dal Monte O, Noble P, Aeverbeck BB. 2014 Intranasal oxytocin effects on social cognition: a critique. *Brain Res.* **1580**, 69–77.
28. Daughters K, Manstead ASR, Hubble K, Rees A, Thapar A, Goozen SHM van. 2015 Salivary Oxytocin Concentrations in Males following Intranasal Administration of Oxytocin: A Double-Blind, Cross-Over Study. *PLOS ONE* **10**, e0145104. (doi:10.1371/journal.pone.0145104)
29. Martins DA, Mazibuko N, Zelaya F, Vasilakopoulou S, Loveridge J, Oates A, Maltezos S, Mehta M, Wastling S, Howard M, McAlonan G, Murphy D, Williams SCR, Fotopoulou A, Schuschnig U, Paloyelis Y. 2020 Effects of route of administration on oxytocin-induced changes in regional cerebral blood flow in humans. *Nat. Commun.* **11**, 1160. (doi:10.1038/s41467-020-14845-5)

30. Neumann ID, Maloumy R, Beiderbeck DI, Lukas M, Landgraf R. 2013 Increased brain and plasma oxytocin after nasal and peripheral administration in rats and mice. *Psychoneuroendocrinology* **38**, 1985–1993. (doi:10.1016/j.psyneuen.2013.03.003)
31. Duque JF, Leichner W, Ahmann H, Stevens JR. 2018 Mesotocin influences pinyon jay prosociality. *Biol. Lett.* **14**, 20180105. (doi:10.1098/rsbl.2018.0105)
32. Striepens N, Kendrick KM, Hanking V, Landgraf R, Wüllner U, Maier W, Hurlemann R. 2013 Elevated cerebrospinal fluid and blood concentrations of oxytocin following its intranasal administration in humans. *Sci. Rep.* **3**, 3440. (doi:10.1038/srep03440)
33. Dogterom J, Greidanus TBVW, Swaab DF. 1977 Evidence for the Release of Vasopressin and Oxytocin into Cerebrospinal Fluid: Measurements in Plasma and CSF of Intact and Hypophysectomized Rats. *Neuroendocrinology* **24**, 108–118. (doi:10.1159/000122702)
34. Keebaugh AC, Barrett CE, Laprairie JL, Jenkins JJ, Young LJ. 2015 RNAi knockdown of oxytocin receptor in the nucleus accumbens inhibits social attachment and parental care in monogamous female prairie voles. *Soc. Neurosci.* **10**, 561–570. (doi:10.1080/17470919.2015.1040893)
35. Goodson JL, Schrock SE, Kingsbury MA. 2015 Oxytocin mechanisms of stress response and aggression in a territorial finch. *Physiol. Behav.* **141**, 154–163. (doi:10.1016/j.physbeh.2015.01.016)
36. Kleszczyńska A, Sokołowska E, Kulczykowska E. 2012 Variation in brain arginine vasotocin (AVT) and isotocin (IT) levels with reproductive stage and social status in males of three-spined stickleback (*Gasterosteus aculeatus*). *Gen. Comp. Endocrinol.* **175**, 290–296. (doi:10.1016/j.ygcen.2011.11.022)
37. Freeman SM, Samineni S, Allen PC, Stockinger D, Bales KL, Hwa GG, Roberts JA. 2016 Plasma and CSF oxytocin levels after intranasal and intravenous oxytocin in awake macaques. *Psychoneuroendocrinology* **66**, 185–194.
38. Triki Z, Bshary R, Grutter AS, Ros AFH. 2017 The arginine-vasotocin and serotonergic systems affect interspecific social behaviour of client fish in marine cleaning mutualism. *Physiol. Behav.* **174**, 136–143. (doi:10.1016/j.physbeh.2017.03.011)
39. Saito A, Nakamura K. 2011 Oxytocin changes primate paternal tolerance to offspring in food transfer. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* **197**, 329–337. (doi:10.1007/s00359-010-0617-2)
40. Lukas M, Toth I, Reber SO, Slattery DA, Veenema AH, Neumann ID. 2011 The Neuropeptide Oxytocin Facilitates Pro-Social Behavior and Prevents Social Avoidance in Rats and Mice. *Neuropsychopharmacology* **36**, 2159–2168. (doi:10.1038/npp.2011.95)
41. Klatt JD, Goodson JL. 2013 Oxytocin-like receptors mediate pair bonding in a socially monogamous songbird. *Proc. R. Soc. B Biol. Sci.* **280**, 20122396. (doi:10.1098/rspb.2012.2396)

42. Thompson RR, Walton JC. 2004 Peptide Effects on Social Behavior: Effects of Vasotocin and Isotocin on Social Approach Behavior in Male Goldfish (*Carassius auratus*). *Behav. Neurosci.* **118**, 620–626. (doi:10.1037/0735-7044.118.3.620)
43. MacLean EL, Gesquiere LR, Gee NR, Levy K, Martin WL, Carter CS. 2017 Effects of Affiliative Human–Animal Interaction on Dog Salivary and Plasma Oxytocin and Vasopressin. *Front. Psychol.* **8**, 1606. (doi:10.3389/fpsyg.2017.01606)
44. Bottje WG, Wang S, Kinzler S, Neldon HL, Koike TI. 1990 Plasma concentrations of arginine vasotocin and mesotocin following pentobarbital anaesthesia and carotid cannulation in domestic fowl. *Br. Poult. Sci.* **31**, 189–195.
45. Gozdowska M, Kulczykowska E. 2004 Determination of arginine–vasotocin and isotocin in fish plasma with solid-phase extraction and fluorescence derivatization followed by high-performance liquid chromatography. *J. Chromatogr. B* **807**, 229–233.
46. Stocker M, Prosl J, Vanhooland L-C, Horn L, Bugnyar T, Canoine V, Massen JJM. 2021 Measuring salivary mesotocin in birds - Seasonal differences in ravens' peripheral mesotocin levels. *Horm. Behav.* **134**, 105015. (doi:10.1016/j.yhbeh.2021.105015)
47. Samuni L, Preis A, Mundry R, Deschner T, Crockford C, Wittig RM. 2017 Oxytocin reactivity during intergroup conflict in wild chimpanzees. *Proc. Natl. Acad. Sci.* **114**, 268–273. (doi:10.1073/pnas.1616812114)
48. Wirobski G, Range F, Schaebs FS, Palme R, Deschner T, Marshall-Pescini S. 2021 Endocrine changes related to dog domestication: Comparing urinary cortisol and oxytocin in hand-raised, pack-living dogs and wolves. *Horm. Behav.* **128**, 104901. (doi:10.1016/j.yhbeh.2020.104901)
49. Hattori T, Kanno K, Nagasawa M, Nishimori K, Mogi K, Kikusui T. 2015 Impairment of interstrain social recognition during territorial aggressive behavior in oxytocin receptor-null mice. *Neurosci. Res.* **90**, 90–94. (doi:10.1016/j.neures.2014.05.003)
50. Kelly AM, Goodson JL. 2014 Hypothalamic oxytocin and vasopressin neurons exert sex-specific effects on pair bonding, gregariousness, and aggression in finches. *Proc. Natl. Acad. Sci.* **111**, 6069–6074. (doi:10.1073/pnas.1322554111)
51. Modi ME, Young LJ. 2012 The oxytocin system in drug discovery for autism: animal models and novel therapeutic strategies. *Horm. Behav.* **61**, 340–350. (doi:10.1016/j.yhbeh.2011.12.010)
52. Franke AA, Li X, Menden A, Lee MR, Lai JF. 2019 Oxytocin analysis from human serum, urine, and saliva by orbitrap liquid chromatography-mass spectrometry. *Drug Test. Anal.* **11**, 119–128. (doi:10.1002/dta.2475)
53. Weitekamp CA, Solomon-Lane TK, Del Valle P, Triki Z, Nugent BM, Hofmann HA. 2017 A Role for Oxytocin-Like Receptor in Social Habituation in a Teleost. *Brain. Behav. Evol.* **89**, 153–161. (doi:10.1159/000464098)

54. Calcagnoli F, de Boer SF, Beiderbeck DI, Althaus M, Koolhaas JM, Neumann ID. 2014 Local oxytocin expression and oxytocin receptor binding in the male rat brain is associated with aggressiveness. *Behav. Brain Res.* **261**, 315–322. (doi:10.1016/j.bbr.2013.12.050)
55. Morhenn VB, Park JW, Piper E, Zak PJ. 2008 Monetary sacrifice among strangers is mediated by endogenous oxytocin release after physical contact. *Evol. Hum. Behav.* **29**, 375–383. (doi:10.1016/j.evolhumbehav.2008.04.004)
56. Gordon I, Zagoory-Sharon O, Leckman JF, Feldman R. 2010 Oxytocin and the Development of Parenting in Humans. *Biol. Psychiatry* **68**, 377–382. (doi:10.1016/j.biopsych.2010.02.005)
57. Ogi A, Mariti C, Baragli P, Sergi V, Gazzano A. 2020 Effects of Stroking on Salivary Oxytocin and Cortisol in Guide Dogs: Preliminary Results. *Anim. Open Access J. MDPI* **10**, 708. (doi:10.3390/ani10040708)
58. McClung JS, Triki Z, Clément F, Bangerter A, Bshary R. 2018 Endogenous oxytocin predicts helping and conversation as a function of group membership. *Proc. R. Soc. B Biol. Sci.* **285**, 20180939. (doi:10.1098/rspb.2018.0939)
59. Wittig RM, Crockford C, Deschner T, Langergraber KE, Ziegler TE, Zuberbühler K. 2014 Food sharing is linked to urinary oxytocin levels and bonding in related and unrelated wild chimpanzees. *Proc. R. Soc. B Biol. Sci.* **281**, 20133096. (doi:10.1098/rspb.2013.3096)
60. Rincon AV, Deschner T, Schülke O, Ostner J. 2020 Oxytocin increases after affiliative interactions in male Barbary macaques. *Horm. Behav.* **119**, 104661. (doi:10.1016/j.yhbeh.2019.104661)
61. Finkenwirth C, van Schaik C, Ziegler TE, Burkart JM. 2015 Strongly bonded family members in common marmosets show synchronized fluctuations in oxytocin. *Physiol. Behav.* **151**, 246–251. (doi:10.1016/j.physbeh.2015.07.034)
62. Feldman R. 2012 Oxytocin and social affiliation in humans. *Horm. Behav.* **61**, 380–391. (doi:10.1016/j.yhbeh.2012.01.008)
63. Pedersen A, Tomaszycski ML. 2012 Oxytocin antagonist treatments alter the formation of pair relationships in zebra finches of both sexes. *Horm. Behav.* **62**, 113–119. (doi:10.1016/j.yhbeh.2012.05.009)
64. Kabelik D, Magruder DS. 2014 Involvement of different mesotocin (oxytocin homologue) populations in sexual and aggressive behaviours of the brown anole. *Biol. Lett.* **10**, 20140566. (doi:10.1098/rsbl.2014.0566)
65. Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F. 2008 Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput. Biol.* **4**, e1000123.

66. Neumann ID, Landgraf R. 2019 Tracking oxytocin functions in the rodent brain during the last 30 years: From push-pull perfusion to chemogenetic silencing. *J. Neuroendocrinol.* **31**, e12695.
67. Kavaliers M, Choleris E. 2017 Out-Group Threat Responses, In-Group Bias, and Nonapeptide Involvement Are Conserved across Vertebrates: (A Comment on Bruintjes et al., 'Out-Group Threat Promotes Within-Group Affiliation in a Cooperative Fish'). *Am. Nat.* **189**, 453–458. (doi:10.1086/690838)
68. Walum H, Young LJ. 2018 The neural mechanisms and circuitry of the pair bond. *Nat. Rev. Neurosci.* **19**, 643–654.
69. Lehmann L, Keller L. 2006 The evolution of cooperation and altruism--a general framework and a classification of models. *J. Evol. Biol.* **19**, 1365–1376. (doi:10.1111/j.1420-9101.2006.01119.x)
70. Green PA, Briffa M, Cant MA. 2021 Assessment during Intergroup Contests. *Trends Ecol. Evol.* **36**, 139–150. (doi:10.1016/j.tree.2020.09.007)
71. De Dreu CK, Fariña A, Gross J, Romano A. 2021 Pro-sociality as a Foundation for Intergroup Conflict. *Curr. Opin. Psychol.*
72. De Dreu CKW, Kret ME. 2016 Oxytocin Conditions Intergroup Relations Through Upregulated In-Group Empathy, Cooperation, Conformity, and Defense. *Biol. Psychiatry* **79**, 165–173. (doi:10.1016/j.biopsych.2015.03.020)
73. De Dreu CKW. 2012 Oxytocin modulates cooperation within and competition between groups: An integrative review and research agenda. *Horm. Behav.* **61**, 419–428. (doi:10.1016/j.yhbeh.2011.12.009)
74. Ziegler TE, Crockford C. 2017 Neuroendocrine control in social relationships in non-human primates: Field based evidence. *Horm. Behav.* **91**, 107–121. (doi:10.1016/j.yhbeh.2017.03.004)
75. Piva M, Chang SWC. 2018 An integrated framework for the role of oxytocin in multistage social decision-making. *Am. J. Primatol.* **80**, e22735. (doi:10.1002/ajp.22735)
76. Balliet D, Wu J, De Dreu CKW. 2014 Ingroup favoritism in cooperation: A meta-analysis. *Psychol. Bull.* **140**, 1556–1581. (doi:10.1037/a0037737)
77. De Dreu CKW, Greer LL, Van Kleef GA, Shalvi S, Handgraaf MJJ. 2011 Oxytocin promotes human ethnocentrism. *Proc. Natl. Acad. Sci.* **108**, 1262–1266. (doi:10.1073/pnas.1015316108)
78. Yang X, Wang W, Wang XT, Wang YW. 2021 A meta-analysis of hormone administration effects on cooperative behaviours: Oxytocin, vasopressin, and testosterone. *Neurosci. Biobehav. Rev.* **126**, 430–443. (doi:10.1016/j.neubiorev.2021.03.033)

79. Sheng F, Liu Y, Zhou B, Zhou W, Han S. 2013 Oxytocin modulates the racial bias in neural responses to others' suffering. *Biol. Psychol.* **92**, 380–386. (doi:10.1016/j.biopsycho.2012.11.018)
80. Levy J, Goldstein A, Infuls M, Masalha S, Zagoory-Sharon O, Feldman R. 2016 Adolescents growing up amidst intractable conflict attenuate brain response to pain of outgroup. *Proc. Natl. Acad. Sci.* **113**, 13696–13701. (doi:10.1073/pnas.1612903113)
81. van Dijk E, De Dreu CKW. 2021 Experimental Games and Social Decision Making. *Annu. Rev. Psychol.* **72**, 415–438. (doi:10.1146/annurev-psych-081420-110718)
82. Jaeggi AV, Burkart JM, Van Schaik CP. 2010 On the psychology of cooperation in humans and other primates: combining the natural history and experimental evidence of prosociality. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 2723–2735. (doi:10.1098/rstb.2010.0118)
83. Bshary R, Grutter AS. 2006 Image scoring and cooperation in a cleaner fish mutualism. *Nature* **441**, 975–978. (doi:10.1038/nature04755)
84. Fraser ON, Bugnyar T. 2012 Reciprocity of agonistic support in ravens. *Anim. Behav.* **83**, 171–177. (doi:10.1016/j.anbehav.2011.10.023)
85. Milinski M. 1987 Tit for tat in sticklebacks and the evolution of cooperation. *nature* **325**, 433–435.
86. Fehr E, Schurtenberger I. 2018 Normative foundations of human cooperation. *Nat. Hum. Behav.* **2**, 458–468. (doi:10.1038/s41562-018-0385-5)
87. Schweinfurth MK, Call J. 2019 Revisiting the possibility of reciprocal help in non-human primates. *Neurosci. Biobehav. Rev.* **104**, 73–86. (doi:10.1016/j.neubiorev.2019.06.026)
88. Raihani NJ, Grutter AS, Bshary R. 2010 Punishers Benefit From Third-Party Punishment in Fish. *Science* **327**, 171–171. (doi:10.1126/science.1183068)
89. Ross HE, Young LJ. 2009 Oxytocin and the neural mechanisms regulating social cognition and affiliative behavior. *Front. Neuroendocrinol.* **30**, 534–547. (doi:10.1016/j.yfrne.2009.05.004)
90. Marlin BJ, Mitre M, D'amour JA, Chao MV, Froemke RC. 2015 Oxytocin enables maternal behaviour by balancing cortical inhibition. *Nature* **520**, 499–504. (doi:10.1038/nature14402)
91. Kendrick KM, Da Costa AP, Broad KD, Ohkura S, Guevara R, Lévy F, Keverne EB. 1997 Neural control of maternal behaviour and olfactory recognition of offspring. *Brain Res. Bull.* **44**, 383–395. (doi:10.1016/s0361-9230(97)00218-9)
92. Crockford C, Wittig RM, Langergraber K, Ziegler TE, Zuberbühler K, Deschner T. 2013 Urinary oxytocin and social bonding in related and unrelated wild chimpanzees. *Proc. R. Soc. B Biol. Sci.* **280**, 20122765. (doi:10.1098/rspb.2012.2765)

93. De Dreu CKW, Greer LL, Handgraaf MJJ, Shalvi S, Van Kleef GA, Baas M, Ten Velden FS, Van Dijk E, Feith SWW. 2010 The Neuropeptide Oxytocin Regulates Parochial Altruism in Intergroup Conflict Among Humans. *Science* **328**, 1408–1411. (doi:10.1126/science.1189047)
94. Velden F, Daughters K, Dreu C. 2017 Oxytocin promotes intuitive rather than deliberated cooperation with the in-group. *Horm. Behav.* (doi:10.1016/j.yhbeh.2016.06.005)
95. Israel S, Weisel O, Ebstein RP, Bornstein G. 2012 Oxytocin, but not vasopressin, increases both parochial and universal altruism. *Psychoneuroendocrinology* **37**, 1341–1344. (doi:10.1016/j.psyneuen.2012.02.001)
96. Zhang H, Gross J, De Dreu C, Ma Y. 2019 Oxytocin promotes coordinated out-group attack during intergroup conflict in humans. *eLife* **8**, e40698. (doi:10.7554/eLife.40698)
97. Terris ET, Beavin LE, Barraza JA, Schloss J, Zak PJ. 2018 Endogenous Oxytocin Release Eliminates In-Group Bias in Monetary Transfers With Perspective-Taking. *Front. Behav. Neurosci.* **12**, 35. (doi:10.3389/fnbeh.2018.00035)
98. Schiller B, Domes G, Heinrichs M. 2020 Oxytocin changes behavior and spatio-temporal brain dynamics underlying inter-group conflict in humans. *Eur. Neuropsychopharmacol.* **31**, 119–130. (doi:10.1016/j.euroneuro.2019.12.109)
99. Mustoe AC, Cavanaugh J, Harnisch AM, Thompson BE, French JA. 2015 Do marmosets care to share? Oxytocin treatment reduces prosocial behavior toward strangers. *Horm. Behav.* **71**, 83–90. (doi:10.1016/j.yhbeh.2015.04.015)
100. Cavanaugh J, Mustoe A, French JA. 2018 Oxytocin regulates reunion affiliation with a pairmate following social separation in marmosets. *Am. J. Primatol.* **80**, e22750. (doi:10.1002/ajp.22750)
101. Soares MC, Bshary R, Mendonça R, Grutter AS, Oliveira RF. 2012 Arginine Vasotocin Regulation of Interspecific Cooperative Behaviour in a Cleaner Fish. *PLoS ONE* **7**, e39583. (doi:10.1371/journal.pone.0039583)
102. Bosch OJ. 2013 Maternal aggression in rodents: brain oxytocin and vasopressin mediate pup defence. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**, 20130085. (doi:10.1098/rstb.2013.0085)
103. Pedersen CA, Ascher JA, Monroe YL, Prange AJ. 1982 Oxytocin induces maternal behavior in virgin female rats. *Science* **216**, 648–650. (doi:10.1126/science.7071605)
104. Madden JR, Clutton-Brock TH. 2011 Experimental peripheral administration of oxytocin elevates a suite of cooperative behaviours in a wild social mammal. *Proc. Biol. Sci.* **278**, 1189–1194. (doi:10.1098/rspb.2010.1675)
105. Reddon AR, O'Connor CM, Marsh-Rollo SE, Balshine S. 2012 Effects of isotocin on social responses in a cooperatively breeding fish. *Anim. Behav.* **84**, 753–760. (doi:10.1016/j.anbehav.2012.07.021)

106. De Dreu CKW, Shalvi S, Greer LL, Van Kleef GA, Handgraaf MJJ. 2012 Oxytocin Motivates Non-Cooperation in Intergroup Conflict to Protect Vulnerable In-Group Members. *PLoS ONE* **7**, e46751. (doi:10.1371/journal.pone.0046751)
107. Egito JH, Nevat M, Shamay-Tsoory SG, Osório AAC. 2020 Oxytocin increases the social salience of the outgroup in potential threat contexts. *Horm. Behav.* **122**, 104733. (doi:10.1016/j.yhbeh.2020.104733)
108. Samuni L, Preis A, Deschner T, Wittig RM, Crockford C. 2019 Cortisol and oxytocin show independent activity during chimpanzee intergroup conflict. *Psychoneuroendocrinology* **104**, 165–173.
109. Shamay-Tsoory SG, Saporta N, Marton-Alper IZ, Gvirts HZ. 2019 Herding Brains: A Core Neural Mechanism for Social Alignment. *Trends Cogn. Sci.* **23**, 174–186. (doi:10.1016/j.tics.2019.01.002)
110. De Dreu CKW, Gross J, Méder Z, Giffin M, Prochazkova E, Krikeb J, Columbus S. 2016 In-group defense, out-group aggression, and coordination failures in intergroup conflict. *Proc. Natl. Acad. Sci.* **113**, 10524–10529. (doi:10.1073/pnas.1605115113)
111. Stallen M, De Dreu CKW, Shalvi S, Smidts A, Sanfey AG. 2012 The herding hormone: oxytocin stimulates in-group conformity. *Psychol. Sci.* **23**, 1288–1292. (doi:10.1177/0956797612446026)
112. Edelson MG, Shemesh M, Weizman A, Yariv S, Sharot T, Dudai Y. 2015 Opposing Effects of Oxytocin on Overt Compliance and Lasting Changes to Memory. *Neuropsychopharmacology* **40**, 966–973. (doi:10.1038/npp.2014.273)
113. Aydogan G, Jobst A, D'Ardenne K, Müller N, Kocher MG. 2017 The Detrimental Effects of Oxytocin-Induced Conformity on Dishonesty in Competition. *Psychol. Sci.* **28**, 751–759. (doi:10.1177/0956797617695100)
114. Xu L, Becker B, Kendrick KM. 2019 Oxytocin Facilitates Social Learning by Promoting Conformity to Trusted Individuals. *Front. Neurosci.* **13**, 56. (doi:10.3389/fnins.2019.00056)
115. Arueti M, Perach-Barzilay N, Tsoory MM, Berger B, Getter N, Shamay-Tsoory SG. 2013 When Two Become One: The Role of Oxytocin in Interpersonal Coordination and Cooperation. *J. Cogn. Neurosci.* **25**, 1418–1427. (doi:10.1162/jocn_a_00400)
116. Gebauer L, Witek M a. G, Hansen NC, Thomas J, Konvalinka I, Vuust P. 2016 Oxytocin improves synchronisation in leader-follower interaction. *Sci. Rep.* **6**, 38416. (doi:10.1038/srep38416)
117. Mu Y, Guo C, Han S. 2016 Oxytocin enhances inter-brain synchrony during social coordination in male adults. *Soc. Cogn. Affect. Neurosci.* **11**, 1882–1893. (doi:10.1093/scan/nsw106)

118. Kret ME, De Dreu CKW. 2017 Pupil-mimicry conditions trust in partners: moderation by oxytocin and group membership. *Proc R Soc B* **284**, 20162554. (doi:10.1098/rspb.2016.2554)
119. Spengler FB, Scheele D, Marsh N, Kofferath C, Flach A, Schwarz S, Stoffel-Wagner B, Maier W, Hurlemann R. 2017 Oxytocin facilitates reciprocity in social communication. *Soc. Cogn. Affect. Neurosci.* **12**, 1325–1333. (doi:10.1093/scan/nsx061)
120. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, Onaka T, Mogi K, Kikusui T. 2015 Oxytocin-gaze positive loop and the coevolution of human-dog bonds. *Science* **348**, 333–336. (doi:10.1126/science.1261022)
121. Ataei Mehr B, Garner SR, Neff BD. 2020 Effect of isotocin on shoaling behaviour of the Guppy (*Poecilia reticulata*). *Anim. Cogn.* **23**, 827–831. (doi:10.1007/s10071-020-01381-4)
122. Daughters K, Manstead ASR, Ten Velden FS, De Dreu CKW. 2017 Oxytocin modulates third-party sanctioning of selfish and generous behavior within and between groups. *Psychoneuroendocrinology* **77**, 18–24. (doi:10.1016/j.psyneuen.2016.11.039)
123. Aydogan G, Furtner NC, Kern B, Jobst A, Müller N, Kocher MG. 2017 Oxytocin promotes altruistic punishment. *Soc. Cogn. Affect. Neurosci.* **12**, 1740–1747. (doi:10.1093/scan/nsx101)
124. Marsh N, Scheele D, Feinstein JS, Gerhardt H, Strang S, Maier W, Hurlemann R. 2017 Oxytocin-enforced norm compliance reduces xenophobic outgroup rejection. *Proc. Natl. Acad. Sci.* **114**, 9314–9319. (doi:10.1073/pnas.1705853114)
125. Stallen M, Rossi F, Heijne A, Smidts A, Dreu CKWD, Sanfey AG. 2018 Neurobiological Mechanisms of Responding to Injustice. *J. Neurosci.* **38**, 2944–2954. (doi:10.1523/JNEUROSCI.1242-17.2018)
126. Han X, Gelfand MJ, Wu B, Zhang T, Li W, Gao T, Pang C, Wu T, Zhou Y, Zhou S, Wu X, Han S. 2020 A neurobiological association of revenge propensity during intergroup conflict. *eLife* **9**, e52014. (doi:10.7554/eLife.52014)
127. Samuni L, Preis A, Deschner T, Crockford C, Wittig RM. 2018 Reward of labor coordination and hunting success in wild chimpanzees. *Commun. Biol.* **1**, 1–9.
128. Bertalanffy L von. 1969 General system theory: Foundations, development, applications.
129. Young LJ, Wang Z. 2004 The neurobiology of pair bonding. *Nat. Neurosci.* **7**, 1048–1054. (doi:10.1038/nn1327)
130. Oliveira RF. 2009 Social behavior in context: Hormonal modulation of behavioral plasticity and social competence. *Integr. Comp. Biol.* **49**, 423–440. (doi:10.1093/icb/icp055)

131. Cunningham JP, Hereward JP, Heard TA, De Barro PJ, West SA. 2014 Bees at War: Interspecific Battles and Nest Usurpation in Stingless Bees. *Am. Nat.* **184**, 777–786. (doi:10.1086/678399)
132. Goodson JL. 2008 Nonapeptides and the evolutionary patterning of sociality. In *Progress in Brain Research* (eds ID Neumann, R Landgraf), pp. 3–15. Elsevier. (doi:10.1016/S0079-6123(08)00401-9)
133. Rose RM, Bernstein IS, Gordon TP. 1975 Consequences of social conflict on plasma testosterone levels in rhesus monkeys. *Psychosom. Med.* **37**, 50–61. (doi:10.1097/00006842-197501000-00006)