

# **Supporting Information – Ecological networks in the Scotia Sea: structural changes across latitude and depth**

Lucía López-López<sup>1,2\*</sup>, MartinJ. Genner<sup>3</sup>, Geraint A. Tarling<sup>4</sup>, RyanA. Saunders<sup>4</sup>, Eoin J. O’Gorman<sup>5</sup>

<sup>1</sup>*Department of Life Sciences, Imperial College London, Ascot, UK*

<sup>2</sup>*Oceanographic Centre of Balearic Islands, Spanish Institute of Oceanography, Palma de Mallorca, Spain*

<sup>3</sup>*School of Biological Sciences, University of Bristol, Bristol, BS8 1TQ, UK*

<sup>4</sup>*British Antarctic Survey, Cambridge, UK*

<sup>5</sup>*School of Life Sciences, University of Essex, Colchester, UK*

\*corresponding author: [lucia.lopez@ieo.es](mailto:lucia.lopez@ieo.es)

**Figure S1.** Overview of the step-wise procedure used to allocate interactions to every node in the network, indicating the percentage of links allocated in each step.






Figure S2. Histograms of consumer-resource body mass ratios for each of the taxonomic groups considered. Only consumer-resource pairs identified at the species level were used to identify the range of consumer-resource body mass ratios for each taxonomic group.



**Figure S3.** Variation in network-level metrics after randomly excluding between 1 and 25 nodes from the Northern and Southern Scotie Sea food webs (after 999 permutations). The horizontal grey line corresponds to the value of the empirical networks (without removing any nodes). Note that the mean value of the metric does not change for most simulations and the linkage density for the NSS is always greater than that for the SSS.



**Table S1.** Summary table of references on trophic interactions obtained for each node at each step of the bibliographic research. The table includes both qualitative and quantitative diet estimates, with references corresponding to quantitative studies highlighted in bold.

Taxonomic level	Species	Genus	Family	Species	Genus	Family	Species	Genus	Family		
Geographic area	Scotia Sea	Scotia Sea	Scotia Sea	Southern Ocean	Southern Ocean	Southern Ocean	Worldwide	Worldwide	Worldwide		
Node\Step	Step1	Step2	Step3	Step4	Step5	Step6	Step7	Step8	Step9	Assumed	Assumed at
Acantharea				2						1	Class; Scotia Sea
<i>Acanthephyra pelagica</i>		3					4;5				
Aetideus						6;7;8		9			
<i>Alacia belgicae</i>				6;10							
<i>Alacia hettacra</i>				6;10							
<i>Alluroteuthis antarcticus</i>	11;12			13;6;10							
Aloricate ciliates										1,15,16	Subphylum; Worldwide
<i>Antarctomysis maxima</i>				6;10							
Appendicularia										17;18	Class, Worldwide
<i>Arctozenus risso</i>							19;20;21				
<i>Atolla wyvillei</i>							22	23			
<i>Austrinoecia isocheira</i>	24					25					
<i>Bathylagus antarcticus</i>	8			26							

<i>Bathyteuthis abyssicola</i>										11;12;13;27; 28;29;30;31;32; <b>33</b> ;34; 35; <b>37;38;39</b> ; 40	Order, SouthernOce an
<i>Batoteuthis skolops</i>										11;12;13;27; 28;29;30;31;32; <b>33</b> ;34; 35; <b>37;38;39</b> ; 40	Order, SouthernOce an
<i>Benthalbella elongata</i>								<b>41</b>			
<i>Benthalbella macropinna</i>								<b>41</b>			
Beroe								42			
<i>Boroecia antipoda</i>				<b>6;10</b>		25					
<i>Borostomias antarcticus</i>	<b>8;28</b>	<b>26</b>									
<i>Brachioteuthis linkovskyi</i>		27		<b>12;28;29</b>			30				
<i>Brachioteuthis picta</i>	<b>12;28;29</b>										
<i>Calanoides acutus</i>	<b>8;43;44</b> ; 45										
<i>Calanus australis</i>		<b>8;43;44</b> ; 45			47						
<i>Calanus propinquus</i>	<b>8;43;44</b> ; 45;47										
<i>Calanus simillimus</i>	44;45	<b>8;43;47</b>									
Callianira						<b>10</b>					
<i>Calycopsis borchgrevinki</i>							48			49;50; <b>51</b>	Subclass; Worldwide

Candacia								52;53;54			
<i>Chiroteuthis veranyi</i>										11;12;13;27;28;29;30;31;32;33;34;35;37;38;39;40	Order, Southern Ocean
<i>Clausocalanus ingens</i>			45								
<i>Clausocalanus laticeps</i>	45										
<i>Clio piatkowskii</i>			8								
<i>Clio pyramidata</i>	8										
<i>Clione limacina</i>		10;55									
Coccolitophorida											Assumed
Codonellopsis							56			1; 58	Order; Worldwide
<i>Conchoecilla chuni</i>			24								
<i>Conchoecissa symmetrica</i>			24								
Corycaeus								54;59;60			
<i>Crossota brunnea</i>										10;51	Subclass; Southern Ocean
<i>Ctenocalanus citer</i>	24										
Cyclothone			8;26;28								
<i>Cyllopus lucasii</i>	8;28										
<i>Cyllopus magellanicus</i>		8;28									
Cymatocylis										1; 58	Order; Worldwide
<i>Cynomacrurus piriei</i>	26										
<i>Cyphocaris faurei</i>				6							

<i>Cyphocaris richardi</i>				<b>6</b>						
<i>Desmonema glaciale</i>				<b>10</b>						
<i>Dimophyes arctica</i>						51		49;50		
<i>Diphyes antarctica</i>								49;50		
<i>Dissostichus eleginoides</i>	<b>61</b>									
<i>Drepanopus forcipatus</i>	44	62;63		46						
<i>Drepanopus pectinatus</i>	62;63	44			46					
	<b>8;28;64; 66;67;68</b>									
<i>Electrona antarctica</i>	<b>8</b>									
<i>Electrona carlsbergi</i>	<b>67;68</b>			<b>65</b>						
<i>Electrona subaspera</i>				26; <b>65</b>						
<i>Euchirella rostrata</i>				<b>8</b>						
<i>Euchirella rostromagna</i>	<b>8</b>									
<i>Eudoxoides spiralis</i>						49;50				
<i>Eukrohnia hamata</i>	<b>8;69;70</b>									
<i>Euphausia crystallorophias</i>				7;10; <b>75</b>						
<i>Euphausia frigida</i>	<b>8</b>			<b>10</b>						
<i>Euphausia superba</i>	<b>8;24;28; ;76</b>									
<i>Euphausia triacantha</i>		<b>8;24;28; ;76</b>								
<i>Euphausia vallentini</i>		<b>8;24;28; ;76</b>								
<i>Eurythenes obesus</i>				<b>6</b>						
<i>Eusiroides stenopleura</i>									<b>10</b>	Infraorder;Southern Ocean



<i>Eusirus antarcticus</i>	<b>24</b>										
<i>Eusirus microps</i>				<b>6</b>							
<i>Eusirus perdentatus</i>				<b>6;10</b>							
<i>Filippovia knipovitchi</i>	<b>29</b>			<b>12;31</b>							
<i>Gaetanus tenuispinus</i>	<b>8</b>										
<i>Galiteuthis glacialis</i>				<b>13</b>						<b>11;12;13;27;28;29;30;31;32;33;34;35;37;38;39;40</b>	Order; SouthernOcean
<i>Gennadas kempi</i>				<b>3</b>							
<i>Gigantocypris muelleri</i>										<b>6;7;10;24;25</b>	Subclass;SouthernOcean
<i>Gilia reticulata</i>							<b>49;50</b>				
<i>Neognathophausia gigas</i>							<b>80</b>				
<i>Gonatus antarcticus</i>	<b>29</b>			<b>10</b>						<b>11;12;13;27;28;29;30;31;32;33;34;35;37;38;39;40</b>	Order; SouthernOcean
<i>Gymnoscopelus bolini</i>	<b>65</b>			<b>26</b>							
<i>Gymnoscopelus braueri</i>	<b>28;66;67;68;81</b>			<b>26</b>							
<i>Gymnoscopelus fraseri</i>	<b>67;81</b>			<b>26</b>							
<i>Gymnoscopelus hintonoides</i>	<b>8;28;65;66;67;81</b>			<b>26</b>							
<i>Gymnoscopelus nicholsi</i>	<b>65;66;67;81</b>			<b>26</b>							

[illegible]

[illegible]

<i>Obtusoecia antarctica</i>							<b>24</b>			25	Class;Southern Ocean
<i>Oithona frigida</i>	<b>24</b> ;44;45										
<i>Oithona similis</i>	<b>24</b> ;44;45										
<i>Oncaea curvata</i>	<b>24</b> ;94										
<i>Oncaea parila</i>		<b>24</b> ;94									
<i>Oncaea prolata</i>	<b>24</b>										
<i>Onykia ingens</i>				<b>33</b> ;37;38							
<i>Pandea rubra</i>										48;49;50;51	Subclass; Worldwide
<i>Pantachogon haeckeli</i>										10; <b>51</b>	Subclass; Southern Ocean
<i>Paradiplospinus gracilis</i>	<b>66</b>										
<i>Paraeuchaeta antarctica</i>	<b>8</b> ;94; <b>95</b>										
<i>Paraeuchaeta barbata</i>	<b>95</b>										
<i>Paraeuchaeta biloba</i>	<b>95</b>										
<i>Paraeuchaeta rasa</i>	<b>95</b>										
<i>Parandania boeckii</i>	<b>36</b> ;96										
<i>Paraphyllina ransoni</i>										<b>10</b> ;22;23;98;99	Order; Worldwide
<i>Pasiphaea scotiae</i>				<b>6</b>						3;4;5;24	Infraorder; Worldwide
<i>Pelagobia longicirrata</i>	<b>24</b>										
<i>Periphylla periphylla</i>							96;98			10;22;23	Order; Worldwide
<i>Petalidium foliaceum</i>			<b>3</b>								

<i>Phronima sedentaria</i>							83;99;100				
<i>Pleuragramma antarcticum</i>	64										
<i>Pleurobrachia</i>				101			42				
<i>Pleuromamma abdominalis</i>		8									
<i>Pleuromamma antarctica</i>		8									
<i>Pleuromamma robusta</i>	8										
<i>Podocopida</i>										10	Order;South
<i>Poromitra crassiceps</i>										102	ernOcean Assumed
<i>Primno macropa</i>				6							
<i>Protomyctophum andriashevi</i>				26;65							
<i>Protomyctophum bolini</i>	65;66;68;103			26;28							
<i>Protomyctophum choriodon</i>	65;67;103										
<i>Protomyctophum luciferum</i>	65										
<i>Protomyctophum parallelum</i>				26							
<i>Protomyctophum tenisoni</i>	103			26;28							
<i>Pseudoamallothrix cenotelis</i>	8										
<i>Pseudoconchoecia serrulata</i>			24							25	Class;Southe rnOcean
<i>Psychroteuthis glacialis</i>	11			10;13;27							

<i>Pyrostephos vanhoeffeni</i>				51							
<i>Racovitzanus antarcticus</i>	24										
<i>Rhabdonella</i>									1;56;86	Order; Worldwide	
<i>Rhincalanus gigas</i>	8;44										
<i>Rhynchonereella bongraini</i>	24										
<i>Rosacea plicata</i>							49				
<i>Rotundoecia teretivalvata</i>			24						25	Class;Southern Ocean	
<i>Sagitta gazellae</i>	8			69;70							
<i>Sagitta marri</i>	8										
<i>Salpa thompsoni</i>	8			28							
<i>Scaphocalanus australis</i>	8;24										
<i>Scaphocalanus farrani</i>	8										
<i>Scaphocalanus verwoorti</i>	24										
<i>Scina</i>			83								
<i>Scolecithricella minor</i>	24										
<i>Sergestes</i>		3									
<i>Slosarczykovia circumantarctica</i>									11;12;37;27;28;29;30;31;32;33;34;35;37;38;39;40	Order; Southern Ocean	
<i>Solmundella bitentaculata</i>				10;51							
<i>Spinocalanus abyssalis</i>	8										
<i>Spongiobranchaea australis</i>				55;104;105							
<i>Stenosemella</i>					1					86	Order; Worldwide

<i>Stoloteuthis leucoptera</i>										Assumed
<i>Stomias gracilis</i>				26						
<i>Stygiomedusa gigantea</i>							106			
<i>Subeucalanus longiceps</i>			8;44							
<i>Symbolophorus boops</i>	65									
<i>Themisto gaudichaudii</i>	8;36									
<i>Thysanoessa macrura</i>	8									
<i>Thysanoessa vicina</i>		8								
<i>Tintinnopsis</i>									1;56;86	Order; worldwide
<i>Tomopteris planktonis</i>		24								
<i>Tomopteris septentrionalis</i>	24									
<i>Triconia antarctica</i>	8									
<i>Vanadis</i>									14	Family;Worl dwide
<i>Vibilia antarctica</i>		8					57			
<i>Zanclonia weldoni</i>									48;49;510;5 1	Subclass;Wo rldwide

## References for Table S1

1. Gowing MM, Garrison DL. 1992. Abundance and feeding ecology of larger protozooplankton in the ice edge zone of the Weddell and Scotia Seas during the austral winter. *Deep Sea Research Part A. Oceanographic Research Papers* 39: 893–919.
2. Gowing MM. 1989. Abundance and feeding ecology of Antarctic phaeodarian radiolarians. *Marine Biology* 103: 107–118.
3. Foxton P, Roe HSJ. 1974. Observations on the nocturnal feeding of some mesopelagic decapod Crustacea. *Marine Biology* 28: 37–49.
4. Cartes JE. 1993. Feeding habits of oplophorid shrimps in the deep western Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 73:193–206.
5. Burukovsky RN, Falkenhaus T. 2015. Feeding of the pelagic shrimp *Acantheephyra pelagica* (Risso, 1816)(Crustacea: Decapoda: Oplophoridae) in the northern Mid-Atlantic Ridge area in 1984 and 2004. *Arthropoda Selecta* 24: 303–316.
6. Hopkins TL. 1985. Food web of an Antarctic midwater ecosystem. *Marine Biology* 89:197–212
7. Hopkins TL. 1987. Midwater food web in McMurdo Sound, Ross Sea, Antarctica. *Marine Biology* 96: 93–106.
8. Hopkins TL, Ainley DG, Torres JJ, Lancraft TM. 1993. Trophic structure in open waters of the marginal ice zone in the Scotia-Weddell confluence region during spring (1983). *Polar Biology* 13: 389–397.
9. Robertson SB, Frost BW. 1977. Feeding by an omnivorous planktonic copepod *Aetideus divergens* Bradford. *Journal of Experimental Marine Biology and Ecology* 29: 231–244.



10. Jacob U, Brey T, Mintenbeck K. (unpublished) cited in Brose U, Cushing L, Berlow EL, Jonsson T, Banasek-Richter C, Bersier LF, Blanchard JL, Brey T, Carpenter SR, Cattin B, Landenier M-F, Cohen JE, Dawah HA, Edwards F, Harper-Smith S, Jacob U, Knapp RA, Ledger ME, Memmott J, Mintenbeck K, Pinnegar JK, Rall CR, Rayner T, Ruess L, Ulrich W, Warren P, Williams RJ, Woodward G, Yodanis P, Martinez ND. 2005. Body sizes of consumers and their resources. *Ecology* 86: 2545–2545.
11. Kear AJ. 1992. The diet of Antarctic squid: comparison of conventional and serological gut contents analyses. *Journal of Experimental Marine Biology and Ecology* 156: 161–178.
12. Nemoto T, Okiyama M, Takahashi M. 1985. Aspects of the roles of squid in food chains of marine Antarctic ecosystems. Sigfried WR, Condry PR, Laws RM, editors. *Antarctic Nutrient Cycles and Food Webs*. Berlin, Heidelberg: Springer-Verlag. p. 415–420.
13. Lu CC, Williams R. 1994. Contribution to the biology of squid in the Prydz Bay region, Antarctica. *Antarctic Science* 6:223–229
14. Jumars PA, Dorgan KM, Lindsay SM. 2015. Diet of worms emended: an update of polychaete feeding guilds. *Annual Review of Marine Science* 7: 497–520.
15. Jonsson PR. 1986. Particle size selection, feeding rates and growth dynamics of marine planktonic oligotrichous ciliates (Ciliophora: Oligotrichina). *Marine Ecology Progress Series* 33:265–277.
16. Pitta P, Giannakourou A, Christaki U. 2001. Planktonic ciliates in the oligotrophic Mediterranean Sea: longitudinal trends of standing stocks, distributions and analysis of food vacuole contents. *Aquatic Microbial Ecology* 24: 297–311.
17. Montú M, de Oliveira IR. 1986. Zooplanktonic associations, trophic relations and standing stock of krill and other groups of the community near Elephant Island (February-March 84/85). *Revista Nérítica* 1.

18. Lombard F, Selander E, Kiørboe T. 2011. Active prey rejection in the filter-feeding appendicularian *Oikopleura dioica*. *Limnology and Oceanography* 56: 1504–1512.
19. Sherwood GD, Rose GA. 2005. Stable isotope analysis of some representative fish and invertebrates of the Newfoundland and Labrador continental shelf food web. *Estuarine, Coastal and Shelf Science* 6: 537–549.
20. Carlier A, Le Guilloux E, Olu K, Sarrazin J, Mastrototaro F, Taviani M, Clavier J. 2009. Trophic relationships in a deep Mediterranean cold-water coral bank (Santa Maria di Leuca, Ionian Sea). *Marine Ecology Progress Series* 397: 125–137.
21. Il'inskiy EN, Balanov AA, Ivanov OA. 1995. Rare mesopelagic fishes *Scopelosaurus harryi*, *Arctozenus rissoi*, *Magnisudis atlantica* and *Tactostoma macropus* from the Northwest Pacific. 2. Spatial distribution and biology. *Journal of Ichthyology* 35: 1–19.
22. Larson RJ. 1979. Feeding in coronate medusae (class scyphozoa, order coronatae). *Marine and Freshwater Behaviour and Physiology* 6: 123–129.
23. Hunt JC, Lindsay DJ. 1998. Observations on the behavior of Atolla (Scyphozoa: Coronatae) and Nanomia (Hydrozoa: Physonectae): use of the hypertrophied tentacle in prey capture. *Plankton Biology and Ecology* 45: 239–242.
24. Hopkins TL, Torres JJ. 1989. Midwater food web in the vicinity of a marginal ice zone in the western Weddell Sea. *Deep Sea Research Part A. Oceanographic Research Papers* 36: 543–560.
25. Angel MV, Blachowiak-Samolyk K. 2014. Halocyprid ostracods of the Southern Ocean. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Robert Coudert, Y, editors. *Biogeographic Atlas of the Southern Ocean*. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

26. Gaskett AC, Bulman C, HeX, Goldsworthy SD. 2001. Diet composition and guild structure of mesopelagic and bathypelagic fishes near Macquarie Island, Australia. *New Zealand Journal of Marine and Freshwater Research* 35: 469–473.
27. Filippova JA. 2002. Review of Soviet/Russian studies on squids in the Antarctic Ocean. *Bulletin of Marine Science* 71: 255–267.
28. Lancraft TM, Hopkins TL, Torres JJ, Donnelly J. 1991. Oceanic micronektonic/macrozooplanktonic community structure and feeding in ice covered Antarctic waters during the winter (AMERIEZ 1988). *Polar Biology* 11:157–167.
29. Rodhouse PG, Prince PA, Trathan PN, Hatfield EMC, Watkins JL, Bone DG, Murphy EJ, White MG. 1996. Cephalopods and mesoscale oceanography at the Antarctic Polar Front: satellite tracked predators locate pelagic trophic interactions. *Marine Ecology Progress Series* 136:37–50.
30. Roper CF, Vecchione M. 1996. In situ observations on *Brachioteuthis beanie* Verrill: paired behavior, probably mating (Cephalopoda, Oegopsida). *American Malacological Bulletin* 13:55–60.
31. Collins MA, Allcock AL, Belchier M. 2004. Cephalopods of the South Georgia slope. *Journal of the Marine Biological Association of the United Kingdom* 84: 415–419.
32. Dickson J, Morley SA, Mulvey T. 2004. New data on *Martialia hyadesi* feeding in the Scotia Sea during winter with emphasis on seasonal and annual variability. *Journal of the Marine Biological Association of the United Kingdom* 84: 785–788.
33. Cherel Y, Duhamel G. 2003. Diet of the squid *Moroteuthis ingens* (Teuthoidea: Onychoteuthidae) in the upper slope waters of the Kerguelen Islands. *Marine Ecology Progress Series* 250: 197–203.

34. Gonzalez AF, Trathan PN, Yau C, Rodhouse PG. 1997. Interactions between oceanography, ecology and fishery biology of the ommastrephid squid *Martialia hyadesi* in the South Atlantic. Marine Ecology Progress Series 152: 205–215.
35. Gonzalez AF, Rodhouse PG. 1998. Fishery biology of the seven star flying squid *Martialia hyadesi* at South Georgia during winter. Polar Biology 19: 231–236.
36. Pakhomov EA, Perissinotto R. 1996. Trophodynamics of the hyperiid amphipod *Themisto gaudichaudi* in the South Georgia region during late austral summer. Marine Ecology Progress Series 134: 91–100.
37. Phillips KL, Jackson GD, Nichols PD. 2001. Predation on myctophids by the squid *Moroteuthis ingens* around Macquarie and Heard Islands: stomach contents and fatty acid analyses. Marine Ecology Progress Series 215: 179–189.
38. Phillips KL, Nichols PD, Jackson GD. 2003. Dietary variation of the squid *Moroteuthis ingens* at four sites in the Southern Ocean: stomach contents, lipid and fatty acid profiles. Journal of the Marine Biological Association of the United Kingdom 83:523–534.
39. Rodhouse PG, White MG, Jones MRR. 1992. Trophic relations of the cephalopod *Martialia hyadesi* (Teuthoidea: Ommastrephidae) at the Antarctic Polar Front, Scotia Sea. Marine Biology 114: 415–421.
40. Xavier JC, Cherel Y, Allcock L, Rosa R, Sabirov RM, Blicher ME, Golikov AV. 2018. A review on the biodiversity, distribution and trophic role of cephalopods in the Arctic and Antarctic marine ecosystems under a changing ocean. Marine Biology 165: 93.
41. Yang MS. 2011. Diet of Nineteen Mesopelagic Fishes in the Gulf of Alaska. US Dep. Commer. –NOAA Tech. Memo. NMFS-AFSC-229.
42. Lebour MV. 1923. The food of plankton organisms. II. Journal of the Marine Biological Association of the United Kingdom 13: 70–92.

43. Burghart SE, Hopkins TL, Vargo GA, Torres JJ. 1999. Effects of a rapidly receding ice edge on the abundance, age structure and feeding of three dominant calanoid copepods in the Weddell Sea, Antarctica. *Polar Biology* 22: 279–288.
44. Atkinson A. 1994. Diets and feeding selectivity among the epipelagic copepod community near South Georgia in summer. *Polar Biology* 14:551–560.
45. Atkinson A. 1996. Subantarctic copepods in an oceanic, low chlorophyll environment: ciliate predation, food selectivity and impact on prey populations. *Marine Ecology Progress Series* 130: 85–96.
46. Antacli JC. 2011. Estrategias de vida de los copépodos *Drepanopus forcipatus* y *Calanus australis* en relación con los recursos tróficos en la plataforma Patagónica Austral (Argentina, 47°-55°S). PhD Thesis. –Universidad Nacional del Mar de Plata, Argentina.
47. Metz C, Schnack- Schiel SB. 1995. Observations on carnivorous feeding in Antarctic calanoid copepods. *Marine Ecology Progress Series* 129:71–75.
48. Alvariño A. (1981). Siphonophorae. Boltovskoy D, editor. Atlas del Zooplancton del Atlántico Sudoccidental y métodos de trabajo con el zooplancton marino. –INJDEP, Mar del Plata, Argentina.
49. Purcell JE. 1981. Dietary composition and diel feeding patterns of epipelagic siphonophores. *Marine Biology* 65: 83–90.
50. Purcell JE. 1982. Feeding and growth of the siphonophore *Muggiaea atlantica* (Cunningham 1893). *Journal of Experimental Marine Biology and Ecology* 62: 39–54.
51. Pagès F, Schnack- Schiel SB. 1996. Distribution patterns of the mesozooplankton, principally siphonophores and medusae, in the vicinity of the Antarctic Slope Front (eastern Weddell Sea). *Journal of Marine Systems* 9:231–248.

52. Ohtsuka S, Onbé T. 1989. Evidence of selective feeding on larvaceans by the pelagic copepod *Candacia bipinnata* (Calanoida: Candaciidae). *Journal of Plankton Research* 11: 869–872.
53. Schnack SB. 1989. Functional morphology of feeding appendages in calanoid copepods. Felgenhauer BE, Watling L, Thistle AB, editors. *Functional morphology of feeding and grooming in Crustacea*. Rotterdam: Balkema. p. 137–151.
54. Benedetti F, Gasparini S, Ayata SD. 2015. Identifying copepod functional groups from species functional traits. *Journal of Plankton Research* 38:159–166.
55. Atkinson A, Ward P, HuntBPV, Pakhomov EA, Hosie GW. 2012. An overview of Southern Ocean zooplankton data: abundance, biomass, feeding and functional relationships. *CCAMLR Science* 19: 171–218.
56. Thompson GA, Alder VA. 2005. Patterns in tintinnid species composition and abundance in relation to hydrological conditions of the southwestern Atlantic during austral spring. *Aquatic Microbial Ecology* 40: 85–101.
57. Madin LP, Harbison GR. 1977. The associations of Amphipoda Hyperiidea with gelatinous zooplankton—I. Associations with Salpidae. *Deep Sea Research* 24: 449–463.
58. Pierce RW, Turner JT. 1992. Ecology of planktonic ciliates in marine food webs. *Reviews in Aquatic Science* 6: 139–181.
59. Lebour MV. 1922. The food of plankton organisms. *Journal of the Marine Biological Association of the United Kingdom* 12: 644–677.
60. Landry MR, Lehner-Fournier JM, Fagerness VL. 1985. Predatory feeding behavior of the marine cyclopoid copepod *Corycaeus anglicus*. *Marine Biology* 85: 163–169.

61. Xavier J, Rodhouse P, Purves M, Daw T, Arata J, Pilling G. 2002. Distribution of cephalopods recorded in the diet of the Patagonian toothfish (*Dissostichus eleginoides*) around South Georgia. *Polar Biology* 25: 323–330.
62. Mayzaud P, Razouls S. 1992. Degradation of gut pigment during feeding by a subantarctic copepod: importance of feeding history and digestive acclimation. *Limnology and Oceanography* 37: 393–404.
63. Alonzo F, Mayzaud P, Razouls S. 2001. Egg production and energy storage in relation to feeding conditions in the subantarctic calanoid copepod *Drepanopuspectinatus*: an experimental study of reproductive strategy. *Marine Ecology Progress Series* 209: 231–242.
64. Williams R. 1985. Trophic relationships between pelagic fish and euphausiids in Antarctic waters. Sigfried WR, Condry PR, Laws RM, editors. *Antarctic nutrient cycles and food webs*. Berlin, Heidelberg: Springer. p. 452–459.
65. Pakhomov EA, Perissinotto R, McQuaid CD. 1996. Prey composition and daily rations of myctophid fishes in the Southern Ocean. *Marine Ecology Progress Series* 134: 1–14.
66. Pusch C, Hulley PA, Kock KH. 2004. Community structure and feeding ecology of mesopelagic fishes in the slope waters of King George Island (South Shetland Islands, Antarctica). *Deep Sea Research Part I: Oceanographic Research Papers* 51: 1685–1708.
67. Shreeve RS, Collins MA, Tarling GA, Main CE, Ward P, Johnston NM. 2009. Feeding ecology of myctophid fishes in the northern Scotia Sea. *Marine Ecology Progress Series* 386: 221–236.
68. Saunders RA, Collins MA, Foster E, Shreeve R, Stowasser G, Ward P, Tarling GA. 2014. The trophodynamics of Southern Ocean Electrona (Myctophidae) in the Scotia Sea. *Polar Biology* 37: 789–807.

69. Froneman PW, Pakhomov EA, Perissinotto R, Meaton V. 1998. Feeding and predation impact of two chaetognath species, *Eukrohnia hamata* and *Sagitta gazellae*, in the vicinity of Marion Island (Southern Ocean). *Marine Biology* 131: 95–101.
70. Froneman PW, Pakhomov EA. 1998. Trophic importance of the chaetognaths *Eukrohnia hamata* and *Sagitta gazellae* in the pelagic system of the Prince Edward Islands (Southern Ocean). *Polar Biology* 19:242–249.
71. Kruse S, Hagen W, Bathmann U. 2010. Feeding ecology and energetics of the Antarctic chaetognaths *Eukrohnia hamata*, *E. bathypelagica* and *E. bathyantartica*. *Marine Biology* 157:2289–2302.
72. Giesecke R, González HE. 2012. Distribution and feeding of chaetognaths in the epipelagic zone of the Lazarev Sea (Antarctica) during austral summer. *Polar Biology* 35:689–703.
73. Øresland V. 1990. Feeding and predation impact of the chaetognath *Eukrohnia hamata* in Gerlache Strait, Antarctic Peninsula. *Marine Ecology Progress Series* 63:201–209.
74. Øresland V. 1995. Winter population structure and feeding of the chaetognath *Eukrohnia hamata* and the copepod *Euchaeta antarctica* in Gerlache Strait, Antarctic Peninsula. *Marine Ecology Progress Series* 119: 77–86.
75. Pakhomov EA, Perissinotto R, Froneman PW. 1998. Abundance and trophodynamics of *Euphausia crystallorophias* in the shelf region of the Lazarev Sea during austral spring and summer. *Journal of Marine Systems* 1: 313–324.
76. Pakhomov EA, Perissinotto R, Froneman PW, Miller DGM. 1997. Energetics and feeding dynamics of *Euphausia superba* in the South Georgia region during the summer of 1994. *Journal of Plankton Research* 19:399–423.
77. Price HJ, Boyd KR, Boyd CM. 1988. Omnivorous feeding behavior of the Antarctic krill *Euphausia superba*. *Marine Biology* 97: 67–77.



78. Meyer B, Fuentes V, Guerra C, Schmidt K, Atkinson A, Spahic S, Cisewski B, Freier U, Olariaga A, Bathmann U. 2009. Physiology, growth, and development of larval krill *Euphausia superba* in autumn and winter in the Lazarev Sea, Antarctica. *Limnology and Oceanography* 54: 1595–1614.
79. Blachowiak-Samolyk K, Angel M. 2008. An Atlas of Southern Ocean Planktonic Ostracods. <http://ocean.iopan.gda.pl/ostracoda/index.php>
80. Hopkins TL, Flock ME, Gartner Jr JV, Torres JJ. 1994. Structure and trophic ecology of a low latitude midwater decapod and mysid assemblage. *Marine Ecology Progress Series* 109:143–156.
81. Saunders RA, Collins MA, Ward P, Stowasser G, Shreeve R, Tarling GA. 2015. Distribution, population structure and trophodynamics of Southern Ocean *Gymnoscopelus* (Myctophidae) in the Scotia Sea. *Polar Biology* 38: 287–308.
82. Gasca R, Hoover R, Haddock SH. 2015. New symbiotic associations of hyperiid amphipods (Peracarida) with gelatinous zooplankton in deep waters off California. *Journal of the Marine Biological Association of the United Kingdom* 95:503–511.
83. Lourenço S, Saunders RA, Collins M, Shreeve R, Assis CA, Belchier M, Watkins JL, Xavier JC. 2017. Life cycle, distribution and trophodynamics of the lanternfish *Krefftichthys anderssoni* (Lönnberg, 1905) in the Scotia Sea. *Polar Biology* 40: 1229–1245.
84. Gasca R, Suárez-Morales E, Haddock SH. 2007. Symbiotic associations between crustaceans and gelatinous zooplankton in deep and surface waters off California. *Marine Biology* 151: 233–242.
85. Zeidler W. 2012. A review of the hyperiidean amphipod families Mimonectidae and Proscinidae (Crustacea: Amphipoda: Hyperiidea: Scinoidea). *Zootaxa* 3533: 1–74.

86. Turner JT. 1984. The feeding ecology of some zooplankters that are important prey items of larval fish. NOAA Technical Report NMFS 7. p. 1–28.
87. Gowing MM, Wishner KF. 1998. Feeding ecology of the copepod *Lucicutia aff. L. grandis* near the lower interface of the Arabian Sea oxygen minimum zone. Deep Sea Research Part II: Topical Studies in Oceanography 45:2433–2459.
88. Burghart SE, Hopkins TL, Vargo GA, Torres JJ. 1999. Effects of a rapidly receding ice edge on the abundance, age structure and feeding of three dominant calanoid copepods in the Weddell Sea, Antarctica. Polar Biology 22: 279–288.
89. Koski M, Kiørboe T, Takahashi K. 2005. Benthic life in the pelagic: aggregate encounter and degradation rates by pelagic harpacticoid copepods. Limnology and Oceanography 50: 1254–1263.
90. Koski M, Møller EF, Maar M, Visser AW. 2007. The fate of discarded appendicularian houses: degradation by the copepod, *Microsetella norvegica*, and other agents. Journal of Plankton Research 29: 641–654.
91. Ohtsuka S, Kubo N, Okada M, Gushima K. 1993. Attachment and feeding of pelagic copepods on larvacean houses. Journal of Oceanography 49: 115–120.
92. Stowasser G, Pond DW, Collins MA. 2009. Using fatty acid analysis to elucidate the feeding habits of Southern Ocean mesopelagic fish. Marine Biology 156:2289–2302. Ecology
93. Metz C. 1998. Feeding of *Oncaea curvata* (Poecilostomatoida, Copepoda). Marine Progress Series 169:229–235.
94. Øresland V. 1991. Feeding of the carnivorous copepod *Euchaeta antarctica* in Antarctic waters. Marine Ecology Progress Series 78: 41–47.
95. Øresland V, Ward P. 1993. Summer and winter diet of four carnivorous copepod species around South Georgia. Marine Ecology Progress Series 98: 73–78.

96. Coleman CO 1990. Anatomy of the alimentary canal of *Parandania boeckii* (Stebbing, 1888) (Crustacea, Amphipoda, Stegocephalidae) from the Antarctic Ocean. *Journal of Natural History* 24:1573–1585.
97. Fosså JH. 1992. Mass occurrence of *Periphylla periphylla* (Scyphozoa, Coronatae) in a Norwegian fjord. *Sarsia* 77:237–251
98. Youngbluth MJ, Båmstedt U. 2001. Distribution, abundance, behaviour and metabolism of *Periphylla periphylla*, a meso-pelagic coronate medusa in a Norwegian fjord. *Hydrobiologia* 451:321–333
99. Laval P. 1968. Observations sur la biologie de *Phronima curvipes* Voss. (Amphipoda, Hyperide) et description du mâleadulte. *Cahiers de Biologie Marine* 9:347–362.
100. Diebel CE. 1988. Observations on the anatomy and behavior of *Phronima sedentaria* (Forskal) (Amphipoda: Hyperiidea). *Journal of Crustacean Biology* 8: 79–90.
101. Alvarino A. 1985. Predation in the plankton realm, mainly with reference to fish larvae. *Investigaciones Marinas CICIMAR* 2: 1–122.
102. Hart JL. 1973. Pacific fishes of Canada. (Bulletin N.180) Fisheries Research board of Canada. p.740.
103. Saunders RA, Collins MA, Ward P, Stowasser G, Shreeve R, Tarling GA. 2015. Trophodynamics of *Protomyctophum* (Myctophidae) in the Scotia Sea (Southern Ocean). *Journal of Fish Biology* 87: 1031–1058.
104. Lalli CM, Gilmer RW. 1989. Pelagic snails: the biology of holoplanktonic gastropod mollusks. Stanford, California: Stanford University Press. p. 253.
105. Phleger CF, Nelson MM, Mooney B, Nichols PD. 1999. Lipids of abducted Antarctic pteropods, *Spongiobranchaea australis*, and their hyperiid amphipod host.

Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology 124: 295–307.

106. Benfield MC, Graham WM. 2010. In situ observations of *Stygiomedusa gigantea* in the Gulf of Mexico with a review of its global distribution and habitat. *Journal of the Marine Biological Association of the United Kingdom* 90:1079–1093.

**Table S2.** Summary table of node taxonomy, taxonomic groups, mean masses, and known distributions in latitude and depth. Superscripts are references for the conversion factors used to calculate mean mass, with the exception of [<sup>A</sup>] which indicates an assumption.

Node	Taxonomy	Group	Mass	Latitudinal distribution	Depth distribution	References
Acantharea	Class	Protist	1.00E-07 <sup>A</sup>	Both <sup>a</sup>	All <sup>a</sup>	
<i>Acantheephyra pelagica</i>	Species	Decapod	7.20E+00	Both	Meso + bathypelagic	1,2
Aetideus	Genus	Copepod	1.03E-03 <sup>b,c</sup>	Both	All	3,4
<i>Alacia belgicae</i>	Species	Ostracod	4.40E-04 <sup>A</sup>	South	Epipelagic	5,6
<i>Alacia hettacra</i>	Species	Ostracod	4.40E-04	Both	Mesopelagic	5,7
<i>Alluroteuthis antarcticus</i>	Species	Squid	2.90E+01	Both	Mesopelagic	5,8
Aloricate ciliates	Subphylum	Protist	1.00E-09	Both <sup>a</sup>	All <sup>a</sup>	
Amphidinium	Genus	Primary producer	1.63E-09 <sup>a</sup>	Both	Epipelagic	
Amphisolenia	Genus	Primary producer	1.95E-08 <sup>a</sup>	Both	Epipelagic	
<i>Antarctomysis maxima</i>	Species	Decapod	7.21E-01	Both	All	9,10
Appendicularia	Class	Mesh feeder	2.57E-03	Both	All	1,11
<i>Arctozenus risso</i>	Species	Fish	3.01E-01	Both	All	12,13
<i>Atolla wyvillei</i>	Species	Gelatinous carnivore	3.18E+01	Both	All	1
<i>Austrinoecia isocheira</i>	Species	Ostracod	3.39E-05 <sup>d</sup>	Both	Mesopelagic	7
Bacteria		Primary producer	1.00E-12	Both <sup>a</sup>	All <sup>a</sup>	
<i>Bathylagus antarcticus</i>	Species	Squid	2.70E+01	South	All	5,14
<i>Bathyteuthis abyssicola</i>	Species	Squid	3.56E+00	Both	All	1,6
<i>Batoteuthis skolops</i>	Species	Squid	9.95E+00 <sup>e</sup>	Both	Meso + bathypelagic	6,15
<i>Benthalbella elongata</i>	Species	Fish	1.65E+01	Both	Meso + bathypelagic	1,14

<i>Benthalbella macropinna</i>	Species	Fish	1.65E+01 <sup>A</sup>	Both	Meso + bathypelagic	14
Beroe	Genus	Gelatinous carnivore	5.88E+01	Both	All	9,11
<i>Boroecia antipoda</i>	Species	Ostracod	4.40E-04	Both	Mesopelagic	5,7
<i>Borostomias antarcticus</i>	Species	Fish	2.30E+01	Both	Meso + bathypelagic	1,14
<i>Brachioteuthis linkovskyi</i>	Species	Squid	2.10E+00 <sup>e</sup>	North	All	15,16
<i>Brachioteuthis picta</i>	Species	Squid	1.15E+00 <sup>e</sup>	Both	All	16
<i>Calanoides acutus</i>	Species	Copepod	5.50E-03	Both	Epi + mesopelagic	5,17
<i>Calanus australis</i>	Species	Copepod	4.92E-03 <sup>b,c</sup>	North	Meso + bathypelagic	3, 4
<i>Calanus propinquus</i>	Species	Copepod	5.70E-03	Both	Epi + mesopelagic	5,17
<i>Calanus simillimus</i>	Species	Copepod	6.30E-03 <sup>b,c</sup>	North	Epi + mesopelagic	3,4
Callianira	Genus	Gelatinous carnivore	5.60E+00	Both	All	5
<i>Calycopsis borchgrevinki</i>	Species	Gelatinous carnivore	1.75E+00	Both	All	1
Candacia	Genus	Copepod	9.04E-03 <sup>b,c</sup>	Both	All	3,4
Ceratium	Genus	Primaryproducer	6.96E-09 <sup>A</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
Chaetoceros	Genus	Primary producer	5.28E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	18
<i>Chroteuthis veranyi</i>	Species	Squid	3.64E+01 <sup>f</sup>	North	Meso + bathypelagic	15,19
<i>Clausocalanus ingens</i>	Species	Copepod	5.69E-04 <sup>b,c</sup>	North	Epi + mesopelagic	3,17
<i>Clausocalanus laticeps</i>	Species	Copepod	3.92E-04 <sup>b,c</sup>	North	All	3,17
<i>Clio piatkowskii</i>	Species	Mesh feeder	1.70E-02	Both	Epipelagic	5,20
<i>Clio pyramidata</i>	Species	Mesh feeder	1.70E-02	Both	Epipelagic	5,20
<i>Clione limacina</i>	Species	Mesh feeder	1.40E-02	Both	Epipelagic	5,11
Coccolitophorida		Primary producer	5.23E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	
Codonellopsis	Genus	Protist	1.00E-08 <sup>A</sup>	Both	All	21

<i>Conchoecilla chuni</i>	Species	Ostracod	7.54E-03 <sup>g</sup>	North	Mesopelagic	7
<i>Conchoecissa symmetrica</i>	Species	Ostracod	4.37E-02 <sup>g</sup>	North	Meso + bathypelagic	7
Corethron	Genus	Primary producer	4.45E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
Corycaeus	Genus	Copepod	1.01E-03 <sup>b,c</sup>	Both <sup>a</sup>	All <sup>a</sup>	3
		Gelatinous			Meso + bathypelagic	
<i>Crossota brunnea</i>	Species	carnivore	9.00E-02	Both		22,
<i>Ctenocalanus citer</i>	Species	Copepod	3.40E-04 <sup>b,c</sup>	North	All	3,4
					Meso + bathypelagic	
Cyclothone	Genus	Fish	5.60E+00	Both		1,24
<i>Cyllopus lucasii</i>	Species	Amphipod	3.20E-01	Both	Epipelagic	5,25
<i>Cyllopus magellanicus</i>	Species	Amphipod	3.20E-01	Both	Epipelagic	5,25
Cymatocylis	Genus	Protist	1.00E-08 <sup>A</sup>	Both	All	21
<i>Cynomacrus piriei</i>	Species	Fish	4.60E+01 <sup>h</sup>	Both	All	14,27
<i>Cyphocaris faurei</i>	Species	Amphipod	1.28E+00	Both	All	1,28
<i>Cyphocaris richardi</i>	Species	Amphipod	2.21E+00	Both	All	1,28
Dactyliosolen	Genus	Primary producer	1.41E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
		Gelatinous				
<i>Desmonema glaciale</i>	Species	carnivore	3.50E+04	Both	Epi + mesopelagic	23,29
Detritus		Primary producer		Both <sup>a</sup>	All <sup>a</sup>	Assumed
		Gelatinous				
<i>Dimophyes arctica</i>	Species	carnivore	2.00E+00 <sup>A</sup>	Both	All	11
Dinophysis	Genus	Primary producer	1.95E-08 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
		Gelatinous				
<i>Diphyes antarctica</i>	Species	carnivore	2.09E+00	Both	All	9,23
<i>Dissostichus eleginoides</i>	Species	Fish	6.14E+03	Both	All	13
<i>Drepanopus forcipatus</i>	Species	Copepod	2.13E-03 <sup>b,c</sup>	North	Epi + mesopelagic	3,4
<i>Drepanopus pectinatus</i>	Species	Copepod	2.84E-03 <sup>b,c</sup>	North	Epi + mesopelagic	3,4
Dictyochophyceae	Class	Primary producer	2.91E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
<i>Electrona antarctica</i>	Species	Fish	9.98E+00 <sup>i</sup>	Both	All	14,27

<i>Electrona carlsbergi</i>	Species	Fish	6.09E+00 <sup>i</sup>	North	All	14,27	
<i>Electrona subaspera</i>	Species	Fish	6.00E+00 <sup>A</sup>	North	All	14,27	
<i>Eucampia</i>	Genus	Primary producer	6.91E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Euchirella rostrata</i>	Species	Copepod	5.10E-03	South	All	3,4	
					Meso + bathypelagic		3
<i>Euchirella rostromagna</i>	Species	Copepod	3.78E-02	Both			
		Gelatinous					
<i>Eudoxoides spiralis</i>	Species	carnivore	2.50E-01 <sup>A</sup>	Both	All		23
		Gelatinous					
<i>Eukrohnia hamata</i>	Species	carnivore	5.20E-03	Both	All	5,30	
<i>Euphausia crystallorophias</i>	Species	Euphausiid	2.49E-01	South	Epi + mesopelagic	9,31	
<i>Euphausia frigida</i>	Species	Euphausiid	1.70E-01	Both	Epi + mesopelagic	5,31	
<i>Euphausia superba</i>	Species	Euphausia superba	1.45E+00	Both	All	9,31	
<i>Euphausiatriacantha</i>	Species	Euphausiid	6.00E-01 <sup>c</sup>	Both	Mesopelagic	31,32	
<i>Euphausia vallentini</i>	Species	Euphausiid	1.70E-01	North	Epipelagic		31
<i>Eurythenes obesus</i>	Species	Amphipod	3.70E-01	Both	All	1,33	
<i>Eusiroides stenopleura</i>	Species	Amphipod	1.63E-01	Both <sup>a</sup>	All <sup>a</sup>		1
<i>Eusirus antarcticus</i>	Species	Amphipod	4.70E-02	South	All	11,28	
<i>Eusirus microps</i>	Species	Amphipod	6.99E-01	South	All	11,28	
<i>Eusirus perdentatus</i>	Species	Amphipod	1.90E+01	Both <sup>a</sup>	All <sup>a</sup>		5
					Meso + bathypelagic		8
<i>Filippovia knipovitchi</i>	Species	Squid	3.90E+02 <sup>e</sup>	Both			
Foraminifera	Phylum	Protist	5.00E-06	Both	All		34
Fragilariopsis	Genus	Primary producer	2.39E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Gaetanus tenuispinus</i>	Species	Copepod	6.36E-03 <sup>b,c</sup>	Both	All	3, 35	
<i>Galiteuthis glacialis</i>	Species	Squid	5.95E+01	Both	All	1,8	
					Meso + bathypelagic		
<i>Gennadas kemp</i>	Species	Decapod	7.35E-01	Both		1,2	
<i>Gigantocypris muelleri</i>	Species	Ostracod	1.22E+00	Both	All	7,28	



<i>Gilia reticulata</i>	Species	Gelatinous carnivore	2.50E-01	Both	Mesopelagic Meso + bathypelagic	23
<i>Neognathophausia gigas</i>	Species	Decapod	6.94E-01	Both	Meso + bathypelagic	1,2
<i>Gonatus antarcticus</i>	Species	Squid	4.28E+02 <sup>e</sup>	Both <sup>a</sup>	bathypelagic <sup>a</sup>	36
<i>Grammatophora</i>	Genus	Primary producer	1.98E-08 <sup>a</sup>	Both	Epipelagic	18
<i>Gymnoscopelus bolini</i>	Species	Fish	2.19E+01	Both	All	14,37
<i>Gymnoscopelus braueri</i>	Species	Fish	2.50E+01	Both	All	5,14
<i>Gymnoscopelus fraseri</i>	Species	Fish	8.73E+00 <sup>j</sup>	North	All	5,14
<i>Gymnoscopelus hintonoides</i>	Species	Fish	2.50E+01	North	All	14
<i>Gymnoscopelus nicholsi</i>	Species	Fish	3.12E+01	Both	All	14,37
<i>Gymnoscopelus opisthopterus</i>	Species	Fish	3.60E+01	Both	All	5,14
<i>Gymnoscopelus piabilis</i>	Species	Fish	2.50E+01 <sup>A</sup>	North	All Meso + bathypelagic	14
<i>Halicreas minimum</i>	Species	Polychaete	3.70E+00	Both	bathypelagic	2,38
<i>Haloptilus ocellatus</i>	Species	Copepod	6.12E-02 <sup>b,c</sup>	South	Epi + mesopelagic	3,4
<i>Hemiaulus</i>	Genus	Primary producer	2.83E-08 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	18
<i>Heterorhabdus austrinus</i>	Species	Copepod	7.29E-03 <sup>b,c</sup>	Both	Meso + bathypelagic	3,4
<i>Vogtia</i>	Genus	Gelatinous carnivore	2.50E-01 <sup>A</sup>	Both	Meso + bathypelagic	39
<i>Histioteuthis eltaninae</i>	Species	Squid	2.02E+01 <sup>e</sup>	Both <sup>a</sup>	Meso + bathypelagic <sup>a</sup>	15
<i>Hyperia macrocephala</i>	Species	Amphipod	4.20E-01	Both	Epi + mesopelagic	5,26
<i>Hyperiella antarctica</i>	Species	Amphipod	5.70E-03 <sup>A</sup>	Both	Epipelagic	26
<i>Hyperiella dilatata</i>	Species	Amphipod	5.70E-03	Both	Epipelagic	5,26
<i>Ihlea racovitzai</i>	Species	Mesh feeder	7.40E-03	South	All	5,40
<i>Kondakovia longimana</i>	Species	Squid	3.30E+02	Both	All	5,8
<i>Krefflichthys anderssoni</i>	Species	Fish	2.77E+00 <sup>j</sup>	Both	All	8
<i>Laackmanniella</i>	Genus	Protist	1.00E-08	Both	All	21

Lampanyctus	Genus	Fish	2.35E+01 <sup>j</sup>	North	All	14,27	
Lanceola	Genus	Amphipod	2.49E-01 <sup>d</sup>	Both	All	1,41	
<i>Limacina helicina</i>	Species	Meshfeeder	2.72E-05	Both	Epipelagic	1,20	
<i>Limacina retroversa</i>	Species	Meshfeeder	2.72E-05	Both	Epipelagic	1,20	
Lucicutia	Genus	Copepod	8.25E-02 <sup>b,c</sup>	Both	All	3,35	
		Gelatinous					
<i>Marrus antarcticus</i>	Species	carnivore	1.75E-01	Both <sup>a</sup>	All <sup>a</sup>		1
<i>Martialia hyadesi</i>	Species	Squid	3.30E+02	North	All	5,8	
					Meso +		
Mastigoteuthis	Genus	Squid	3.50E+02	Both	bathypelagic		8
Megalocranchia	Genus	Squid	3.50E+02	Both	All	Assumed	
<i>Melanostigma gelatinosum</i>	Species	Fish	2.50E+01	Both	All		13
Melosira	Genus	Primary producer	3.97E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Mesonychoteuthis hamiltoni</i>	Species	Squid	1.10E+05	Both	Bathypelagic	5,8	
<i>Metaconchoecia skogsbergi</i>	Species	Ostracod	3.57E-04 <sup>d</sup>	Both	Mesopelagic	6,7	
					Meso +		
<i>Metridia curticauda</i>	Species	Copepod	5.61E-04 <sup>d</sup>	Both	bathypelagic	3,17	
<i>Metridia gerlachei</i>	Species	Copepod	2.30E-03 <sup>d</sup>	Both	All	3,17	
<i>Metridia lucens</i>	Species	Copepod	3.12E-04 <sup>d</sup>	Both	Epi + mesopelagic	3,17	
<i>Microcalanus pygmaeus</i>	Species	Copepod	1.53E-04 <sup>b,c</sup>	Both	All	3,35	
<i>Microsetella norvegica</i>	Species	Copepod	6.28E-05 <sup>c</sup>	Both	All	17,35	
		Gelatinous					
<i>Muggiaea bargmannae</i>	Species	carnivore	2.50E-01	Both <sup>a</sup>	All <sup>a</sup>		39
<i>Nannobrachium achirus</i>	Species	Fish	2.39E+01 <sup>j</sup>	Both	All	14,27	
					Meso +		
<i>Nansenia antarctica</i>	Species	Fish	1.82E+01 <sup>j</sup>	North	bathypelagic	14,27	
<i>Nematocarcinus lanceopes</i>	Species	Decapod	1.20E+00	Both	All	5,42	
<i>Neocalanus tonsus</i>	Species	Copepod	1.15E-02 <sup>b,c</sup>	North	All	3,4	
Nitzschia	Genus	Primaryproducer	1.99E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Notolepis coatsi</i>	Species	Fish	2.43E+01	Both	Mesopelagic	14,37	

<i>Notoscopelus resplendens</i>	Species	Fish	1.09E+00 <sup>j</sup>	North	All	13,27	
<i>Obtusoecia antarctica</i>	Species	Ostracod	4.51E-04 <sup>d</sup>	North	Mesopelagic	6,7	
<i>Oithona frigida</i>	Species	Copepod	2.64E-05 <sup>c</sup>	Both	Epipelagic	17,43	
<i>Oithona similis</i>	Species	Copepod	2.26E-05 <sup>c</sup>	Both	Epipelagic	17,43	
<i>Oncaea curvata</i>	Species	Copepod	5.67E-05 <sup>b,c</sup>	Both	Epi + mesopelagic	3,17	
<i>Oncaea parila</i>	Species	Copepod	5.40E-05 <sup>b,c</sup>	Both	Meso + bathypelagic	3,17	
<i>Oncaea prolata</i>	Species	Copepod	5.31E-05 <sup>b,c</sup>	Both	Meso + bathypelagic	3,17	
<i>Onykia ingens</i>	Species	Squid	8.33E+02 <sup>d</sup>	North	All		8
<i>Pandea rubra</i>	Species	Gelatinous carnivore	3.00E+00	Both	Meso + bathypelagic	23,38	
<i>Pantachogon haeckeli</i>	Species	Gelatinous carnivore	1.70E+00	Both	All	23,38	
<i>Paradiplospinus gracilis</i>	Species	Fish	1.24E+01	Both	All	1,14	
<i>Paraeuchaeta antarctica</i>	Species	Copepod	7.32E-02 <sup>b,c</sup>	Both	All	3,4	
<i>Paraeuchaeta barbata</i>	Species	Copepod	1.70E-01 <sup>b,c</sup>	Both	Meso + bathypelagic	3,35	
<i>Paraeuchaeta biloba</i>	Species	Copepod	3.64E-02 <sup>b,c</sup>	Both	All	3,4	
<i>Paraeuchaeta rasa</i>	Species	Copepod	3.85E-02 <sup>b,c</sup>	Both	Meso + bathypelagic	3,4	
<i>Parandania boeckii</i>	Species	Copepod	5.00E-01	Both	Meso + bathypelagic	1,2	
<i>Paraphyllina ransoni</i>	Species	Amphipod	3.20E+02	Both <sup>a</sup>	All <sup>a</sup>		1
<i>Pasiphaea scotiae</i>	Species	Gelatinous carnivore	3.68E+00	Both	All	28,42	
<i>Pelagobia longicirrata</i>	Species	Decapod	1.90E-03	Both	All	5,11	
<i>Peridinium</i>	Species	Polychaete	8.36E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		
<i>Periphylla periphylla</i>	Genus	Primaryproducer	1.36E+02	Both <sup>a</sup>	All <sup>a</sup>		1
	Species	Gelatinous carnivore					

<i>Petalidium foliaceum</i>	Species	Decapod	1.16E+00	Both	Meso + bathypelagic	2,28	
Phaeocystis	Genus	Primary producer	1.67E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		44
<i>Phronima sedentaria</i>	Species	Amphipod	8.22E-01 <sup>d</sup>	Both <sup>a</sup>	All <sup>a</sup>		45
Plagiotropus	Genus	Primary producer	1.15E-08 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Pleuragramma antarcticum</i>	Species	Fish	3.00E+02	Both	All	4,5	
		Gelatinous					
Pleurobrachia	Genus	carnivore	1.00E+00 <sup>A</sup>	Both	All		11
<i>Pleuromamma abdominalis</i>	Species	Copepod	5.93E-03 <sup>b,c</sup>	Both	All	3,35	
<i>Pleuromamma antarctica</i>	Species	Copepod	8.26E-03 <sup>b,c</sup>	Both	Epi + mesopelagic	3,35	
					Meso + bathypelagic	3,35	
<i>Pleuromamma robusta</i>	Species	Copepod	7.10E-03 <sup>b,c</sup>	Both	bathypelagic	3,35	
Pleurosigma	Genus	Primary producer	2.17E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
Podocopida	Order	Ostracod	3.29E-04 <sup>A</sup>	Both <sup>a</sup>	All <sup>a</sup>		
					Meso + bathypelagic	13,14	
<i>Poromitra crassiceps</i>	Species	Fish	2.80E+00 <sup>j</sup>	North	bathypelagic	13,14	
Porosira	Genus	Primary producer	1.48E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Primno macropa</i>	Species	Amphipod	2.40E-01	Both	All	5,26	
Prorocentrum	Genus	Primary producer	8.92E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
<i>Protomyctophum andriashevi</i>	Species	Fish	6.25E-01 <sup>j</sup>	North	All	14,46	
<i>Protomyctophum bolini</i>	Species	Fish	5.60E+00	Both	All	5,46	
<i>Protomyctophum choriodon</i>	Species	Fish	3.44E+00 <sup>j</sup>	Both	All	14,27	
<i>Protomyctophum luciferum</i>	Species	Fish	2.02E-01 <sup>j</sup>	North	All	14,46	
<i>Protomyctophum parallelum</i>	Species	Fish	6.42E-01 <sup>j</sup>	North	All	14,27	
<i>Protomyctophum tenisoni</i>	Species	Fish	4.96E-01 <sup>j</sup>	North	All	14,27	
<i>Pseudoamalthothrix cenotelis</i>	Species	Copepod	1.89E-03 <sup>b,c</sup>	Both <sup>a</sup>	All <sup>a</sup>		3
<i>Pseudoconchoecia serrulata</i>	Species	Ostracod	3.96E-04 <sup>d</sup>	North	Mesopelagic	6,7	
Pseudo-nitzschia	Genus	Primary producer	3.51E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>		1
					Meso + bathypelagic	5,8	
<i>Psychroteuthis glacialis</i>	Species	Squid	7.70E+02	Both	bathypelagic	5,8	

Ptychodiscus	Genus	Primary producer Gelatinous	1.63E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	
<i>Pyrostephos vanhoeffeni</i>	Species	carnivore	1.35E-01	Both <sup>a</sup>	All <sup>a</sup>	47
<i>Racovitzanus antarcticus</i>	Species	Copepod	1.87E-03 <sup>b,c</sup>	Both <sup>a</sup>	All <sup>a</sup>	3
Rhabdonella	Genus	Protist	1.00E-08 <sup>A</sup>	North	All	21
<i>Rhincalanus gigas</i>	Species	Copepod	2.00E-02	Both	All	5,17
Rhizosolenia	Genus	Primary producer	1.18E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
<i>Rhynchonereella bongraini</i>	Species	Polychaete Gelatinous	1.90E-03	Both <sup>a</sup>	All <sup>a</sup>	5
<i>Rosacea plicata</i>	Species	carnivore	2.58E-01 <sup>d</sup>	Both	All	23,48
<i>Rotundoechia teretivalvata</i>	Species	Ostracod Gelatinous	7.60E-05 <sup>d</sup>	North	Mesopelagic	6,7
<i>Sagitta gazellae</i>	Species	carnivore Gelatinous	2.34E-01	Both	All	22,49
<i>Sagitta marri</i>	Species	carnivore	1.30E-03	Both	All	5,49
<i>Salpa thompsoni</i>	Species	Meshfeeder	1.40E-01	Both	All	5,50
<i>Scaphocalanus australis</i>	Species	Copepod	6.11E-04 <sup>b,c</sup>	Both <sup>a</sup>	All <sup>a</sup>	3
<i>Scaphocalanus farrani</i>	Species	Copepod	4.35E-03 <sup>b,c</sup>	Both	Meso + bathypelagic	3,4
<i>Scaphocalanus verwoorti</i>	Species	Copepod	1.37E-03 <sup>b,c</sup>	South	All	3,4
Scina	Genus	Amphipod	2.84E-02 <sup>d</sup>	Both	All	26,41
<i>Scolecithricella minor</i>	Species	Copepod	5.77E-04 <sup>b,c</sup>	Both	All	3,35
Sergestes	Genus	Decapod	1.16E+00 <sup>A</sup>	Both	Meso + bathypelagic	2
<i>Slosarczykovia circumantarctica</i>	Species	Squid Gelatinous	1.06E+02 <sup>e</sup>	Both	All	8,15
<i>Solmundella bitentaculata</i>	Species	carnivore	3.30E+00	Both	All	5,11
<i>Spinocalanus abyssalis</i>	Species	Copepod	7.78E-04 <sup>b,c</sup>	Both <sup>a</sup>	Meso + bathypelagic <sup>a</sup>	3
<i>Spongiobranchia australis</i>	Species	Gelatinouscarnivore	1.30E-01	Both	Epipelagic	11,51

<i>Stenosemella</i>	Genus	Protist	1.43E-08	Both <sup>a</sup>	All <sup>a</sup>	52
<i>Stephanopyxis</i>	Genus	Primary producer	1.86E-07 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	18
<i>Stoloteuthis leucoptera</i>	Species	Squid	1.00E-01 <sup>A</sup>	Both <sup>a</sup>	All <sup>a</sup>	
<i>Stomias gracilis</i>	Species	Fish	1.00E+01	Both	All	1,14
		Gelatinous				
<i>Stygiomedusa gigantea</i>	Species	carnivore	6.50E+03	Both <sup>a</sup>	All <sup>a</sup>	1
<i>Subeucalanus longiceps</i>	Species	Copepod	1.71E-02 <sup>b,c</sup>	North	Epi + mesopelagic	3,4
<i>Symbolophorus boops</i>	Species	Fish	4.78E+00 <sup>j</sup>	North	All	13,27
<i>Thalassionema</i>	Genus	Primary producer	2.00E-10 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
<i>Thalassiosira</i>	Genus	Primary producer	9.03E-09 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
<i>Thalassiothrix</i>	Genus	Primary producer	3.71E-08 <sup>a</sup>	Both <sup>a</sup>	Epipelagic <sup>a</sup>	1
<i>Themisto gaudichaudii</i>	Species	Amphipod	1.20E-01	Both	Epi + mesopelagic	5,26
<i>Thysanoessa macrura</i>	Species	Euphausiid	1.50E-01	Both	Mesopelagic	28,31
<i>Thysanoessa vicina</i>	Species	Euphausiid	2.13E-02 <sup>d</sup>	Both	Epi + mesopelagic	5,31
<i>Tintinnopsis</i>	Genus	Protist	1.46E-08	North	All	21,52
<i>Tomopteris planktonis</i>	Species	Polychaete	1.03E-03 <sup>d</sup>	Both	All	54,54
<i>Tomopteris septentrionalis</i>	Species	Polychaete	7.09E-03 <sup>d</sup>	Both	All	53
<i>Triconia antarctica</i>	Species	Copepod	2.79E-04 <sup>b,c</sup>	Both	Epi + mesopelagic	3,17
<i>Vanadis</i>	Genus	Polychaete	1.03E-01 <sup>d</sup>	Both	All	11,53
<i>Vibilia antarctica</i>	Species	Amphipod	1.10E-04	Both	All	5,26
		Gelatinous				
<i>Zanclonia weldoni</i>	Species	carnivore	3.00E+00 <sup>A</sup>	Both	Epipelagic	23

## References for Table S2

[<sup>A</sup>] Assumed

[a] Strathmann RR. (1967). Estimating the organic carbon content of phytoplankton from cell volume or plasma volume. *Limnology and oceanography*, 12(3): 411–418.

[b] Uye SI. (1982). Length- weight relationships of important zooplankton from the Inland Sea of Japan. *Journal of the Oceanographical Society of Japan* 38: 149–158.

[c] Wiebe PH, Boyd S, Cox JL. 1975. Relationships between zooplankton displacement volume, wet weight, dry weight and carbon. *Fisheries Bulletin* 78:777– 786.

[d] Mizdalski E. 1988. Weight and length data of zooplankton in the Weddell Sea in austral spring 1986 (ANT V/3). *Berichte zur Polarforschung*: 55.

[e] Lu CC, Williams R. 1994. Contribution to the biology of squid in the Prydz Bay region, Antarctica. *Antarctic Science* 6: 223–229.

[f] Arkhipkin AI, Bjørke H. 1999. Ontogenetic changes in morphometric and reproductive indices of the squid *Gonatus fabricii* (Oegopsida, Gonatidae) in the Norwegian Sea. *Polar Biology* 22:357–365.

[g] Kaeriyama H, Ikeda T. 2002. Body allometry and developmental characteristics of the three dominant pelagic ostracods (*Discoconchoeciapseudodiscophora*, *Orthoconcheocia haddoni* and *Metaconchoecia skogsbergi*) in the Oyashio region, western North Pacific. *Plankton Biology and Ecology* 49:97–100.

[h] Giussi AR, Sánchez F, Wohler OC, Bernardele JC. 2010. Grenadiers (Pisces, Macrouridae) of the Southwest Atlantic Ocean: biologic and fishery aspects. *Revista de Investigación y Desarrollo Pesquero* 20: 19–33.

- [i] Karuppasamy PK, George S, Menon NG. 2008. Length–weight relationship of *Benthosema pterotum* (myctophid) in the deep scattering layer (DSL) of the eastern Arabian Sea. *Indian Journal of Fisheries* 55: 301–303.
- [j] Linkowski TB. 1985. Population biology of the myctophid fish *Gymnoscopelus nicholsi* (Gillbert, 1911) from the western South Atlantic. *Journal of Fish Biology* 27: 683–698.
- [1] Obtained from BAS database. Unpublished data.
- [2] Piatkowski U, Rodhouse PG, White MG, Bone DG, Symon C. 1994. Nekton community of the Scotia Sea as sampled by the RMT 25 during austral summer. *Marine Ecology Progress Series* 112: 13–28.
- [3] Razouls C, de Bovée F, Kouwenberg J, Desreumaux N. 2018. Diversity and Geographic Distribution of Marine Planktonic Copepods. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed 01/2018]
- [4] Park ET, Ferrari FD. 2009. Species diversity and distributions of pelagic calanoid copepods (Crustacea) from the Southern Ocean. Krupnik I, Lang MA, Miller S, editors. *Smithsonian at the poles: contributions to international polar year science*. Washington, D.C: Smithsonian Institution Scholarly Press. p. 143–180.
- [5] O'Gorman EJ, Yearsley JM, Crowe TP, Emmerson MC, Jacob U, Petchey OL. 2010. Loss of functionally unique species may gradually undermine ecosystems. *Proceedings of the Royal Society of London B: Biological Sciences*: rspb20102036.
- [6] Angel MV, Blachowiak-Samolyk K. 2014. Halocyprid ostracods of the Southern Ocean. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert, Y, editors. *Biogeographic Atlas of the Southern Ocean*. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.



- [7] Blachowiak-Samolyk K, Angel M. 2008. An Atlas of Southern Ocean Planktonic Ostracods. <http://ocean.iopan.gda.pl/ostracoda/index.php> [Accessed 01/ 2018]
- [8] Collins MA, Rodhouse PG. 2006. Southern ocean cephalopods. *Advances in marine biology* 50: 191–265.
- [9] Ikeda T, Bruce B. 1985. Metabolic activity and elemental composition of krill and other zooplankton from Prydz Bay, Antarctica, during early summer (November–December). *Marine Biology* 92: 545–555.
- [10] Ward P. 1984. Aspects of the biology of *Antarctomysis maxima* (Crustacea: Mysidacea). *Polar Biology* 3: 85–92.
- [11] Mackey AP, Atkinson A, Hill SL, Ward P, Cunningham NJ, Johnston NM, Murphy EJ. 2012. Antarctic macrozooplankton of the southwest Atlantic sector and Bellingshausen Sea: Baseline historical distributions (Discovery Investigations, 1928–1935) related to temperature and food, with projections for subsequent ocean warming. *Deep Sea Research Part II: Topical Studies in Oceanography* 59: 130–146.
- [12] Post A. 1990. Paralepididae. In Quéro, J.C., Hureau, C. Karrer, A. Post & Saldanha, L. (Eds.). Check-list of the fishes of the eastern tropical Atlantic (CLOFETA) (Vol. 1). JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. p. 373–384.
- [13] Froese R, Pauly D, editors. 2018. FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org) [Accessed 05/2018]
- [14] Collins MA, Stowasser G, Fielding S, Shreeve R, Xavier JC, Venables HJ, Enderlein P, Cherel, Y, Van de Putte A. 2012. Latitudinal and bathymetric patterns in the distribution and abundance of mesopelagic fish in the Scotia Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 59: 189–198.
- [15] Tree of Life Web Project. <http://tolweb.org> [Accessed 05/2018]

- [16] Sealife Base. [www.sealifebase.com](http://www.sealifebase.com) [Accessed 05/2018]
- [17] Ward P, Atkinson A, Tarling G. 2012. Mesozooplankton community structure and variability in the Scotia Sea: a seasonal comparison. *Deep Sea Research Part II: Topical Studies in Oceanography* 59: 78–92.
- [18] Leblanc K, Arístegui J, Armand L, Assmy P, Beker B, Bode A, Breton E, Cornet V, Gibson J, Gosselin M-P, Kopczynska E, Marshall H, Peloquin J, Piontkovski S, Poulton AJ, Quéginer B, Shiebel R, Shipe R, Stefels J, van Leeuwe MA, Varela M, Widdicombe C, Yallop M. 2012. A global diatom database—abundance, biovolume and biomass in the world ocean. *Earth Systems Science Data* 4: 149–165.
- [19] Rodhouse PG, LuCC. 1998. *Chiroteuthis veranyi* from the Atlantic sector of the Southern Ocean (Cephalopoda: Chiroteuthidae). *South African Journal of Marine Science* 20: 311–322.
- [20] Roberts D, Hopcroft R, Hosie GW. 2014. Southern Ocean Pteropods. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d’Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert Y, editors. *Biogeographic Atlas of the Southern Ocean*. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.
- [21] Thompson GA, Alder VA, Boltovskoy D, Brandini F. 1999. Abundance and biogeography of tintinnids (Ciliophora) and associated microzooplankton in the Southwestern Atlantic Ocean. *Journal of Plankton Research* 21: 1265–1298.
- [22] Parker ML. 2012. Micronekton and Macrozooplankton of the Western Antarctic Peninsula and the Eastern Ross Sea: contrast between two different thermal regimes. PhD Thesis. University of South Florida.
- [23] Lindsay D, Guerrero E, Grossmann M, Fuentes V. 2014. Southern Ocean Gelatinous Zooplankton. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d’Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert

Coudert, Y, editors. Biogeographic Atlas of the Southern Ocean. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

[24] Duhamel G., Hulley P.-A, Causse R., Koubbi P., Vacchi M., Pruvost P., ... Van de Putte A.P. 2014. Biogeographic patterns of fish. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert, Y, editors. Biogeographic Atlas of the Southern Ocean. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

[26] Zeidler W. & De Broyer C. (2014). 6.8. Amphipoda: Hyperiida. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert, Y, editors. Biogeographic Atlas of the Southern Ocean. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

[27] Collins, M. A., Xavier, J. C., Johnston, N. M., North, A. W., Enderlein, P., Tarling, G. A., ... Cunningham, N. J. (2008). Patterns in the distribution of myctophid fish in the northern Scotia Sea ecosystem. *Polar Biology*, 31, 837–851.

[28] Torres, J. J., Aarset, A. V., Donnelly, J., Hopkins, T. L., Lancraft, T. M. & Ainley, D. G. (1994). Metabolism of Antarctic micronektonic Crustacea as a function of depth of occurrence and season. *Marine Ecology Progress Series*, 113, 207–219.

[29] Ward, P. (1989). The distribution of zooplankton in an Antarctic fjord at South Georgia during summer and winter. *Antarctic Science*, 1, 141–150.

[30] Kruse, S., Bathmann, U. & Brey, T. (2009). Meso-and bathypelagic distribution and abundance of chaetognaths in the Atlantic sector of the Southern Ocean. *Polar Biology*, 32, 1359–1376.

[31] Cuzin-Roudy J., Irisson J.-O., Penot F., Kawaguchi S. & Vallet C. (2014). Southern Ocean Euphausiids. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d'Acoz C, Van de

Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert, Y, editors. Biogeographic Atlas of the Southern Ocean. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

[32] Ikeda T. 1988. Metabolism and chemical composition of crustaceans from the Antarctic mesopelagic zone. Deep-Sea Research Part A- Oceanographic Research Papers 35:1991–2002.

[33] Cherel Y, Bocher P, Trouvé C, Weimerskirch H. 2002. Diet and feeding ecology of blue petrels *Halobaenacaerulea* at Iles Kerguelen, Southern Indian Ocean. Marine Ecology Progress Series 228: 283–299.

[34] Metcalfe B. 2013. Planktonic Foraminifera: From production to preservation of the oceanographic signal. PhD Thesis. –Vrije Universiteit Amsterdam.

[35] Kouwenberg JHM., Razouls C, Desreumaux N. 2014. Southern Ocean Pelagic Copepods. De Broyer C, Koubbi P, Griffiths HJ, Raymond B, Udekem d’Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J, Held C, Hosie G, Huettmann F, Post A, Ropert Coudert, Y, editors. Biogeographic Atlas of the Southern Ocean. Cambridge, UK: Scientific Committee on Antarctic Research. p. 297–302.

[36] Jacob U, Brey T, Mintenbeck K. unpublished data. In: Brose U, Jonsson T, Berlow EL, Warren P, Banasek-Richter C, Bersier LF, Blanchard JL, Brey T, Carpenter SR, Cattin B, Landenier M-F, Cushing L, Dawah H.A, Dell T, Edwards F, Harper-Smith S, Jacob U, Ledger ME, Martinez ND, Memmott J, Mintenbeck K, Pinnergar JK, Rall BC, Rayner TS, Ulrich W, Williamns RJ, Woodward G, Cohen JE. (2006). Consumer-Resource body-size relationships in natural food-webs. Ecology 87: 2411–2417.

[37] Artigues B, Morales-Nin B, Balguerías E. 2003. Fish length-weight relationships in the Weddell Sea and Bransfield Strait. Polar Biology 26: 463–467.

- [38] Kawabata T, Lindsay DJ, Kitamura M, Konishi S, Nishikawa J, Nishida S, Kamio M, Nagai H. 2013. Evaluation of the bioactivities of water-soluble extracts from twelve deep-sea jellyfish species. *Fisheries Science* 79: 487–494.
- [39] Pagès F, Kurbjeweit F. 1994. Vertical distribution and abundance of mesoplanktonic medusae and siphonophores from the Weddell Sea, Antarctica. *Polar Biology* 14: 243–251.
- [40] Atkinson A, Ward P, Hunt BPV, Pakhomov EA, Hosie GW. 2012. An overview of Southern Ocean zooplankton data: abundance, biomass, feeding and functional relationships. *CCAMLR Science* 19: 171–218.
- [41] Zeidler W. 1992. Hyperiid amphipods (Crustacea: Amphipoda: Hyperiidae) collected recently from Eastern Australian waters. *Records of the Australian Museum* 44: 85–133.
- [42] Gorny M. 1999. On the biogeography and ecology of the Southern Ocean decapod fauna. *Scientia Marina* 63: 367–382.
- [43] Metz C. 1996. Life strategies of dominant Antarctic Oithonidae (Cyclopoida, Copepoda) and Oncaeidae (Poecilostomatoida, Copepoda) in the Bellingshausen Sea. *Berichte zur Polarforschung* 207: 1–123.
- [44] Peperzak L, Duin R, Colijn F, Gieskes W. 2000. Growth and mortality of flagellates and non-flagellate cells of *Phaeocystis globosa* (Prymnesiophyceae). *Journal of Plankton Research* 22: 107–119.
- [45] Zooplankton Guide. Scripps Institution of Oceanography. <https://scripps.ucsd.edu/zooplanktonguide/> [Accessed 07/2018]
- [46] Pakhomov EA, Perissinotto R, McQuaid CD. 1996. Prey composition and daily rations of myctophid fishes in the Southern Ocean. *Marine Ecology Progress Series* 134: 1–14.
- [47] Mizdalski E. 1988. Weight and length data of zooplankton in the Weddell Sea in austral spring 1986 (ANT V/3). *Berichte zur Polarforschung*, 55.

- [48] Gershwin L, Lewis M, Gowlett-Holmes K, Kloser R. 2014. The Siphonophores. Gershwin L, editor. Pelagic Invertebrates of South-Eastern Australia: A field reference guide. Version 1.1. CSIRO Marine and Atmospheric Research. p. 31.
- [49] Kruse S, Bathmann U, Brey T. 2009. Meso- and bathypelagic distribution and abundance of chaetognaths in the Atlantic sector of the Southern Ocean. *Polar Biology* 32: 1359–1376.
- [50] Atkinson A, Siegel V, Pakhomov E, Rothery P. 2004. Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature* 432: 100.
- [51] Suprenand PM, Ombres EH, Torres JJ. 2015. Metabolism of gymnosomatous pteropods in waters of the western Antarctic Peninsula shelf during austral fall. *Marine Ecology Progress Series* 518: 69–83.
- [52] Capriulo GM, Carpenter EJ. 1983. Abundance, species composition and feeding impact of tintinnid micro-zooplankton in central Long Island Sound. *Marine Ecology Progress Series* 10: 277–288.
- [53] O'Sullivan D. 1983. A guide to pelagic polychaetes of the Southern Ocean and adjacent waters. –ANARE Research Notes, Antarctic Division, Australia.
- [54] Crelier AM, Dadon JR, Isbert-Perlender HG, Nahabedian DE, Daponte MC. 2010. Distribution patterns of chaetognata, polychaeta, pteropoda and salpidae off South Georgia and South Orkney