

Table 5

Median, Interquartile Range, 25th and 75th Percentile for Total Symptom Score and Total Number of Symptoms by Group by Exposure for Open Provocation Trials

	IEI-EMF (n = 102)				Control (n = 237)				IEI-EMF vs. Control		Sham vs. Real		
	Sham	Real	Sham	Real	Sham	Real	Sham	Real	Sham	Real	IEI-EMF	Control	All
	Median (IQR ^a)	[P25, P75 ^b]	Median (IQR)	[P25, P75]	Median (IQR)	[P25, P75]	Median (IQR)	[P25, P75]	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>
Total Symptom Score	2.00 (7.00)	[0.00, 7.00]	5.00 (9.50)	[2.00, 11.50]	0.00 (1.00)	[0.00, 1.00]	0.00 (1.50)	[0.00, 1.50]	<0.001	<0.001	<0.001	0.358	<0.001
Total Number of Symptoms	2.00 (5.25)	[0.00, 5.25]	4.00 (6.63)	[2.00, 8.6.3]	0.00 (1.00)	[0.00, 1.00]	0.00 (1.50)	[0.00, 1.50]	<0.001	<0.001	<0.001	0.438	<0.001

P-values in bold are significant at the corrected level $\alpha = 0.008$ for sham vs. real comparisons and $\alpha = 0.013$ for IEI-EMF vs. control comparisons.

^a Interquartile Range

^b 25th and 75th percentile