

1 **Symbiosis-specific changes in dimethylsulphoniopropionate concentrations in *Stylophora***
2 ***pistillata* along a depth gradient**

3
4

5 Esther M. Borell¹, Daniel T. Pettay², Michael Steinke³, Mark Warner², Maoz Fine^{1,4}

6
7

8 Corresponding author:

9 Esther M. Borell, Email: estherborell@yahoo.co.uk, tel.: 0034 633 735 425

10 ¹ The Interuniversity Institute for Marine Sciences, 88000 Eilat, Israel

11 ² College of Earth, Ocean, and Environment, University of Delaware, Lewes, DE 19958, USA

12 ³ Coral Reef Research Unit, School of Biological Sciences, University of Essex, Colchester CO4
13 3SQ, United Kingdom

14 ⁴ The Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat-Gan
15 52900, Israel

16
17

18
19

20

21 Keywords: Coral transplantation, DMSP, Light, Photoacclimation, *Symbiodinium*

22
23

1 **Abstract**

2 Scleractinian corals are prolific producers of dimethylsulphoniopropionate (DMSP), but
3 ecophysiological mechanisms influencing cellular concentrations are uncertain. DMSP is often
4 proposed to function as an antioxidant but interactions between specific host–symbiont genotype
5 associations, plasticity in DMSP concentrations and environmental conditions that can either
6 exert or alleviate oxidative stress are unclear. We used long-term (6 months) reciprocal
7 transplantation of *Stylophora pistillata* hosting two distinct symbiont phylotypes along a depth
8 gradient, clades A (< 20 m) and C (> 20 m), to assess the effect of change in depth (light
9 intensity) on DMSP concentrations in relation to symbiont genotype and photoacclimation in
10 corals between 3 and 50 m in the Gulf of Aqaba. Bathymetric distribution of total DMSP
11 (DMSPt) per cell varied significantly while particulate DMSP (DMSPp) appeared to be
12 unaffected by depth. Highest DMSPt concentrations in control corals occurred at 20 m. While 3
13 m transplants showed a significant increase in DMSPt concentration at 20 m and became
14 affiliated with an additional genotype (C72), 50 m transplants largely persisted with their original
15 genotype and exhibited no significant changes in DMSPt concentrations. DMSPt concentrations
16 in transplants at both 3 and 50 m, on the other hand, increased significantly while all corals
17 maintained their original symbiont genotypes. Photoacclimation differed significantly with
18 transplantation direction relative to the controls. Symbionts in 3 m transplants at 20 m exhibited
19 no changes in chlorophyll a (chl a) concentration, cell density or cell diameter while symbiont
20 densities decreased and chl a concentrations increased significantly at 50 m. In contrast,
21 symbiont densities in 50 m transplants remained unaffected across depths while symbiont
22 diameters decreased. Chl a concentrations decreased at 20 m and increased at 3 m. Our results

1 indicate that DMSPt concentrations following changes in depth are not only a function of
2 symbiont genotype but result from different acclimation abilities of both symbiotic partners.

3

4 **Introduction**

5 Dimethylsulfoniopropionate (DMSP) is a secondary metabolite that is produced and
6 accumulated at high intracellular concentrations by many marine microalgae (Keller et al. 1989).
7 It is the precursor of dimethylsulfide (DMS), whose oxidation products play a major role in the
8 formation of clouds, cloud albedo, and thus in the regulation of global climate (Vallina and Simó
9 2007). Scleractinian corals maintain an obligate symbiosis with DMSP-producing dinoflagellates
10 in the genus *Symbiodinium* (Hill et al. 1995) and are thought to make a substantial contribution
11 to the amount of DMS entering the atmosphere (Broadbent and Jones 2004; Fischer and Jones
12 2012). However the production of DMSP is complex and cellular concentrations are influenced
13 by a suite of environmental variables including salinity, nutrients, light, and temperature (Stefels
14 2000; Stefels et al. 2007). While the functional role of DMSP in macroalgae and free living
15 phytoplankton is fairly well apprehended (Stefels 2000), fundamental questions remain
16 concerning the biosynthesis, regulation and metabolic roles of DMSP in symbiotic cnidarians
17 (Yost et al. 2012; Raina et al. 2013; Tout et al. 2015). Among various physiological functions
18 (Stefels 2000; Stefels et al. 2007), DMSP and its enzymatic breakdown products are potent
19 scavengers of reactive oxygen species (ROS) (Sunda et al. 2002). Because high concentrations of
20 DMSP were observed in coral host and symbionts following exposure to oxidative stress
21 inducing agents such as light, temperature or copper (Yost et al. 2010; McLenon and DiTullio
22 2012; Deschaseaux et al. 2014a,b) as well as in bleached corals (Jones et al. 2014), DMSP is
23 thought to play an important role in the antioxidant system of the holobiont (the animal and

1 symbiont in combination). Exposure to high levels of visible light and ultraviolet radiation,
2 which in clear water can penetrate to depths of 20 m (Fleischmann 1989), leads to the
3 photodynamic production of ROS within the holobiont (Smith et al. 2005), which is thought to
4 be the principal cause of coral bleaching (Weis 2008). Concentrations of antioxidant defences in
5 corals are therefore often greatest in shallow-water corals and decrease with increasing depth
6 (Shick et al. 1995; Banaszak et al. 1998; Richier et al. 2008).

7 The genus *Symbiodinium* currently comprises nine phylogenetic clades with numerous
8 independently evolving lineages (Sampayo et al. 2009; Pochon and Gates 2010). Each symbiont
9 is adapted to a particular light regime; this is thought to play a major role in structuring host
10 species distributions over reef slopes (Iglesias-Prieto and Trench 1994; Warner and Berry-Lowe
11 2006; Finney et al. 2010) but a corresponding depth-related pattern of DMSP concentrations in
12 different coral host–symbiont genotype associations has not been established so far (Yost et al.
13 2012). Some symbionts exhibit within-genotype variation in DMSP concentrations between
14 different coral hosts (Yost et al. 2012), indicating that DMSP concentrations are a function of
15 both the endosymbiont and coral host. To date, there have been no studies on interactions
16 between specific host–symbiont genotype associations and the plasticity in DMSP concentrations
17 in response to environmental conditions such as light intensity that can either induce or alleviate
18 oxidative stress.

19 The coral *Stylophora pistillata* (Esper, 1797) in the Gulf of Aqaba hosts two distinct
20 *Symbiodinium* from clades A and C, each of which displays well-defined vertical zonation in this
21 region. Deep-water colonies (>20 m) host symbionts belonging to clade C (typical for low
22 irradiance habitats), whereas colonies growing in shallow-water host clade A (typical for high
23 irradiance habitats) (Rowan and Knowlton 1995; Winters et al. 2009). We used reciprocal

1 transplantation of *S. pistillata* along a depth gradient between 3 and 50 m in the Gulf of Aqaba to
2 assess the effect of changes in light intensity on DMSP concentrations in relation to symbiont
3 genotype. Specifically we hypothesized that if transplanted corals remain affiliated with their
4 respective symbiont genotypes, DMSP concentrations in deep-water corals hosting clade C
5 *Symbiodinium* will increase with decreasing depth/increasing light intensity as they will be more
6 sensitive to high light levels than corals hosting clade A *Symbiodinium* and thus be more prone to
7 experience photo-oxidative stress. Conversely, we assumed that DMSP concentrations in
8 shallow-water corals hosting clade A *Symbiodinium* will not be significantly greater than DMSP
9 concentrations in deep-water corals as shallow-water *S. pistillata* in the Gulf of Aqaba is
10 extremely bleaching resistant (Fine et al. 2013) suggesting that these corals are well adapted to
11 high light conditions. DMSP concentrations in shallow-water transplants were expected to
12 remain either unaffected or, because the production of DMSP in corals is thought to be linked to
13 symbiont photosynthesis (Deschaseaux et al. 2014a), to decrease following transplantation to 50
14 m.

15

16 **Materials and methods**

17 **Coral transplantation**

18 Thirty fragments from independent colonies of *S. pistillata* (at least 10 m apart) were collected
19 from depths of 3 and 50 m, and ten fragments were collected from 20 m from the waters in front
20 of the Interuniversity Institute for Marine Sciences (IUI), Eilat, Gulf of Aqaba, Red Sea
21 (29°30'N, 34°56'E) in May 2012 using Trimix SCUBA. Fragments were collected under a
22 special permit from the Israeli Natural Parks Authority. The fragments (~7 cm length) were
23 placed in black labeled plastic bags filled with ambient seawater and transported directly to the

1 laboratory. Under dim light conditions ($< 50 \mu\text{mol m}^{-2} \text{s}^{-1}$), each fragment was tied to a piece of
2 nylon string allowing the fragments to hang freely in the water column. The fragments were
3 labeled to identify colony number and original depth. The following day, all corals were
4 transferred back to the sea to the depth from which they were initially collected and left to
5 recover for one week, after which corals from 3 and 50 m were slowly acclimated to changing
6 light regimes (average light intensities at 3, 20 and 50 m are 1700, 700 and 180 $\mu\text{mol m}^{-2} \text{s}^{-1}$)
7 respectively (Mass et al. 2007).

8 The corals were kept hanging on a horizontal iron cross positioned parallel to the sea floor. The
9 cross, lodged in the ground via a metal pole, protruded 70 cm into the water and corals were
10 hung from each of the four arms 50 cm off the ground with sufficient space between each
11 fragment to ensure that the corals did not touch each other. For the light acclimation a metal
12 cross was installed at 3 m and then at 10 m increments down to 50 m depth.

13 Twenty fragments from 3 m were directly transplanted to a depth of 20 m while 20 fragments
14 from 50 m were transplanted up to 40 m depth where they remained for a week. After that the
15 fragments were reciprocally transplanted in 10 m intervals. Fragments from 3 m were acclimated
16 to low light conditions over a period of 8 d, remaining at 30 and 40 m for 4 d respectively.

17 Fragments from 50 m depth were acclimated to higher light conditions over a period of 24 d to
18 avoid excessive stress or mortality of the corals as has been observed previously in ‘upward’
19 transplantations due to the sensitivity to acute high light exposure for many corals collected from
20 low light conditions (Baker 2001; Cohen and Dubinsky 2015). Specifically, corals were
21 acclimated to 30 and 20 m for 4 d each, after which they were acclimated to 10 m and finally to 3
22 m over 8 d at each depth.

1 After reciprocal transplantation was complete, coral fragments at 3 m comprised ten control
2 fragments and ten transplanted fragments from 50 m, corals at 20 m comprised ten control corals
3 from 20 m, ten transplants from 3 m and ten transplants from 50 m, and fragments at 50 m
4 consisted of ten controls from 50 m and ten transplants from 3m. Algae were cleaned from the
5 iron frames and nylon strings every fortnight. No algal growth occurred on the corals. All
6 fragments were kept in the sea for 6 months until the end of November 2013, after which they
7 were transported to the lab using black plastic bags with ambient seawater. Once ashore, the
8 fragments were immediately frozen in liquid nitrogen and then stored at -80 °C pending tissue
9 analyses.

10

11 **Sample processing**

12 Coral tissues were stripped from the skeletons under dim light using an airbrush and sterile
13 filtered seawater (FSW, 0.2 µm) to obtain a slurry that was collected in a 5 mL Falcon tube. The
14 slurry (20–50 mL) was homogenized for 10 s with an electric homogenizer (DIAX 100
15 homogenizer Heidolph Instruments GmbH & Co. KG, Schwabach, Germany). Sub-samples were
16 immediately removed for analyses of multiple indices using sterile transfer pipettes. We
17 measured algal genotype, total DMSP (DMSPt = sum of algal and host DMSP) and particulate
18 DMSP (DMSPp = algal DMSP), chlorophyll a (chl a), symbiont densities and cell sizes
19 (diameter). The remaining coral skeleton was saved for surface area measurements using the
20 paraffin wax technique (Stimson and Kinzie 1991).

21

22 **Analyses of algal indices and algal genotype**

1 To quantify symbiont numbers, diameters and chl a concentrations, 2 mL of homogenate was
2 centrifuged (1900 g at 4 °C) and the algal pellets resuspended three times in FSW. Resuspended
3 algae were used for chl a extraction in acetone (100%) at 4 °C in the dark for 24 h.
4 Concentrations were determined spectrophotometrically (Jeffrey and Humphrey 1975).
5 Symbiont densities were quantified from four replicate counts using a Neubauer hemocytometer.
6 Densities and chl a were normalized to coral surface area. Cell diameters were determined
7 microscopically with an eyepiece graticule from 30 replicate measurements per sample.

8 Nucleic acid extractions from five randomly chosen replicates were conducted using a
9 modified Promega Wizard genomic DNA extraction protocol (LaJeunesse et al. 2003). Symbiont
10 identity was characterized by denaturing gradient gel electrophoresis (DGGE) fingerprinting of
11 the partial 5.8S and internal transcribed spacer region 2 (ITS2; LaJeunesse 2002). The region
12 was amplified using a touch-down thermal cycle profile with the primers “ITS2clamp” and
13 “ITSintfor2” (Lajeunesse and Trench 2000), and the PCR products resolved on denaturing gels
14 (45–80% of 7 mol L⁻¹ urea and 40% formamide) using a CBS Scientific system (Del Mar, CA)
15 for 16 h at 95 V. The dominant band of the symbiont’s DGGE profile was excised, reamplified,
16 and cycle-sequenced to provide the ITS2 sequence that dominates the symbiont’s genome.

17

18 **DMSPt and DMSPp analyses**

19 For the quantification of DMSPt, 1 mL of homogenate was added to 2 mL 0.5 mol L⁻¹ NaOH in
20 a gas-tight, screw-cap headspace vial. To isolate the algal component (DMSPp) from the
21 homogenate, 1 mL of each sample homogenate was centrifuged (2900 g at 4 °C) for 20 min, the
22 pellets resuspended in 1 mL FSW and immediately added to 2 mL 0.5 mol L⁻¹ NaOH in a
23 headspace vial. The supernatant of the control corals from 3 and 20 m was transferred into

1 another headspace vial containing 0.5 mol L^{-1} NaOH to analyze DMSP concentrations in the host
2 tissue (DMSP_h). DMSP_t concentrations were compared to the sum of DMSP_p and DMSP_h
3 concentrations to account for the potential loss of DMSP due to the conversion to DMS during
4 centrifugation and handling. The addition of NaOH produces an alkaline hydrolysis that rapidly
5 converts DMSP to equimolar concentrations of DMS, which can be quantified using gas
6 chromatographic methods with direct injection of headspace (Steinke et al. 2011). Results were
7 expressed as femtomole DMSP_t and DMSP_p per symbiont cell and as nmol cm^{-2} surface area.

8

9 **Data analyses**

10 Data were checked for homogeneity of variances using the Cochran's C-test and $\ln(x)$ -
11 transformed if necessary before using one-way ANOVA. Student–Newman–Keuls (SNK) tests
12 were used for post hoc multiple comparisons. At the end of the experiment two fragments were
13 missing from the control corals at 20 m and one fragment from the control corals at 3 m. To
14 avoid confounding effects due to unbalanced sample sizes all measured variables were analyzed
15 using eight replicates. Data were analyzed using WinGMAV (EICC, University of Sydney,
16 Australia). An additional regression analysis to test for correlations between DMSP_p and DMSP_t
17 (normalized to surface area) and symbiont densities was performed and analyzed by ANOVA
18 using JMP Pro 10.0.2 (SAS).

19

20 **Results**

21 ***Symbiodinium* genotypes in *S. pistillata***

22 Three distinct symbionts belonging to clades A and C were found in *S. pistillata* fragments that
23 were maintained at their native depths (Table 1). *Symbiodinium* A1 dominated all fragments at 3

1 and 20 m depths, while C167a and C168a were found in the fragments at 50 m. Except for some
2 transplanted corals at 20 m, all transplanted fragments persisted in association with symbionts
3 corresponding to their original collection depth. Three fragments transplanted from 3 to 20m
4 possessed C72a-b, while one fragment transplanted from 50 to 20m possessed C169a-b.

6 **Photoacclimation of *Symbiodinium* along the depth gradient**

7 Symbiont densities, chl a concentrations and cell diameters in control corals and coral transplants
8 varied significantly with depth (Table 2). Algae in control corals at 3 and 20 m occurred at
9 similar densities but displayed a significant decrease at 50 m (Fig. 1a). This trend was paralleled
10 by cell densities in corals transplanted from 3 to 20 and 50 m. By contrast, change in depth had
11 no effect on symbiont densities in 50 m transplants. Chl a concentrations in control corals did not
12 vary significantly between depths (Fig. 1b). Concentrations in 3 m transplants at 20 m were
13 similar to the control corals while concentrations in 3 m transplants at 50 m exhibited a
14 significant increase. Chl a concentrations in 50 m transplants changed significantly at both
15 depths, with an increase in concentration at 3 m and a decrease at 20 m. The symbiont cell
16 diameter in control corals significantly increased with increasing depth (Fig. 2).
17 Correspondingly, algae in 50 m transplants exhibited a decrease in diameter with decreasing
18 depth. By contrast, the cell diameter of symbionts in 3 m transplants remained unaffected by
19 changes in depth.

20 **DMSP concentrations**

21 Potential loss of DMSPp due to centrifugation and handling was insignificant (Table 3).
22 The bathymetric distribution of DMSPp and DMSPt varied significantly depending on the
23 normalization index used (Table 2). Concentrations of DMSPt normalized to algal cell in control

1 corals were significantly higher at 20 m than at 3 and 50 m (Fig. 3a). DMSPt in 3 m transplants
2 increased significantly at both 20 and 50 m relative to the control corals. Likewise there was a
3 significant increase in DMSPt concentrations in 50 m transplants at 3 m. However DMSPt
4 concentrations at 20 m were not significantly different from the controls at 50 m (Fig. 3a).
5 DMSPt normalized to surface area exhibited a similar trend to that of symbiont densities (Fig.
6 3b). DMSPt concentrations in control corals at 3 and 20m were significantly higher than control
7 corals at 50 m with the highest concentrations occurring at 20 m. DMSPt concentrations in 3m
8 transplants paralleled those of the control corals while there were no significant differences in
9 DMSPt between depths for 50 m transplants. No significant effects were detected for
10 concentrations of DMSPP normalized to cell (Fig. 4a). In contrast, DMSPP normalized to surface
11 area followed the pattern of the depth distribution of symbiont densities (Fig. 4b). Regression
12 analyses revealed a significant correlation between DMSPt (normalised to surface area) ($r^2 =$
13 0.60, ANOVA, $F_{1,55} = 80.00$, $P < 0.01$; Fig. 5a) and DMSPP (normalized to surface area) ($r^2 =$
14 0.52, ANOVA, $F_{1,55} = 57.72$, $P < 0.01$; Fig. 5b) and symbiont density.

15

16 **Discussion**

17 **DMSP concentration and normalization index**

18 To date, there is no clear consensus on which index best conveys DMSP concentrations in corals.
19 Concentrations are frequently normalized to multiple indices, each of which can reveal a
20 different mechanistic process that may influence DMSP concentrations in the holobiont (Yost et
21 al. 2012; Deschaseaux et al. 2014a). Normalization to surface area can capture potential effects
22 of symbiont densities. Normalization to cell, on the other hand, can reflect the ecophysiological
23 effects on DMSP concentrations in the holobiont. Although DMSP is commonly perceived to be

1 produced by the algal symbionts (Van Alstyne et al. 2009), the location of DMSP production
2 within the coral remains ambiguous. While concentrations of DMSP are found in the tissue of
3 the host implying that DMSP either leaks or is translocated from symbiont to host (Yost et al.
4 2012), there is also evidence showing that juvenile corals lacking symbionts are able to
5 synthesize DMSP directly (Raina et al. 2013). Our results show that regardless of the
6 normalization index used, concentrations of DMSPt were consistently greater than
7 concentrations of DMSPp, consistent with previous studies (Yost et al. 2010, 2012). The
8 regression analyses, together with the fact that the bathymetric distribution pattern of DMSPt
9 and DMSPp normalized to surface area closely paralleled that of the symbiont densities, indicate
10 that accumulation of DMSP in the host tissue occurred in a cell density-dependent fashion.
11 Normalization of the data to algal cell on the other hand demonstrated a clear effect of depth on
12 DMSPt concentrations in *S. pistillata*, while DMSPp concentrations appeared to be unaffected by
13 depth. This suggests that DMSP translocation from symbiont to host may be to some extent
14 mediated by the coral animal. DMSPt and DMSPp in the following discussion refer to
15 concentrations normalized to cell.

16

17 **Depth distribution of DMSP**

18 The depth distribution of DMSPt in control corals was in line with our predictions, showing that
19 concentrations in corals at 3 m hosting clade A symbionts and in corals at 50 m hosting clade C
20 symbionts did not differ significantly. In contrast, the high DMSPt concentrations measured in
21 both the control corals and 3 m transplants at 20 m were unexpected.

22 Combined with chl a concentrations and symbiont densities of the control corals at 3 m, which
23 provided no signs of bleaching, the data indicate that *S. pistillata* at 3 m was well adapted to high

1 levels of irradiance supporting the tenet of high bleaching resistance of this species in our study
2 area (Fine et al. 2013). Compared to the shallow-water light environment, corals at 20 m are
3 exposed to enormous variation in light intensity (Dishon et al. 2012).
4 Such fluctuations may have rendered corals at this depth more susceptible to stress than shallow-
5 water corals that were exposed to high but continuous levels of irradiance (Allahverdiyeva et al.
6 2015). Further indication that corals at 20 m experienced more fluctuation than those at 3 m is
7 suggested by the fact that chl a concentrations in 50 m transplants were significantly lower at 20
8 m than at 3 m, and by the emergence of additional genotypes (Jones et al. 2008) in coral
9 transplants which only occurred at this depth while transplants at 3 and 50 m maintained their
10 original genotypes.

11

12 **DMSP distribution in coral transplants and bathymetrically-driven association with** 13 ***Symbiodinium***

14 DMSPt concentrations displayed a significant increase relative to the controls at each
15 transplantation depth in all but deep-water transplants at 20 m. While increases in DMSPt at both
16 3 and 50 m occurred independently of symbiont genotype, high DMSPt concentrations in 3 m
17 transplants at 20 m coincided with the emergence of an additional genotype. The presence of
18 C72-a-b symbiont genotypes in the 3 m transplants is consistent with previous observations of
19 the alternation and co-occurrence of symbiont types C72 and A1 in *S. pistillata* at this depth in
20 the Gulf of Aqaba (Byler et al. 2013). Against this background it is intriguing that the 50 m
21 transplants at 20 m exhibited no increase in DMSPt concentrations and remained largely
22 affiliated with the same types of *Symbiodinium* as their 50 m controls. The relevance of the

1 emergence of the genotype C169-a-b in 50 m transplants remains to be determined, as it has not
2 been observed in *S. pistillata* before and did not appear to be very abundant.

3 One reason for the contrasting DMSPt concentrations at 20 m may be related to intrinsic
4 differences in the capacities to acclimatize to changing light levels between shallow- and deep-
5 water holobionts. This is supported by the variation in symbiont population characteristics
6 relative to the control corals at each transplantation depth. While shallow-water transplants at 20
7 m exhibited no changes in chl a concentration, cell density or cell diameter, symbionts in deep-
8 water transplants decreased in cell diameter (hence volume) and contained significantly less chl a
9 relative to the control corals. Differential acclimation abilities of coral hosts may have also
10 accounted for the symbiont recombination in the 3 m transplants at 20 m relative to the control
11 corals at 20 m, which were affiliated with *Symbiodinium* type A1 only. This is consistent with
12 previous observations indicating that the ability of the algal symbionts to acclimate to specific
13 light regimes does not occur in isolation but is influenced by morphological and/or physiological
14 constraints of the coral host, which in turn affects physiological responses and thus relative stress
15 tolerance of the holobiont (Goulet et al. 2005; Frade et al. 2008a,b). The fact that DMSPP did not
16 change as a function of depth together with the observed variation in photoacclimation responses
17 indicate that DMSPt concentrations were to some extent regulated by the host. Indeed, this
18 agrees with observations by Yost et al. (2012) showing that DMSPt concentrations varied
19 between different coral species hosting the same symbiont genotype.

20 Diverging photoacclimation strategies were also evident in coral transplants at 3 and 50
21 m, which maintained their original symbiont genotypes. The photoacclimation response of
22 shallow-water transplants hosting clade A symbiont genotypes was similar to that of the control
23 corals showing a classic pattern of photoacclimation to low light levels with a decrease in

1 symbiont numbers (Dustan 1982) and concomitant increase in chl a concentrations (Falkowski
2 and Dubinsky 1981; Cohen and Dubinsky 2015) corroborating other observations for *S. pistillata*
3 along the same depth gradient in the Gulf of Aqaba (Mass et al. 2007; Winters et al. 2009; Cohen
4 and Dubinsky 2015). In contrast, in the current study, deep-water corals hosting clade C78 and
5 C161a symbionts responded to high light intensity with a complete switch to C161a at 3 m. A
6 switch in the relative dominance of genotype, referred to as symbiont shuffling, is thought to
7 present an acclimation mechanism in response to changes in environmental conditions (Jones et
8 al. 2008) as different genotypes vary markedly in their photophysiological characteristics
9 (Iglesias-Prieto et al. 2004; Frade et al. 2008b). In contrast to 3 m transplants, photoacclimation
10 in 50 m transplants did not entail changes in symbiont densities but a decrease in cell diameter
11 and curiously, an increase in chl a concentrations at 3 m.

12 While DMSPt concentrations in deep-water transplants at 3 m conform to the notion that
13 corals hosting clade C were more sensitive to high light than shallow-water controls hosting
14 clade A, high chl a concentrations in these corals rebut this suggestion. A lesser-known function
15 for DMSp, proposed by Stefels (2000), is that of an overflow mechanism for excess reduced
16 sulphur and as a means to dissipate excess energy under conditions where carbon incorporation
17 exceeds the rate of protein synthesis, a common phenomenon in shallow-water corals (Falkowski
18 et al. 1984). Although copious amounts of fixed carbon are translocated from symbiont to host,
19 very little of it is assimilated as the translocated products are deficient in nitrogen, which
20 necessitates release of large quantities of carbon to the environment (Falkowski et al. 1984).
21 Carbon translocation and secretion, however, differ with environmental conditions (Falkowski et
22 al. 1984; Tremblay et al. 2014), as well as between host or symbiont genotypes (Davy et al.
23 1996; Loram et al. 2007). Differential amounts of released carbon due to specific symbiont–host

1 interactions and distinct photophysiological characteristics could therefore account for the high
2 concentrations of DMSP in deep-water corals at 3 m relative to the controls.

3 An intriguing aspect of our results was the significant increase in DMSPt concentrations in
4 shallow-water corals transplanted to 50 m which contrasted with our assumption that DMSP
5 concentrations would decrease with a decrease in the potential to experience photo-oxidative
6 stress (e.g., Shick et al. 1995). Likewise, our results contradict findings by Deschaseaux et al.
7 (2014b) who observed no significant effect of light depletion on DMSPt concentrations over the
8 course of 3 d. One possibility is that DMSPt concentrations in shallow-water transplants in our
9 study did not change at 50 m and that the observed increase in DMSPt is merely an artifact
10 resulting from a decrease in symbiont densities in transplants at 50 m. Although the increase in
11 DMSPt in our study coincided with a significant increase in chl a concentration, it seems
12 unlikely that these were linked to the high levels of DMSPt since, despite photoacclimation, the
13 photosynthetic efficiency of symbionts generally decreases significantly with depth (Lesser et al.
14 2000; Titlyanov et al. 2001; Mass et al. 2007). An alternative explanation may be that the
15 physiological function of DMSP may vary with depth. An important metabolic function of
16 DMSP in both algae (Kirst 1996) and symbiotic cnidarians (Yancey et al. 2010) is that of an
17 osmolyte. Though not very well investigated as osmoconformers, corals maintain high levels of
18 compatible solutes (Stefels 2000; Mayfield and Gates 2007). However, the distribution of these
19 solutes, including DMSP, is likely to change with depth because rates of carbon assimilation and
20 quality of photoassimilates are not uniform across different depths (Falkowski et al. 1984;
21 Muscatine et al. 1984; Alamaru et al. 2009), thus leading to changes in the relative concentration
22 of each solute (Stefels 2000).

23 Regardless of the metabolic role of DMSP in *S. pistillata*, our results indicate that DMSPt

1 concentrations following changes in light intensity are not only a function of symbiont genotype
2 but result from complex interactions between both symbiotic partners. As coral reefs are
3 subjected to enormous environmental pressures via pollution (Fabricius 2005) and climate
4 change (Hoegh-Guldberg et al. 2007), which influence light levels (Wooldridge 2008;
5 Reopanichkul et al. 2009), symbiont population characteristics and genetic composition (Baker
6 et al. 2008; Jones et al. 2008), our data present important considerations for ongoing efforts to
7 elucidate how environmental variability can influence plasticity in DMSP concentrations in the
8 coral holobiont.

9

10

11 **Acknowledgements**

12 We thank Guy Levy, Genadi Salzman and Oded Ben-Shaprut for help with the transplantation
13 experiments and are grateful for the continuous support of the IUI staff. Thanks to Tracy Lawson
14 for assistance with the preparation of the figures. Much appreciated are the constructive
15 comments of two anonymous reviewers, which significantly improved this manuscript. EMB
16 was funded by the Minerva fellowship program, DTP was supported by the NSF and MS
17 financial support to MF was provided by the EU MedSeA project.

18

19 **References**

20 Alamaru A, Loya Y, Brokovich E, Yam R, Shemesh A (2009) Carbon and nitrogen utilization in
21 two species of Red Sea corals along a depth gradient: insights from stable isotope analysis
22 of total organic material and lipids. *Geochim Cosmochim Acta* 73:5333–5342
23 Allahverdiyeva Y, Suorsa M, Tikkanen M, Aro E-M (2015) Photoprotection of photosystems in
24 fluctuating light intensities. *J Exp Bot* 66:2427–2436
25 Baker AC (2001) Ecosystems: reef corals bleach to survive change. *Nature* 411:765–766

- 1 Baker AC, Glynn PW, Riegel B (2008) Climate change and coral reef bleaching: an ecological
2 assessment of long-term impacts, recovery trends and future outlook. *Estuar Coast Shelf*
3 *Sci* 80:435–471
- 4 Banaszak AT, Lesser MP, Kuffner IB, Ondrusek M (1998) Relationship between ultraviolet
5 (UV) radiation and mycosporine-like amino acids (MAAs) in marine organisms. *Bull Mar*
6 *Sci* 63:617–628
- 7 Broadbent AD, Jones GB (2004) DMS and DMSP in mucus ropes, coral mucus, surface films
8 and sediment pore waters from coral reefs in the Great Barrier Reef. *Mar Freshw Res*
9 55:849–855
- 10 Byler KA, Carmi-Veal M, Fine M, Goulet TL (2013) Multiple symbiont acquisition strategies as
11 an adaptive mechanism in the coral *Stylophora pistillata*. *PLoS One* 8:e59596
- 12 Cohen I, Dubinsky Z (2015) Long term photoacclimation responses of the coral *Stylophora*
13 *pistillata* to reciprocal deep to shallow transplantation: photosynthesis and calcification.
14 *Front Mar Sci* 2:45
- 15 Davy SK, Lucas IA, Turner JR (1996) Carbon budgets in temperate anthozoan-dinoflagellate
16 symbioses. *Marine Biology* 126:773–783
- 17 Deschaseaux ESM, Beltran VH, Jones GB, Deseo MA, Swan HB, Harrison PL, Eyre BD
18 (2014a) Comparative response of DMS and DMSP concentrations in *Symbiodinium* clades
19 C1 and D1 under thermal stress. *J Exp Mar Bio Ecol* 459:181–189
- 20 Deschaseaux ESM, Jones GB, Deseo MA, Shepherd KM, Kiene RP, Swan HB, Harrison PL,
21 Eyre BD (2014b) Effects of environmental factors on dimethylated sulfur compounds and
22 their potential role in the antioxidant system of the coral holobiont. *Limnol Oceanogr*
23 59:758–768
- 24 Dishon G, Dubinsky Z, Fine M, Iluz D (2012) Underwater light field patterns in subtropical
25 coastal waters: a case study from the Gulf of Eilat (Aqaba). *Isr J Plant Sci* 60:265–275
- 26 Dustan P (1982) Depth-dependent photoadaptation by zooxanthellae of the reef coral *Montastrea*
27 *annularis*. *Mar Biol* 68:253–264
- 28 Fabricius KE (2005) Effects of terrestrial runoff on the ecology of corals and coral reefs: review
29 and synthesis. *Mar Pollut Bull* 50:125–146
- 30 Falkowski PG, Dubinsky Z (1981) Light-shade adaptation of *Stylophora pistillata*, a hermatypic
31 coral from the Gulf of Eilat. *Nature* 289:172–174
- 32 Falkowski PG, Dubinsky Z, Muscatine L, Porter JW (1984) Light and the bioenergetics of a
33 symbiotic coral. *Bioscience* 34:705–709
- 34 Fine M, Gildor H, Genin A (2013) A coral reef refuge in the Red Sea. *Glob Chang Biol*
35 19:3640–3647
- 36 Finney JC, Pettay DT, Sampayo EM, Warner ME, Oxenford HA, LaJeunesse TC (2010) The
37 relative significance of host-habitat, depth, and geography on the ecology, endemism, and
38 speciation of coral endosymbionts in the genus *Symbiodinium*. *Microb Ecol* 60:250–263
- 39 Fischer E, Jones G (2012) Atmospheric dimethylsulphide production from corals in the Great
40 Barrier Reef and links to solar radiation, climate and coral bleaching. *Biogeochemistry*
41 110:31–46
- 42 Fleischmann EM (1989) The measurement and penetration of ultraviolet radiation into tropical
43 marine water. *Limnol Oceanogr* 34:1623–1629
- 44 Frade PR, Bongaerts P, Winkelhagen AJS, Tonk L, Bak RPM (2008a) In situ photobiology of
45 corals over large depth ranges: a multivariate analysis on the roles of environment, host,
46 and algal symbiont. *Limnol Oceanogr* 53:2711–2723

- 1 Frade PR, De Jongh F, Vermeulen F, Van Bleijswijk J, Bak RPM (2008b) Variation in symbiont
2 distribution between closely related coral species over large depth ranges. *Mol Ecol*
3 17:691–703
- 4 Goulet TL, Cook CB, Goulet D (2005) Effect of short-term exposure to elevated temperatures
5 and light levels on photosynthesis of different host–symbiont combinations in the *Aiptasia*
6 *pallida/Symbiodinium* symbiosis. *Limnol Oceanogr* 50:1490–1498
- 7 Hill RW, Dacey JWH, Krupp DA (1995) Dimethylsulfoniopropionate in reef corals. *Bull Mar*
8 *Sci* 57:489–494
- 9 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD,
10 Sale PF, Edwards AJ, Caldeira K (2007) Coral reefs under rapid climate change and ocean
11 acidification. *Science* 318:1737–1742
- 12 Iglesias-Prieto R, Trench RK (1994) Acclimation and adaptation to irradiance in symbiotic
13 dinoflagellates. I. Responses of the photosynthetic unit to changes in photon flux density.
14 *Mar Ecol Prog Ser* 113:163–175
- 15 Iglesias-Prieto R, Beltrán VH, LaJeunesse TC, Reyes-Bonilla H, Thomé PE (2004) Different
16 algal symbionts explain the vertical distribution of dominant reef corals in the eastern
17 Pacific. *Proc R Soc Lond B Biol Sci* 271:1757–1763
- 18 Jeffrey SW, Humphrey GF (1975) New spectrophotometric equations for determining
19 chlorophylls *a*, *b*, *c*₁ and *c*₂ in higher plants, algae and natural phytoplankton. *Biochem*
20 *Physiol Pflanz* 167:191–194
- 21 Jones AM, Berkelmans R, van Oppen M.J.H, Mieog J.C, Sinclair W (2008) A community change
22 in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field
23 evidence of acclimatization. *Proc R Soc Lond B Biol Sci* 275:1359–1365
- 24 Jones GB, Fischer E, Deschaseaux ESM, Harrison PL (2014) The effect of coral bleaching on
25 the cellular concentration of dimethylsulphoniopropionate in reef corals. *J Exp Mar Bio*
26 *Ecol* 460:19–31
- 27 Keller MD, Bellows WK, Guillard RRL (1989) Dimethyl sulfide production in marine
28 phytoplankton. In: Saltzman ES, Cooper WJ (eds.) *Biogenic sulfur in the environment*.
29 American Chemical Society Washington DC, pp 167–182
- 30 Kirst GO (1996) Osmotic adjustment in phytoplankton and macroalgae. In: Kiene R, Visscher P,
31 Keller M, Kirst G, Kirst GO (eds) *Biological and environmental chemistry of DMSP and*
32 *related sulfonium compounds*. Springer US, pp 121–129
- 33 LaJeunesse T (2002) Diversity and community structure of symbiotic dinoflagellates from
34 Caribbean coral reefs. *Mar Biol* 141:387–400
- 35 Lajeunesse TC, Trench RK (2000) Biogeography of two species of *Symbiodinium* (Freudenthal)
36 inhabiting the intertidal sea anemone *Anthopleura elegantissima* (Brandt). *Biol Bull*
37 199:126–134
- 38 LaJeunesse TC, Loh WKW, Van Woesik R, Hoegh-Guldberg O, Schmidt GW, Fitt WK (2003)
39 Low symbiont diversity in southern Great Barrier Reef corals, relative to those of the
40 Caribbean. *Limnol Oceanogr* 48:2046–2054
- 41 Lesser MPL, Mazel C, Phinney D, Yentsch CS (2000) Light absorption and utilization by
42 colonies of the congeneric hermatypic corals *Montastraea faveolata* and *Montastraea*
43 *cavernosa*. *Limnol Oceanogr* 45:76–86
- 44 Loram JE, Trapido-Rosenthal HG, Douglas AE (2007) Functional significance of genetically
45 different symbiotic algae *Symbiodinium* in a coral reef symbiosis. *Molecular Ecology*
46 16:4849–4857

- 1 Mass T, Einbinder S, Brokovich E, Shashar N, Vago R, Erez J, Dubinsky Z (2007)
2 Photoacclimation of *Stylophora pistillata* to light extremes: metabolism and calcification.
3 Mar Ecol Prog Ser 334:93–102
- 4 Mayfield AB, Gates RD (2007) Osmoregulation in anthozoan–dinoflagellate symbiosis. Comp
5 Biochem Physiol A Mol Integr Physiol 147:1–10
- 6 McLendon AL, DiTullio GR (2012) Effects of increased temperature on
7 dimethylsulfoniopropionate (DMSP) concentration and methionine synthase activity in
8 *Symbiodinium microadriaticum*. Biogeochemistry 110:17–29
- 9 Muscatine L, Falkowski PG, Porter JW, Dubinsky Z (1984) Fate of photosynthetic fixed carbon
10 in light- and shade-adapted colonies of the symbiotic coral *Stylophora pistillata*. Proc R
11 Soc Lond B Biol Sci 222:181–202
- 12 Pochon X, Gates RD (2010) A new *Symbiodinium* clade (Dinophyceae) from soritid foraminifera
13 in Hawai'i. Mol Phylogenet Evol 56:492–497
- 14 Raina J-B, Tapiolas DM, Forêt S, Lutz A, Abrego D, Ceh J, Seneca FO, Clode PL, Bourne DG,
15 Willis BL (2013) DMSP biosynthesis by an animal and its role in coral thermal stress
16 response. Nature 502:677–680
- 17 Reopanichkul, P, Schlacher TA, Carter RW, Worachananant S (2009) Sewage impacts coral
18 reefs at multiple levels of ecological organization. Mar Pollut Bull 58:1356–1362
- 19 Richier S, Cottalorda J-M, Guillaume MMM, Fernandez C, Allemand D, Furla P (2008) Depth-
20 dependant response to light of the reef building coral, *Pocillopora verrucosa*: implication
21 of oxidative stress. J Exp Mar Bio Ecol 357:48–56
- 22 Rowan R, Knowlton N (1995) Intraspecific diversity and ecological zonation in coral–algal
23 symbiosis. Proc R Soc Lond B Biol Sci 92:2850–2853
- 24 Sampayo EM, Dove S, LaJeunesse TC (2009) Cohesive molecular genetic data delineate species
25 diversity in the dinoflagellate genus *Symbiodinium*. Mol Ecol 18:500–519
- 26 Shick JM, Lesser MP, Dunlap WC, Stochaj WR, Chalker BE, Won JW (1995) Depth-dependent
27 responses to solar ultraviolet radiation and oxidative stress in the zooxanthellate coral
28 *Acropora microphthalma*. Mar Biol 122:41–51
- 29 Smith DJ, Suggett DJ, Baker NR (2005) Is photoinhibition of zooxanthellae photosynthesis the
30 primary cause of thermal bleaching in corals? Glob Chang Biol 11:1–11
- 31 Stefels J (2000) Physiological aspects of the production and conversion of DMSP in marine
32 algae and higher plants. Netherlands Journal of Sea Research 43:183–197
- 33 Stefels J, Steinke M, Turner S, Malin G, Belviso S (2007) Environmental constraints on the
34 production and removal of the climatically active gas dimethylsulphide (DMS) and
35 implications for ecosystem modelling.. Biogeochemistry 83:245–275
- 36 Steinke M, Brading P, Kerrison P, Warner ME, Suggett DJ (2011) Concentrations of
37 dimethylsulfoniopropionate and dimethyl sulfide are strain-specific in symbiotic
38 dinoflagellates (*Symbiodinium* sp., Dinophyceae). J Phycol 47:775–783
- 39 Stimson J, Kinzie RA (1991) The temporal release of zooxanthellae from the reef coral
40 *Pocillopora damicornis* (Linnaeus) under nitrogen-enrichment and control conditions. J
41 Exp Mar Bio Ecol 153:63–74
- 42 Sunda W, Kieber DJ, Kiene RP, Huntsman S (2002) An antioxidant function for DMSP and
43 DMS in marine algae. Nature 418:317–320
- 44 Titlyanov EA, Titlyanova TV, Yamazato K, Van Woesik R (2001) Photo-acclimation dynamics
45 of the coral *Stylophora pistillata* to low and extremely low light. J Exp Mar Bio Ecol
46 263:211–225

- 1 Tout J, Jeffries TC, Petrou K, Tyson GW, Webster NS, Garren M, Stocker R, Ralph PJ, Seymour
2 JR (2015) Chemotaxis by natural populations of coral reef bacteria. *ISME J* 9:1764–1777
- 3 Tremblay P, Grover R, Maguer JF, Hoogenboom M, Ferrier-Pagès C (2014) Carbon
4 translocation from symbiont to host depends on irradiance and food availability in the
5 tropical coral *Stylophora pistillata*. *Coral Reefs* 33:1–13
- 6 Vallina SM, Simó R (2007) Strong relationship between DMS and the solar radiation dose over
7 the global surface ocean. *Science* 315:506–508
- 8 Van Alstyne KL, Dominique III VJ and Muller-Parker G (2009). Is dimethylsulfoniopropionate
9 (DMSP) produced by the symbionts or the host in an anemone–zooxanthella
10 symbiosis? *Coral Reefs* 28: 167-176
- 11 Warner ME, Berry-Lowe S (2006) Differential xanthophyll cycling and photochemical activity
12 in symbiotic dinoflagellates in multiple locations of three species of Caribbean coral. *J Exp*
13 *Mar Bio Ecol* 339:86–95
- 14 Weis VM (2008) Cellular mechanisms of cnidarian bleaching: stress causes the collapse of
15 symbiosis. *J Exp Biol* 211:3059–3066
- 16 Winters G, Beer S, Zvi BB, Brickner I, Loya Y (2009) Spatial and temporal photoacclimation of
17 *Stylophora pistillata*: zooxanthella size, pigmentation, location and clade. *Mar Ecol Prog*
18 *Ser* 384:107–119
- 19 Wooldridge SA (2008) Water quality and coral bleaching thresholds: formalising the linkage for
20 the inshore reefs of the Great Barrier Reef, Australia. *Mar Pollut Bull* 58:745–751
- 21 Yancey PH, Heppenstall M, Ly S, Andrell RM, Gates RD, Carter VL, Hagedorn M (2010)
22 Betaines and dimethylsulfoniopropionate as major osmolytes in cnidaria with
23 endosymbiotic dinoflagellates. *Physiol Biochem Zool* 83:167–173
- 24 Yost DM, Jones RJ, Mitchelmore CL (2010) Alterations in dimethylsulfoniopropionate (DMSP)
25 levels in the coral *Montastrea franksi* in response to copper exposure. *Aquat Toxicol*
26 98:367–373
- 27 Yost DM, Jones R, Rowe CL, Mitchelmore CL (2012) Quantification of total and particulate
28 dimethylsulfoniopropionate (DMSP) in five Bermudian coral species across a depth
29 gradient. *Coral Reefs* 31:561–570
- 30

31 **Figure legends**

32 **Fig. 1** Mean ($n = 8, \pm$ SE) (a) symbiont densities and (b) chlorophyll a concentrations in control
33 corals (*grey*), corals transplanted from 3 to 20 and 50 m (T3, *striped*) and from 50 to 20 and 3 m
34 (T50, *white*) in the Gulf of Aqaba. Letters above error bars indicate significant differences
35 between groups (SNK tests, $p < 0.05$)

36

37 **Fig. 2** Mean ($n = 8, \pm$ SE) diameter of symbionts in control corals (*grey*), corals transplanted
38 from 3 to 20 and 50 m (T3, *striped*) and from 50 to 20 and 3 m (T50, *white*) in the Gulf of

1 Aqaba. *Letters above error bars* indicate significant differences between groups (SNK tests, $p <$
2 0.05)

3


4 **Fig. 3** Mean concentrations ($n = 8, \pm \text{SE}$) of **(a)** DMSPt expressed as fmol cell^{-1} and of **(b)**
5 DMSPt expressed as nmol cm^{-2} in control corals (*grey*), corals transplanted from 3 to 20 and 50
6 m (T3, *striped*) and from 50 to 20 and 3 m (T50, *white*) in the Gulf of Aqaba. *Letters above error*
7 *bars* indicate significant differences between groups (SNK tests, $p < 0.05$)

8


9 **Fig. 4** Mean concentrations ($n = 8, \pm \text{SE}$) of **(a)** DMSPp expressed as fmol cell^{-1} and **(b)** DMSPp
10 expressed as fmol cm^{-2} in control corals (*grey*), corals transplanted from 3 to 20 and 50 m
11 (*striped*) and from 50 to 20 and 3 m (*white*) in the Gulf of Aqaba. *Letters above error bars*
12 indicate significant differences between groups (SNK tests, $p < 0.05$)

13


14 **Fig. 5** Regression analyses of the relationship between symbiont cell densities and **(a)** DMSPt
15 and **(b)** DMSPp ($n = 56$)




Borell et al. 2016 - Figure 1




Borell et al. 2016 - Figure 2



Borell et al. 2016 - Figure 3



Borell et al. 2016 - Figure 4



Borell et al. 2016 - Figure 5