

Free backbone carbonyls function as linchpins in the activation of rhodopsin

Naoki Kimata^{1,3}, Andreyah Pope^{1,3}, Omar B. Sanchez-Reyes¹, Markus Eilers¹, Chikwado A. Opefi², Martine Ziliox¹, Philip J. Reeves² and Steven O. Smith¹

¹Department of Biochemistry and Cell Biology, 11794-5215, USA, ²School of Biological Sciences, University of Essex, Wivenhoe Park, Essex, United Kingdom CO4 3SQ.

³These authors contributed equally to this work.

Correspondence should be addressed to S.O.S. (email: steven.o.smith@stonybrook.edu)

Abstract

Conserved prolines in the transmembrane helices of G protein-coupled receptors (GPCRs) are often considered to function as hinges that divide the helix into two segments capable of independent motion. Depending on their potential to hydrogen bond, the free C=O groups associated with these prolines can facilitate conformational flexibility, conformational switching or stabilize receptor structure. Solid-state NMR spectroscopy of rhodopsin, a GPCR in the visual receptor subfamily, reveals that the free conserved backbone C=O groups on helices H5 and H7 stabilize the inactive rhodopsin structure and suggest that they serve as molecular linchpins. In response to light-induced isomerization of the retinal chromophore, hydrogen-bonding interactions involving these C=O groups are released allowing H5 and H7 to repack onto the transmembrane core of the receptor. These results provide insights into the multiple structural and functional roles prolines play in membrane proteins.

Prolines within the transmembrane (TM) helices of membrane proteins can function in several ways. First, they may facilitate protein dynamics as flexible hinges and play a functional role in guiding large scale conformational changes ¹. Second, prolines may be key elements in stabilizing protein structure. Proline-induced kinks can facilitate tighter packing of membrane proteins by allowing the helices to adopt optimal side chain interactions ^{2,3} and the backbone carbonyls at the *i-4* position relative to TM prolines are free to form strong stabilizing interhelical hydrogen bonds. Third, prolines and the associated free C=O groups can facilitate (reversible) switching between distinct protein conformations ⁴. A key question in G protein-coupled receptors (GPCRs) is whether conserved prolines play functional roles as hinges that allow helix motion, and/or structural roles in stabilizing helix-helix interactions in this important class of membrane receptors ^{5,6}.

To address the role of prolines in the structure and function of GPCRs, we focus on the visual receptor rhodopsin. Rhodopsin serves as an on-off switch for light detection in the vertebrate retina ⁷. Light energy absorbed by the retinal chromophore in rhodopsin drives the receptor from an inactive to an active conformation. The hallmark of the active state of rhodopsin (Metarhodopsin II or Meta II) is the outward rotation of the intracellular end of TM helix H6. Coupled with H6 motion are changes in the orientations of the adjacent helices H5 and H7. Helices H5, H6, and H7 each contains a proline residue in the middle of the TM sequence: Pro215^{5.50}, Pro267^{6.50} and Pro303^{7.50}, respectively (**Fig. 1**). These residues are the most conserved in each of these helices in the family A GPCRs and according to the Ballesteros-Weinstein universal numbering system ⁸ are designated 5.50, 6.50 and 7.50, respectively. An additional proline (Pro291^{7.38}) with high sequence conservation within the visual receptor subfamily occurs at the extracellular end of H7.

The defining feature of a proline in a TM helix is that it is unable to form a backbone hydrogen bond to the carbonyl group one helical turn away. In the case of Pro215^{5.50} and

Pro303^{7.50} on H5 and H7, respectively, the free *i*-4 backbone carbonyls form hydrogen-bonds with strongly polar residues on adjacent helices. However, the free backbone carbonyls associated with Pro267^{6.50} and Pro291^{7.38} are oriented toward the membrane lipids and do not hydrogen bond in the crystal structures of inactive rhodopsin^{9,10}, active opsin¹¹ or Meta II^{12,13}, suggesting instead that they allow the helical segments to easily swivel. Coordinated motion of the extracellular ends of H6 and H7 has been proposed as part of a general mechanism for GPCR activation^{14,15} raising the possibility that the sequence stretching from Pro267^{6.50} and Pro291^{7.38}, which includes extracellular loop (EL3), pivots upon activation.

Early studies on rhodopsin using FTIR spectroscopy indicated that conformational changes at one or more prolines occurred upon activation¹⁶. FTIR difference spectra revealed a large shift of the amide I vibration associated with a peptide bond adjacent to the amino-terminal side of a proline. The authors left open the possibility that the observed shift was due to *cis-trans* proline isomerization. More recent studies¹⁷ using non-native amino acid substitutions at conserved prolines in the D2 dopamine receptor ruled out the idea that *cis-trans* proline isomerization occurs, at least in this specific GPCR. They found that the main function of proline was to introduce a break in the helix by removing a backbone NH. Introducing either cyclic, R-hydroxy, or N-methyl residues unable to form NH hydrogen bonds at the position of proline resulted in receptors with wild-type function. Consistent with this role, in several GPCRs mutation of proline to residues able to form backbone hydrogen bonds disrupts expression¹⁸ and/or function^{18,19}.

Solid-state NMR spectroscopy in combination with isotope labeling of rhodopsin can be used to clarify the role of specific prolines by targeting the carbonyl groups that are four amino acids upstream of conserved prolines. These C=O groups are free to form alternative hydrogen bonds due to the lack of an NH hydrogen bond partner along their own helix backbone. Our focus is on the highly conserved *i*-4 free C=O groups on helices H5, H6 and H7 whose ¹³C chemical

shifts are sensitive to both secondary structure and hydrogen bonding. Each of these free carbonyls is in a functionally important region of the protein (**Supplementary Fig. 1**). The His211^{5.46} carbonyl associated with conserved Pro215^{5.50} on H5 is located within the retinal-binding site, and is a key determinant of the high sensitivity of the dim light photoreceptors in rod cells, as compared to the color photoreceptors in cone cells²⁰. The Ile263^{6.46} carbonyl associated with conserved Pro267^{6.50} is bracketed by conserved residues (Phe261^{6.44} and Trp265^{6.48}) on TM helix H6 that are part of a transmission switch that couples retinal isomerization to conformational changes on the intracellular side of the receptor^{21,22}. Finally, the Ala299^{7.46} carbonyl associated with conserved Pro303^{7.50} serves to orient the side chain of Asn55^{1.50}, the most conserved residue within the family A GPCRs⁷. The ¹³C chemical shift changes of these free carbonyls upon receptor activation reveal that they play both structural roles in guiding how the TM helices pack in the inactive receptor, as well as functional roles in allowing the helices to reassemble in response to light-induced isomerization of the retinal chromophore. These results highlight the importance of hydrogen-bonding interactions involving the free C=O groups associated with TM prolines in membrane proteins.

RESULTS

The TM core of rhodopsin is highly conserved and serves to couple structural changes on the extracellular side of the receptor containing the retinal chromophore to the intracellular G protein binding region (**Fig. 1**). The crystal structures of the inactive^{9,10} and active¹¹⁻¹³ states of rhodopsin reveal the position of conserved amino acids and provide the framework for understanding how changes in hydrogen bonding guide the rearrangement of TM helices H5-H7. NMR spectroscopy is used here to target the free carbonyl groups within the TM core associated with conserved proline residues. These studies rely extensively on rhodopsin crystal structures that show the positions of the prolines and the free C=O groups relative to surrounding amino acids and water, which can participate in stabilizing hydrogen bonds. NMR measurements of specific ¹³C=O chemical shifts report on the strength of the hydrogen bonding interactions and how they change upon activation. Measurements are made on rhodopsin and the Meta II intermediate with and without the G α peptide to address flexibility at the sites of the free carbonyls.

Pro215^{5,50}

The conserved proline on helix H5 at position 215 in rhodopsin frees the carbonyl of His211^{5,46}. There are six histidine residues in rhodopsin that must be considered. The magic angle spinning (MAS) NMR difference spectrum between the inactive (dark) state of rhodopsin and the active Meta II intermediate shows at least one His ¹³C=O resonance shifts to lower frequency (**Fig. 2a**). The ¹³C=O resonance that changes chemical shift can be assigned to His211^{5,46} by using rotational echo double resonance (REDOR)²³ NMR filtering of rhodopsin labeled with 1-¹³C histidine and ¹⁵N-phenylalanine. There is only a single His-Phe pair in the rhodopsin sequence, namely His211^{5,46}-Phe212^{5,47}. The labeling strategy generates a single ¹³C-¹⁵N labeled peptide bond. The REDOR experiment allows one to measure ¹³C...¹⁵N dipolar couplings in the solid-state NMR experiments, and the strong dipolar coupling resulting from the directly bonded ¹³C-¹⁵N pair allows one to selectively observe only the single ¹³C=O resonance

in the REDOR filtered spectrum²³. The ¹³C=O frequency is observed at 172.5 ppm in rhodopsin and shifts to 170.2 ppm upon conversion to Meta II.

The decrease in the His211^{5,46} ¹³C=O chemical shift in the transition to Meta II can result from changes in hydrogen bonding, backbone torsion angles or both. Table 1 lists the ¹³C chemical shifts of the free C=O groups on H5, H6 and H7 along with the backbone torsion angles associated with these residues in inactive and active structures. The chemical shift changes associated with changes in backbone conformation versus hydrogen bonding can be comparable in magnitude (4-5 ppm)²⁴⁻²⁶. The X-ray crystal structure of rhodopsin shows that the C=O of His211^{5,46} is directly hydrogen-bonded to the Glu122^{3,37} COOH side chain^{9,10}. Nevertheless, the observed chemical shift of 172.5 ppm is lower than that normally observed for hydrogen-bonded C=O groups in α -helices suggesting that the lower chemical shift is caused by non-helical backbone torsion angles (bold, italics in Table 1). Indeed in the crystal structure of rhodopsin, the ϕ and ψ torsion angles show a strong distortion compared to standard α -helices ($\phi = -57^\circ$ and $\psi = -47^\circ$). Upon activation, the His211^{5,46} C=O hydrogen bond with Glu122^{3,37} is broken and a new hydrogen bond is formed between Glu122^{3,37} and the His211^{5,46} imidazole nitrogen^{11,27} (see **Fig. 2e**). The direct His211^{5,46}-Glu122^{3,37} interaction can already be sensed using FTIR spectroscopy in the Meta I intermediate²⁸. In the Meta II crystal structure^{12,13}, the helix is less distorted than in rhodopsin (the backbone torsion angles of His211^{5,46} are closer to standard α -helix values), which would favor a downfield shift in the His211^{5,46} ¹³C=O resonance. The observed upfield chemical shift of the His211^{5,46} C=O in Meta II is in the opposite direction and consequently is attributed to the loss of Glu122^{3,37} hydrogen bonding. This change suggests that the His211^{5,46} C=O functions as a hydrogen-bonding switch in receptor activation.

Pro267^{6,50}

Pro267^{6,50}, the conserved proline in TM helix H6, is associated with the free carbonyl of Ile263^{6,46}. There are 22 isoleucine residues in rhodopsin. In contrast to the histidine C=O

difference spectrum (**Fig. 2a**), the MAS NMR difference between rhodopsin and Meta II containing 1-¹³C isoleucine shows that several isoleucines change chemical shift upon activation (**Fig. 2b**). We take advantage of the unique Ile263^{6.46}-Cys264 pair to identify the Ile263^{6.46} resonance in this spectrum. The REDOR NMR filtered spectrum of rhodopsin (middle panel) labeled with 1-¹³C-isoleucine and ¹⁵N-cysteine exhibits a single resonance at 171.9 ppm assigned to Ile263^{6.46}. The Ile263^{6.46} C=O is oriented away from the helical bundle, toward the lipids, and the 171.9 ppm chemical shift likely reflects the absence of a hydrogen-bonding partner since the backbone ϕ and ψ torsion angles of Ile263^{6.46} are close to values observed in standard α -helices.

Upon activation, the Ile263^{6.46} resonance splits into two components at 170.5 and 175.6 ppm. In the Meta II crystal structure ^{12,13}, the backbone ϕ and ψ torsion angles are still close to values in standard α -helices (Table 1). The upfield (170.5 ppm) component of the Ile263^{6.46} ¹³C=O resonance is consistent with a non-hydrogen bonded C=O and the small (1.4 ppm) upfield chemical shift would be consistent with a small distortion of the backbone from standard ϕ and ψ torsion angles. In contrast, the most likely explanation for the resonance with the large downfield chemical shift is that the Ile263^{6.46} C=O establishes a strong hydrogen bonding interaction upon activation. The crystal structures of Meta II ^{12,13} reveal that the only side chain near the Ile263^{6.46} C=O is the aromatic ring of Phe294^{7.41}.

Pro303^{7.50}

Conserved Pro303^{7.50} on helix H7 frees the carbonyl of Ala299^{7.46} in rhodopsin. With the HEK293S expression system used for ¹³C-labeling rhodopsin, alanine is scrambled and cannot be specifically labeled. To target this carbonyl, we have mutated Ala299^{7.46} to serine. The A299S rhodopsin mutant exhibits a 500 nm absorption band, and the photobleaching behavior and stability of Meta II are similar to the wild-type pigment (unpublished results). The new Ser299^{7.46}-Val300^{7.47} dipeptide sequence is unique. The 1D ¹³C difference spectrum of 1-¹³C

Ser, ^{15}N -Val-labeled rhodopsin along with the REDOR filtered spectra of rhodopsin and Meta II reveals the position of the Ser299 $^{13}\text{C}=\text{O}$ resonance (**Fig. 2c**).

The Ser299^{7.46} $^{13}\text{C}=\text{O}$ chemical shifts from 173.1 ppm to 168.9 ppm upon activation. This group is hydrogen bonded to the side chain NH_2 of Asn55^{1.50}. Like His211^{5.46}, the backbone ϕ and ψ torsion angles of Ala299^{7.46} in rhodopsin are different from those in standard α -helices (Table 1), but in Meta II the values are consistent with α -helix. The large decrease in chemical shift upon activation is therefore attributed to a loss of hydrogen bonding to Asn55^{1.50} (indicated by bold highlight in Table 1). Given the high conservation of both Asn55^{1.50} and the free $\text{C}=\text{O}$ group at position 299 across the family A GPCRs, this structural change upon activation is likely related to receptor function.

Pro291^{7.38}

Pro291^{7.38} at the extracellular end of H7 is highly conserved (82%) in the visual receptor subfamily. The Phe287^{7.34} $\text{C}=\text{O}$ is oriented toward the surrounding lipid and does not form inter-residue hydrogen bonds in the rhodopsin crystal structure^{9,10}. There are two Phe-Met pairs, and correspondingly two peaks in the REDOR spectrum of rhodopsin at 174.1 ppm and 169.3 ppm (**Fig. 2d**). We assign the 174.1 ppm to Phe85 in helix H2. This chemical shift is consistent with the helical secondary structure and backbone hydrogen-bonding of Phe85 in the rhodopsin crystal structure. The 169.3 ppm resonance is assigned to the Phe287^{7.34} $\text{C}=\text{O}$ on the basis of the ϕ and ψ torsion angles and the lack of a clear hydrogen bonding interaction. Upon activation, the 174.1 ppm resonance splits into two components at 174 and 175.3 ppm and the resonance at 169 ppm broadens. We attribute the broadening (and lower intensity) to increased disorder in the backbone structure at Phe287^{7.34}, and unlike the other conserved prolines discussed above, this result is consistent with Pro291^{7.38} functioning as a flexible hinge. The splitting of the 174.1 ppm resonance is tentatively assigned to conformational changes in the

region of Gly89^{2.56}-Gly90^{2.57}, one helical turn from Phe85^{2.52} in the rhodopsin sequence (see Discussion).

Induced fit of the C-terminal G α peptide influences H6

The change in the ¹³C=O resonance of Ile263^{6.46} upon light activation is the most unusual of the three sites associated with the conserved prolines described above. This ¹³C=O resonance shifts downfield in frequency, but is not associated with a clear hydrogen-bonding partner in the crystal structures of active opsin¹¹ or Meta II^{12,13}. Comparison of the crystal structures of rhodopsin and Meta II shows that the backbone torsion angles of Ile263^{6.46} do not change appreciably upon activation. To better understand the structural changes occurring in this region of H6, we have obtained REDOR NMR spectra of the free carbonyls at His211^{5.46}, Ile263^{6.46} and Ser299^{7.46} in the presence of the transducin G α peptide.

It is known that binding of the transducin G α peptide to Meta II stabilizes the active structure²⁹. The C=O chemical shifts of His211^{5.46}, Ile263^{6.46} and Ser299^{7.46} do not move appreciably upon G α peptide binding (**Fig. 3a-c**). However, there is a marked increase in the intensity of the Ile263^{6.46} C=O with the unusual downfield chemical shift. The change in populations suggests there is an induced fit of the G α peptide into its intracellular binding site, which is allosterically coupled to the region containing the Ile263^{6.46} C=O.

DISCUSSION

Prolines are unique in lacking a backbone NH, which effectively eliminates the ability of the carbonyl group at the *i-4* position to form an α -helical hydrogen bond. The three prolines in rhodopsin on helices H5, H6 and H7 are among the most conserved residues in the family A GPCRs, and consequently the free C=O groups at the *i-4* positions associated with these prolines are highly conserved as well. Since breaking a backbone hydrogen bond in a hydrophobic environment is estimated to cost ~4-5 kcal/mol of energy³², the free *i-4* carbonyl represents an energetically favorable site for hydrogen bond formation. Depending on their ability to hydrogen bond, the free C=O groups can stabilize protein structure (strong hydrogen bonds) or facilitate conformational dynamics (weak or no hydrogen bonds). Conformational switching of hydrogen-bonding interactions may significantly lower the energetic barrier for breaking interhelical contacts and as such facilitates the conformational changes that underlie protein function⁴. Our studies focus on the hydrogen bonding interactions of the conserved free C=O groups on H5, H6 and H7 in rhodopsin to address their role in receptor structure and function. The prolines studied cover many of the different types of interactions possible and consequently provide insights into possible functions of prolines in membrane proteins.

The structure of rhodopsin reflects the evolutionary requirements of night vision: a photoreceptor finely tuned to balance low dark noise (via a stable structure) and high sensitivity through rapid receptor dynamics upon single photon absorption. The first four TM helices provide the stable framework onto which helices H5, H6 and H7 pack in the dark-state structure⁷. Helices H5, H6, and H7 each contain highly conserved prolines (Pro215^{5,50}, Pro267^{6,50} and Pro303^{7,50}) and undergo the largest structural changes upon activation (**Fig. 4**). The free C=O groups associated with these prolines are all roughly in the same plane within or near the receptor's TM core (**Fig. 1**). The TM core is highly conserved and involved in both receptor structure and function⁷.

Structural roles likely exist for all three highly conserved prolines. As the receptor folds during protein synthesis, the free carbonyls at His211^{5.46} and Ala299^{7.46} likely latch onto the strongly polar Glu122^{3.37} and Asn55^{1.50} side chains on helices H3 and H1, respectively. Computational studies on receptor stability and unfolding highlight His211^{5.46} and Ala299^{7.46} in the TM region of rhodopsin as the major TM residues that contribute to the overall structural integrity³³. These studies, which assessed receptor stability after removing *single* TM hydrogen bonds, provide strong support for the role of the free C=O groups on H5 and H7 in stabilizing receptor structure. Although His211^{5.46} and Ala299^{7.46} were identified using two slightly different computational approaches, both results are consistent with His211^{5.46} and Ala299^{7.46} being sites of autosomal dominant retinitis pigmentosa (ADRP) mutations, which are typically associated with protein misfolding³⁴. Mutation of Pro267^{6.50} also leads to protein misfolding and ADRP. Since the free C=O associated with Pro267^{6.50} does not hydrogen bond in inactive rhodopsin, its structural role may be to allow a large pivoting motion to optimize contacts of other residues and possibly prevent non-native (mis-folded) interactions³⁵.

Functional roles of the three highly conserved prolines likely involve conformational switching rather than conformational dynamics. Upon activation, both the His211^{5.46} and Ser299^{7.46} C=O groups lose their stabilizing hydrogen bonding interactions, which appear to act as molecular linchpins in rhodopsin. That is, they are stabilizing when in place, but result in a rearrangement of interhelical interactions and helix orientations when broken. The balance between the active and inactive conformations is dependent on whether the retinal binding site is occupied by the 11-*cis* retinal PSB (inverse agonist) or all-*trans* retinal SB (agonist) chromophores. For H5, the new helix orientation is stabilized not by an alternative hydrogen bonding partner for the His211^{5.46} C=O, but instead by the formation of two new interhelical hydrogen bonds, one involving the His211^{5.46} *side chain* with Glu122^{3.37}, and the other involving the interaction of Tyr223^{5.58} with Arg135^{3.50}. Conservative mutations of either Glu122^{3.37} or Tyr223^{5.58} decrease the stability of Meta II^{20,36}, i.e. the active H5 conformation reverts back to an inactive

conformation more readily. For H7, the loss of the hydrogen bonding interaction with Asn55^{1.50} allows H7 to become more helical. Mutations that favor a helical (non-distorted) conformation, as in the P303A mutant, lead to hyperactivity of the receptor³⁷. The presence of molecular switches involving free C=O groups may also exist at other positions that have high subfamily conservation in GPCRs. For example, a recent NMR study suggested that the free C=O associated with conserved Pro4.60 in the β 1-adrenergic receptor is part of a complex hydrogen bonding network linking H4 and H5 that is modulated by ligand binding³⁸.

The observation of two distinct chemical shifts for Ile263^{6.46} C=O suggests that Pro267^{6.50} provides a hinge in H6, but that it is not conformationally dynamic. The C=O resonances are sharp, rather than broadened as observed for Phe287^{7.34}. Consistent with this idea, position 6.46 is 80% conserved as Ile, Val or Leu and only 2% conserved as glycine, which is thought to facilitate dynamics⁶. The mixture of chemical shift changes of the Ile263^{6.46} C=O indicate that H6 does not toggle to a fully active conformation until binding of the G α peptide. In contrast to the C=O groups on H5 and H7, the Ile263^{6.46} C=O shifts downfield consistent with an increase in hydrogen bonding. Furthermore, the chemical shifts or intensities of the free C=O groups associated with H5 and H7 do not markedly change with the addition of the G α peptide, suggesting that H5 and H7 have adopted their fully active conformations prior to G protein binding.

Of the four prolines studied, only Pro291^{7.38} appears to be conformationally dynamic in the active state. The broadened NMR resonances indicate that there is not a distinct conformation in this region of the active receptor. Conformational flexibility may be aided by adjacent glycines or prolines. For example, the conserved PxP motif at the intracellular end of the S6 helix is thought to influence voltage gating in several voltage-gated ion channels via conformational dynamics³⁹. In this case, conformational flexibility may be a consequence of water solvation of the exposed C=O groups within the ion channel. In rhodopsin, a conserved proline upstream of

Pro291^{7.38} at the i-6 position may contribute to the overall flexibility of the extracellular loop 3 connecting H6 and H7. For Pro215^{5.50} and Pro303^{6.50} on H5 and H7, the conservation of the side chain at the i-4 may also contribute to the hinging motion at these positions. For position 5.46, the highest conservation is for glycine (17%) in the family A GPCRs if the olfactory receptors are excluded. Position 5.50 is only moderately conserved (38%) as proline in the olfactory receptor subfamily. Serine has the next highest conservation at 11%. For position 7.46, the highest conservation in the family A GPCRs excluding the olfactory receptors is for serine (56%). Interestingly, in the olfactory receptors position 7.46 is 94% conserved as a proline. The presence of glycine at position and proline at 7.46 may allow increased flexibility where these residues are conserved.

In a survey of high-resolution membrane protein structures from the Protein Data Bank, we have found that ~75% of the free carbonyls have a polar residue nearby in a position to directly hydrogen bond or hydrogen bond via a water molecule (**Supplementary Fig. 2**). Many of these prolines occur in regions that are important in protein function, and they appear in membrane proteins as diverse as transporters, enzymes, and ion channels (**Supplementary Fig. 2**).

Together, these NMR data provide deeper insight into the diverse roles that proline can perform in rhodopsin. Depending on the need, they can provide kinks, conformational switches and hinges in transmembrane helices. As such, these findings are of broad significance because the strategic positioning of prolines and the utilization of their unique properties is important in the structure and function of membrane proteins in general.

Materials and Methods

Expression and purification of ^{13}C labeled rhodopsin

Isotope enriched rhodopsin was extracted from HEK293S membranes using 1% [w/v] n- β -D dodecyl maltopyranoside (DDM) in PBS pH 7.4 and immunoaffinity-purified using Rho1D4-Sepharose. Resin-bound rhodopsin was washed with 50 column volumes of 0.02% DDM in PBS pH 7.2, equilibrated with 0.02% DDM, 2 mM sodium phosphate pH 6.0. Rhodopsin was eluted in 0.02% DDM, sodium phosphate pH 6.0, 100 μM 9-mer elution peptide (TETSQVAPA) and concentrated to a volume of 1 ml using centricon (Millipore) centrifugation devices with a molecular weight cutoff of 30 kDa. The volume was further reduced to < 60 μl under a gentle stream of argon gas and packed into a 4 mm MAS rotor.

Solid-state NMR spectroscopy

Solid-state NMR experiments were conducted at static field strengths of either 500 or 600 MHz using a three-channel 4 mm MAS probe with a spinning rate of 10 kHz. Spectra were collected using a 2 ms contact pulse during cross polarization. SPINAL64 decoupling was used during acquisition with a ^1H RF field strength between 70-90 kHz. For REDOR spectra, the experiment was conducted as previously described⁴⁰, with a dephasing period of 20 rotor cycles at 10 KHz MAS rate (2 ms). The REDOR filtered spectra (ΔS) were obtained by subtracting spectra with (S) and without (S_0) rotor-synchronized ^{15}N π pulses (10-11 μs). To reduce artifacts, S and S_0 spectral acquisition was interleaved and difference spectra were acquired scan-by-scan. ΔS spectra were summed over 60-100K scans using ~5-6 mgs of rhodopsin in a typical sample.

All spectra were acquired at 190 K. For rhodopsin and Meta II spectra, states were cryo-trapped as described previously⁴¹. The ^{13}C high-resolution NMR and solid-state MAS NMR spectra were externally referenced to the ^{13}C resonance of neat TMS at 0 ppm at room temperature. Using TMS as the external reference, we calibrated the carbonyl resonance of

solid glycine at 176.46 ppm. The chemical shift difference between ^{13}C of DSS in D_2O relative to neat TMS is 2.01 ppm.

G α peptide synthesis and reconstitution with rhodopsin

The 15-mer C-terminal peptide of the G α subunit of transducin (TDIIKENLKDCGLF) was synthesized using solid-phase methods (Keck Small Scale Peptide Synthesis Facility, Yale University) and purified by reverse phase HPLC. The experiments on rhodopsin with added G α peptide were carried in mixed micelles using DDM and dioleoylphosphoserine (DOPS). Solubilization of rhodopsin in DDM/DOPS mixed micelles has been shown to facilitate the interaction between rhodopsin and the full heterotrimeric form of transducin⁴². DOPS was added to rhodopsin in DDM micelles in a 1:100 rhodopsin to lipid ratio, and the G α peptide to rhodopsin ratio was 8:1. The stability of the Meta II state was monitored by fluorescence quenching interaction between the indole of Trp265 and the β -ionone ring of the retinal chromophore⁴³.

Acknowledgements

This work was supported by the National Institutes of Health (NIH) grant GM41412 (to S.O.S.) We thank Hiroe Sasaki and Dr. Xin Zhou (Institute of Protein Research, Osaka University) for expression and purification of several of the ^{15}N , ^{13}C labeled rhodopsin samples, and Dr. Joseph Goncalves for preliminary experiments with the $\text{G}\alpha$ peptide.

Author Contributions

M.E., P.J.R and S.O.S conceived the study; N.K, A.P, and M.E. made samples; M.E. and M.Z. collected and analyzed NMR data; A.P. and O.B.S.R. analyzed the protein database for proline interactions; C.A.O constructed rhodopsin mutants; N.K., A.P., P.J.R. and S.O.S wrote the manuscript.

Competing Financial Interests

The authors declare no competing financial interests.

Table 1. NMR $^{13}\text{C}=\text{O}$ chemical shifts and crystal structure Φ , Ψ torsion angles in rhodopsin and Meta II

Residue	State	Chemical shift (ppm) ¹	Φ ²	Ψ
His211	Rho	172.5	<i>-115/-106</i> ⁴	<i>53/22</i> ⁴
	Meta II	170.2 ³	<i>-107</i> ⁴	<i>-3</i> ⁴
Ile263	Rho	171.9 ³	-55/-66	-44/-41
	Meta II	170.5/175.6	-48	-41
Ser299	Rho	173.1	<i>-86/-82</i> ⁴	<i>0/-8</i> ⁴
	Meta II	168.9 ³	-65	-20
Phe287	Rho	169.3 ³	-35/-59	-60/-41
	Meta II	169.3 ³	-56	-34
Standard α -helix		175-177	-57	-47

¹ ^{13}C Chemical shifts (ppm) for rhodopsin and Meta II are taken from REDOR filtering experiments.

² Torsion angles (degrees) are taken from the rhodopsin (PDB ID 1U19/1GZM) and Meta II (PDB ID 3PQR) crystal structures.

³ Low chemical shift values in bold are attributed to loss of or weaker C=O hydrogen bonding since the torsion angles are close to standard helices. For His211, the torsion angles are less distorted from helical values in Meta II than rhodopsin and the lower chemical shift in Meta II is attributed to loss of C=O hydrogen bonding.

⁴ Torsion angles in bold, italics are outside of the range of standard α -helices.

REFERENCES

1. Williams, K.A. & Deber, C.M. Proline residues in transmembrane helices: structural or dynamic role? *Biochemistry* **30**, 8919-8923 (1991).
2. von Heijne, G. Proline kinks in transmembrane α -helices. *J. Mol. Biol.* **218**, 499-503 (1991).
3. Fu, Q. et al. Structural Basis and Functional Role of Intramembrane Trimerization of the Fas/CD95 Death Receptor. *Mol. Cell* **61**, 602-613 (2016).
4. Cao, Z. & Bowie, J.U. Shifting hydrogen bonds may produce flexible transmembrane helices. *Proc. Natl. Acad. Sci. USA* **109**, 8121-8126 (2012).
5. Sansom, M.S.P. & Weinstein, H. Hinges, swivels and switches: The role of prolines in signalling via transmembrane α -helices. *Trends Pharmacol. Sci.* **21**, 445-451 (2000).
6. Cordes, F.S., Bright, J.N. & Sansom, M.S.P. Proline-induced distortions of transmembrane helices. *J. Mol. Biol.* **323**, 951-960 (2002).
7. Smith, S.O. Structure and activation of the visual pigment rhodopsin. *Ann. Rev. Biophys.* **39**, 309-328 (2010).
8. Ballesteros, J.A. & Weinstein, H. Integrated methods for the construction of three dimensional models and computational probing of structure-function relations in G-protein coupled receptors. *Meth. Neurosci.* **25**, 366-428 (1995).
9. Palczewski, K. et al. Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* **289**, 739-745 (2000).
10. Li, J., Edwards, P.C., Burghammer, M., Villa, C. & Schertler, G.F.X. Structure of bovine rhodopsin in a trigonal crystal form. *J. Mol. Biol.* **343**, 1409-38 (2004).
11. Park, J.H., Scheerer, P., Hofmann, K.P., Choe, H.W. & Ernst, O.P. Crystal structure of the ligand-free G-protein-coupled receptor opsin. *Nature* **454**, 183-187 (2008).
12. Deupi, X. et al. Stabilized G protein binding site in the structure of constitutively active metarhodopsin-II. *Proc. Natl. Acad. Sci. USA* **109**, 119-124 (2012).
13. Choe, H.W. et al. Crystal structure of metarhodopsin II. *Nature* **471**, 651-655 (2011).

14. Elling, C.E. et al. Metal ion site engineering indicates a global toggle switch model for seven-transmembrane receptor activation. *J. Biol. Chem.* **281**, 17337-17346 (2006).
15. Schwartz, T.W., Frimurer, T.M., Holst, B., Rosenkilde, M.M. & Elling, C.E. Molecular mechanism of 7TM receptor activation - A global toggle switch model. *Annu. Rev. Pharmacol. Toxicol.* **46**, 481-519 (2006).
16. Ganter, U.M., Gärtner, W. & Siebert, F. Rhodopsin-lumirhodopsin phototransition of bovine rhodopsin investigated by Fourier transform infrared difference spectroscopy. *Biochemistry* **27**, 7480-7488 (1988).
17. Van Arnam, E.B., Lester, H.A. & Dougherty, D.A. Dissecting the functions of conserved prolines within transmembrane helices of the D2 dopamine receptor. *ACS Chem. Biol.* **6**, 1063-1068 (2011).
18. Wess, J., Nanavati, S., Vogel, Z. & Maggio, R. Functional role of proline and tryptophan residues highly conserved among G protein-coupled receptors studied by mutational analysis of the m3 muscarinic receptor. *EMBO J.* **12**, 331-338 (1993).
19. Stitham, J., Martin, K.A. & Hwa, J. The critical role of transmembrane prolines in human prostacyclin receptor activation. *Mol. Pharmacol.* **61**, 1202-1210 (2002).
20. Imai, H. et al. Single amino acid residue as a functional determinant of rod and cone visual pigments. *Proc. Natl. Acad. Sci. USA* **94**, 2322-2326 (1997).
21. Deupi, X. & Standfuss, J. Structural insights into agonist-induced activation of G-protein-coupled receptors. *Curr. Opin. Struct. Biol.* **21**, 541-551 (2011).
22. Deupi, X. Relevance of rhodopsin studies for GPCR activation. *Biochimica et biophysica acta* **1837**, 674-82 (2014).
23. Gullion, T. & Schaefer, J. Rotational-echo double-resonance NMR. *J. Magn. Reson.* **81**, 196-200 (1989).
24. Saito, H. Conformation-dependent C-13 chemical-shifts - a new means of conformational characterization as obtained by high-resolution solid-state C-13 NMR. *Magnetic Resonance in Chemistry* **24**, 835-852 (1986).

25. Gu, Z.T., Zambrano, R. & McDermott, A. Hydrogen-bonding of carboxyl groups in solid state amino acids and peptides: comparison of carbon chemical shielding, infrared frequencies, and structures. *J. Am. Chem. Soc.* **116**, 6368-6372 (1994).
26. Szilagyi, L. Chemical shifts in proteins come of age. *Progress in NMR Spect.* **27**, 325-443 (1995).
27. Scheerer, P. et al. Crystal structure of opsin in its G-protein-interacting conformation. *Nature* **455**, 497-502 (2008).
28. Beck, M., Sakmar, T.P. & Siebert, F. Spectroscopic evidence for interaction between transmembrane helices 3 and 5 in rhodopsin. *Biochemistry* **37**, 7630-7639 (1998).
29. Hamm, H.E. et al. Site of G protein binding to rhodopsin mapped with synthetic peptides from the alpha subunit. *Science* **241**, 832-835 (1988).
30. Vishnivetskiy, S.A. et al. How does arrestin respond to the phosphorylated state of rhodopsin? *J. Biol. Chem.* **274**, 11451-11454 (1999).
31. Krupnick, J.G. & Benovic, J.L. The role of receptor kinases and arrestins in G protein-coupled receptor regulation. *Annu. Rev. Pharmacol. Toxicol.* **38**, 289-319 (1998).
32. White, S.H. & Wimley, W.C. Membrane protein folding and stability: Physical principles. *Annu. Rev. Biophys. Biomol. Struct.* **28**, 319-365 (1999).
33. Rader, A.J. et al. Identification of core amino acids stabilizing rhodopsin. *Proc. Natl. Acad. Sci. USA* **101**, 7246-7251 (2004).
34. Sung, C.H., Davenport, C.M. & Nathans, J. Rhodopsin mutations responsible for autosomal-dominant retinitis-pigmentosa - clustering of functional classes along the polypeptide-chain. *J. Biol. Chem.* **268**, 26645-26649 (1993).
35. Wigley, W.C. et al. A protein sequence that can encode native structure by disfavoring alternate conformations. *Nat. Struct. Biol.* **9**, 381-388 (2002).
36. Goncalves, J.A. et al. Highly conserved tyrosine stabilizes the active state of rhodopsin. *Proc. Natl. Acad. Sci. USA* **107**, 19861-19866 (2010).

37. Fritze, O. et al. Role of the conserved NPxxY(x)(5,6)F motif in the rhodopsin ground state and during activation. *Proc. Natl. Acad. Sci. USA* **100**, 2290-2295 (2003).
38. Isogai, S. et al. Backbone NMR reveals allosteric signal transduction networks in the beta(1)-adrenergic receptor. *Nature* **530**, 237-+ (2016).
39. Labro, A.J., Raes, A.L., Bellens, I., Ottschytsch, N. & Snyders, D.J. Gating of Shaker-type channels requires the flexibility of S6 caused by prolines. *J. Biol. Chem.* **278**, 50724-50731 (2003).
40. Gullion, T. & Schaefer, J. Detection of weak heteronuclear dipolar coupling by rotational-echo double-resonance NMR. in *Advances in Magnetic Resonance, Vol. 13; Conference on "High Resolution NMR in Solids", January 19-21, 1989* (ed. Warren, W.S.) 57-84 (Academic Press, Inc., San Diego, California, USA; London, England, UK, 1989).
41. Eilers, M., Ying, W.W., Reeves, P.J., Khorana, H.G. & Smith, S.O. Magic angle spinning nuclear magnetic resonance of isotopically labeled rhodopsin. *Methods Enzymol.* **343**, 212-222 (2002).
42. Jastrzebska, B., Goc, A., Golczak, M. & Palczewski, K. Phospholipids are needed for the proper formation, stability, and function of the photoactivated rhodopsin-transducin complex. *Biochemistry* **48**, 5159-5170 (2009).
43. Farrens, D.L. & Khorana, H.G. Structure and function in rhodopsin. Measurement of the rate of metarhodopsin II decay by fluorescence spectroscopy. *J. Biol. Chem.* **270**, 5073-5076 (1995).

Figure Legends

Figure 1. Crystal structure of the visual receptor rhodopsin (PDB ID 1U19⁹) showing the positions of Pro215^{5.50}, Pro267^{6.50}, Pro291^{7.38} and Pro303^{7.50}. These prolines are located on helices H5, H6 and H7. The lack of an NH group results in a carbonyl group at the i-4 position from these prolines that is free to form interhelical hydrogen-bonds. The residues (His211^{5.46}, Ile263^{6.46}, Phe287^{7.34}, and Ala299^{7.46}) with the free carbonyl are shown in grey. The three free C=O groups that are conserved across the family A GPCRs (His211^{5.46}, Ile263^{6.46}, and Ala299^{7.46}) lie within or near the TM core of the receptor.

Figure 2. REDOR NMR as a probe of hydrogen bonding changes of carbonyl residues at the i-4 positions of Pro215^{5.50}, Pro267^{6.50}, Pro291^{7.38} and Pro303^{7.50}. (a) One-dimensional MAS NMR difference spectrum of the histidine ¹³C=O resonances between rhodopsin and Meta II (top panel) and REDOR filtered spectra of rhodopsin (middle panel) and Meta II (lower panel) labeled with 1-¹³C His, ¹⁵N Phe. (b) One-dimensional MAS difference spectrum of the isoleucine ¹³C=O resonances between rhodopsin and Meta II (top panel) and REDOR filtered spectra of rhodopsin (middle panel) and Meta II (lower panel) labeled with 1-¹³C Ile, ¹⁵N Cys. (c) One-dimensional MAS difference spectrum of the serine ¹³C=O resonances between rhodopsin and Meta II (top panel) and REDOR filtered spectra of rhodopsin (middle panel) and Meta II (lower panel) labeled with 1-¹³C Ser, ¹⁵N Val. (d). One-dimensional MAS difference spectrum of the phenylalanine ¹³C=O resonances between rhodopsin and Meta II (top panel) and REDOR filtered spectra of rhodopsin (middle panel) and Meta II (lower panel) labeled with 1-¹³C Phe, ¹⁵N Met. (e-h) Below each set of experimental data are the crystal structures of rhodopsin (gray) and Meta II (cyan) in the region of interest.

Figure 3. ¹³C...¹⁵N REDOR NMR experiments of Meta II with (blue) and without (red) the G α peptide of transducin bound. REDOR filtered spectra are shown for His211^{5.46} (a), Ile263^{6.46} (b) and Ser299^{7.46} (c). The addition of G α peptide shifts the equilibrium to a predominantly a single

conformation (blue) in the case of Ile263^{6.46}. The high frequency of the Ile263^{6.46} ¹³C=O chemical shift (175.6 ppm) is characteristic of strong hydrogen-bonding suggesting stabilization of the receptor. The experiments on rhodopsin without and with added G α peptide were carried out in DDM micelles and mixed DDM/DOPS micelles, respectively.

Figure 4. Receptor activation leads to repacking of helices H5-H7 on the TM core of rhodopsin.

(a) Inactive structure of rhodopsin. Helices H1-H4 form a scaffold onto which helices H5-H7 pack. The key packing contacts on H5 and H7 are associated with Pro215^{5.50} and Pro303^{7.50} and their corresponding *i-4* carbonyls. Retinal isomerization disrupts both interactions. For H5, the β -ionone ring has a steric clash at the position of the His211^{5.46}-Glu122^{3.37} hydrogen bond upon conversion to the all-*trans* configuration. For H7, the retinal is covalently attached to Lys296^{7.43}. Upon isomerization, Trp265^{6.48} on H6 rotates away from H7 and disrupts a water mediated hydrogen bond with Asn302^{7.49}, which is part of a hydrogen bonding network stretching from Asn55^{1.50} and Asp83^{2.50} to the Ala299^{7.46} C=O. (b) Overlap of the crystal structures of rhodopsin (1GZM¹⁰, purple) and Meta II (3PQR¹³, light purple) showing the positions of the TM helices. The disruption of the interactions of the His211^{5.46} C=O and Ala299^{7.46} C=O with the H1-H4 scaffold allows helices H5-H7 to reorient. (c,d) Schematic of the hydrogen bonding changes occurring between inactive rhodopsin and active Meta II rhodopsin.

Figure 1




Figure 2




Figure 3




Figure 4

