

1 **Management of recent onset tendon-related pain in a primary contact setting:**
2 **A survey of practice**

3

4 **Kieran Mitham**

5 Dynamic Health, Physiotherapy Department, Cambridgeshire Community Services NHS Trust,
6 Huntingdon, Cambridgeshire, UK

7 **Adrian Mallows**

8 School of Sport, Rehabilitation & Exercise Sciences, University of Essex, Colchester, UK

9 **Gillian Yeowell**

10 Department of Health Professions, Faculty of Health, Psychology & Social Care, Manchester
11 Metropolitan University, Manchester, UK

12 **Chris Littlewood**

13 Department of Health Professions, Faculty of Health, Psychology & Social Care, Manchester
14 Metropolitan University, Manchester, UK

15

INTRODUCTION

16 Tendon-related pain is a common musculoskeletal condition characterised by pain during an
17 activity which applies load to a tendon (Rio et al., 2014). For example, pain related to Achilles
18 tendon-related pain can be commonly experienced with heel raises and pain related to the
19 rotator cuff can be commonly experienced with lifting the arm away from the body. Tendon-
20 related pain affects a wide range of people and affects both function and quality of life (Grimaldi
21 et al., 2015; Malliaras et al., 2015). A wide range of treatments are used to treat people with
22 tendon-related pain, including exercise therapy, injection therapy, shockwave therapy and
23 acupuncture (Mitham et al., 2020; Van Der Vlist et al., 2020).

24 Why some people with recent onset tendon-related pain recover and others go on to
25 experience persistent pain and disability is unclear. One unknown factor is whether
26 management decisions during early consultations influence prognosis. Passive treatments,
27 such as complete rest, have a high variability in tendon load and may lead to persistent pain
28 and disability in runners with Achilles tendon-related pain (Cardoso et al., 2019; Lagas et al.,
29 2020), suggesting a possible link between treatment and persisting symptoms, and the need
30 to consider this further.

31 To allow early access to expert musculoskeletal treatment, a recent change has been the
32 evolution of the role of First Contact Practitioners (FCP within the NHS (Addley et al., 2010;
33 NHS England, 2019). FCPs are physiotherapists working directly in GP practices and treating
34 people with musculoskeletal problems.

35 The practice of this wide range of healthcare professionals within a primary contact setting for
36 people with recent onset tendon-related pain has not been investigated. Given the potential
37 importance of initial management strategies, it is important to understand healthcare
38 professional's current practice and any variability between FCPs and other clinicians (OCs).
39 This survey aimed to report the practice used by different healthcare professionals for two

40 recent onset tendon-related pain scenarios in a primary contact setting. A secondary aim of
41 the survey was to understand if practice differed between locations of pain.

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

MATERIALS AND METHODS

61

Design

62

63 We undertook a cross-sectional online survey hosted by Qualtrics ([X](#)).

Survey Development

64

65 The survey was designed by the study team with reference to two scenarios that reflect recent
66 onset tendon-related scenarios in the shoulder and Achilles (*Table 1*). Recent onset was
67 defined as less than three month duration. The shoulder and Achilles were chosen as locations
68 due to the high prevalence in upper limb and lower limb tendon pain (Littlewood et al., 2013;
69 Riel et al., 2019).

70 The survey was piloted by three members of the study team and two clinical physiotherapists.

71 The survey was then modified accordingly, including adding 'not applicable' responses where
72 indicated and amalgamating questions and changing wording to improve flow and
73 understanding. Based on the findings from the pilot testing, the survey took less than 10
74 minutes to complete. The survey was open for one month to 17 December 2020.

Participants

75

76 We recruited a convenience sample of healthcare professionals, of any professional
77 background, involved in the management of recent onset musculoskeletal conditions in a
78 primary contact setting, for example, GPs, physiotherapists including FCPs, and chiropractors.
79 Potential respondents were invited to participate via personal email, professional networks,
80 and Twitter.

Data Analysis

81

82 Ethical approval to conduct this research was granted by the University Ethics Science and
83 Health Faculty, University of X (ETH2021-0325). Data were exported from Qualtrics to IBM
84 SPSS Statistics, version 25, and Microsoft Excel. The difference in proportion of responses
85 between the shoulder and Achilles scenarios, and between FCPs and OCs were analysed

86 using the Chi-Square test. Statistical significance was set at $p \leq 0.05$. For questions which
87 allowed multiple answers, analysis between FCPS and OCs, descriptive statistics were
88 presented.

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

RESULTS

109 In total, 118 surveys were completed; 103 (87%) respondents completed both scenarios and
110 118 (100%) respondents completed just the first. The response by professional background
111 is shown in *Table 2*. The descriptive data from the survey is displayed in *Appendix 1*.

Diagnosis

113 *Rotator Cuff Related Shoulder Pain (RCRSP)* was the preferred diagnostic term by 64/118
114 (54.2%) for the shoulder scenario. *Achilles tendinopathy* was the preferred term by 86/103
115 (83.5%) for the Achilles scenario.

Management

117 Rest was not advised for the shoulder or Achilles scenario by 96/118 (81.4%) and 73/103
118 (70.9%) respondents, respectively. The difference between scenarios for rest was not
119 statistically significant ($p=0.07$). Rest was not advised by 18/23 (78.3%) FCPs, and 77/95
120 (81.1%) OCs for the shoulder scenario. Rest was not advised by 13/23 (65.0%) FCPs and
121 60/83 (72.3%) OCs for the Achilles scenario. The difference between FCPs and OCs advising
122 rest for the shoulder and Achilles was not statistically significant ($p=.0.44$; $p=0.54$)

123 Medication was not recommended by 53/118 (44.9%) for the shoulder scenario and 66/103
124 (64.1%) for the Achilles. Non-opioid medication was recommended by 53/118 (44.9%) for the
125 shoulder scenario and 28/103 (27.2%) for the Achilles. The difference between scenarios
126 regarding medication was not statistically significant ($p=0.1$). NSAIDs were recommended by
127 3/23 (13.0%) FCPs compared to 5/95 (5.3%) OCs for the shoulder scenario ($p=0.006$).
128 NSAIDs were recommended by 4/20 (20.0%) FCPs compared to 4/83 (4.8%) OCs for the
129 Achilles scenario ($p=0.046$) (*Table 3*).

130

131 Injections were not recommended by 113/118 (95.8%) for the shoulder scenario and 103/103
132 (100%) for the Achilles.

133 Amended duties were advised for manual workers by 37/118 (31.4%) in the shoulder scenario
134 and 14/103 (13.6%) for the Achilles. The difference between scenarios regarding return to
135 work was statistically significant ($p<0.0001$). This difference was not observed for office
136 workers ($p=0.49$). There was no statically significant difference in the management of return
137 to work for manual workers between FCPs and OCs for the shoulder scenario ($p=0.65$) or the
138 Achilles ($p=0.97$).

139 Treatment Modalities

140 Adjuncts to exercise were not recommended by 44/118 (37.3%) for the shoulder scenario and
141 39/103 (37.9%) for the Achilles, respectively. Nine different adjuncts were recommended at
142 least once. “*Ice and/or heat*” was the most popular modality advised, 52/118 (44.1%) for the
143 shoulder scenario and 52/103 (50.5%) for the Achilles. “*Massage*” was selected by 4/118
144 (3.4%) for the shoulder scenario and 14/103 (13.6%) respondents for the Achilles. There was
145 no statistically significant difference between the two scenarios regarding the use of adjuncts
146 to exercise ($p=0.782$).

147 Exercise was recommended by 116/118 (98.3%) for the shoulder scenario and 102/103 (99%)
148 for the Achilles. Isometric and isotonic exercises were the most popular treatment advised for
149 both scenarios (*Fig. 2*). Eccentric exercises were recommended by 22/118 (18.6%) of the
150 respondents for the shoulder scenario compared to 35/103 (34.0%) for the Achilles. Heavy
151 slow resistance exercises were recommended by 17/118 (14.4%) of the respondents for the
152 shoulder scenario compared to 29/103 (28.2%) respondents for the Achilles. There was no
153 statistically significant difference between the two scenarios ($p=0.086$).

154 Lifestyle factors

155 The role of obesity was discussed by 82/118 (69.5%) for the shoulder scenario compared to
156 95/103 (94.1%) for the Achilles (*Fig. 3*). The role of alcohol was discussed by 60/118 (51.7%)
157 for the shoulder scenario and 55/103 (54.5%) for the Achilles.

158 Psychosocial factors

159 Occupational factors were discussed by 111/118 (94.1%) for the shoulder scenario compared
160 to 87/118 (84.5%) for the Achilles. Support of friends and family was discussed by 70/118
161 (59.3%) for the shoulder scenario and 59/103 (57.3%) for the Achilles.

162 Prognosis

163 Resolution of symptoms within one month was expected by 18/118 (15.3%) for the shoulder
164 scenario compared to 8/103 (7.8%) for the Achilles. Resolution of symptoms within one to two
165 months was expected by 51/118 (43.2%) for the shoulder scenario compared to 29/103
166 (28.2%) for the Achilles (*Fig. 4*). Resolution of symptoms within one month was expected for
167 the shoulder scenario by 1/23 (4.3%) FCP compared to 17/95 (17.9%) OCs (*Fig. 5*).

168 A follow up visit with a healthcare professional was advised recommended by 102/118 (86.4%)
169 for the shoulder scenario and 91/103 (88.3%) for the Achilles. A follow-up appointment was
170 advised by 84/95 (88.4%) and 75/83 (90.4%) OCs for the shoulder and Achilles scenarios,
171 respectively. A follow-up appointment was advised by 18/23 (78.3%) and 16/20 (80.0%) FCPs
172 for the shoulder and Achilles scenarios, respectively.

173

174

175

176

177

178

179

180

181

182

DISCUSSION

183 We conducted a survey to investigate current practice in relation to recent onset tendon-
184 related pain in a primary contact setting. To the authors' knowledge, this is the first survey to
185 investigate this and compare both FCPs and OCs, and different locations of pain for
186 management of recent onset tendon-related pain.

187 Despite both scenarios being identical except the location in pain, there were significant
188 differences in the approach to the management, prognosis, and recommendations for
189 modifications in workload. Reasons for the difference of management are uncertain, but it is
190 suggested that a similar approach between healthcare professionals to recent-onset
191 atraumatic musculoskeletal conditions should be adopted (Caneiro et al., 2020).

192 Diagnostic labels may influence patients understanding, expectations and beliefs regarding
193 musculoskeletal conditions (Carroll et al., 2016). This survey highlighted some uncertainty
194 regarding diagnostic label due to a wider range of diagnostic labels suggested for the shoulder
195 scenario compared to the Achilles. This is unsurprising; recent expert consensus was gained
196 for the diagnostic label for the Achilles tendon-related pain, but consensus could not be
197 reached for tendon-related pain at the shoulder (Scott et al., 2020). Nevertheless, clinicians
198 should be aware the impact of language and diagnostic labels they use with patients and how
199 it may influence understanding, behaviour and therefore outcomes from treatment (Cuff &
200 Littlewood, 2018).

201 NSAIDs are frequently prescribed drugs to ease pain and reduce inflammation for recent-
202 onset musculoskeletal conditions (NHS, 2019). Paracetamol and/or NSAIDs are
203 recommended in the early stage of both shoulder tendon-related pain and Achilles tendon-
204 related pain, but are not recommended in the longer term (NICE, 2017, 2020). Maquirriain &
205 Kokalj (2014) found significant improvement in pain intensity for recent-onset Achilles tendon-
206 related pain when treated with NSAIDs. NSAIDs have been shown to be superior to placebo
207 but there is no evidence about how they compare to other treatments such as exercise for

208 shoulder tendon-related pain (Steuri et al., 2017). In this survey, FCPs were more likely to
209 recommend NSAIDs for both scenarios compared to OCs.

210 Imaging is not considered necessary for a clinical diagnosis of tendon-related pain, and should
211 only be used for differential diagnosis (Scott et al., 2020). The majority of respondents did not
212 recommend further investigations or steroid injections for the either scenario. This compares
213 favourably to the management of shoulder tendon-related pain by GPs with research reporting
214 up to 53% of new shoulder tendon-related pain presentations being referred by GPs for
215 imaging (Naunton et al., 2020).

216 Healthcare professionals have an important role in the return-to-work process for people
217 complaining of shoulder and Achilles tendon-related pain and early intervention has been
218 suggested a key recommendation (Doiron-Cadrin et al., 2020). Amended duties were more
219 likely to be recommended for the shoulder scenario compared to the Achilles for manual
220 workers ($p < 0.0001$). Given our understanding of the negative impact of not being at work for
221 other musculoskeletal conditions (Foster et al., 2018), exploring reasons for this advice for
222 would be beneficial.

223 Isometric exercises were the most popular for both scenarios, despite the limited evidence of
224 effectiveness (Mitham et al., 2020; van der Vlist et al., 2020). Both eccentric and heavy slow
225 resisted exercises were preferred for Achilles scenario, despite current research suggesting
226 no superiority between exercises (Head et al., 2019; Van Der Vlist et al., 2020). This is
227 particularly interesting given the majority of respondents recommended a follow up visit with
228 a healthcare professional for both scenarios. The wide variety of exercise modalities chosen
229 by the participants, suggests further research is needed to improve our understanding around
230 treatment effectiveness for recent onset tendon-related pain, including a wait and see
231 approach.

232 Moderate alcohol consumption has been suggested as a potential risk factor for Achilles
233 tendon-related pain (Van Der Vlist et al., 2019), as well as metabolic factors for both shoulder

234 and Achilles tendon-related pain (Burne et al., 2019; O'Neill et al., 2016). Respondents in this
235 survey were more likely to discuss obesity, smoking and physical activity for the Achilles
236 scenario compared to the shoulder, despite the association between these factors and
237 shoulder pain (Özkuk & Ateş, 2020; Rechart et al., 2010). In non-athletic individuals,
238 metabolic factors have been suggested to be more prevalent and can negatively influence the
239 recovery of tendon-related pain with exercise treatment (Millar et al., 2021).

240 In regards to resolution of symptoms, the difference between FCPs and OCs was not
241 statistically significant, but we observed a trend towards OCs recommending a more
242 favourable recovery. Improvement in Achilles tendon-related pain can be observed after 12
243 weeks of a loading exercise program, but some people may still have ongoing pain and
244 reduced function (Murphy et al., 2018). Discussions between clinicians and patients are
245 important in healthcare. The respondents in this survey may have been too optimistic for an
246 earlier recovery as 43.2% of respondents expected resolution of symptoms within one to two
247 months for the shoulder scenario. Patients want clear and consistent information regarding
248 prognosis for musculoskeletal pain (Lim et al., 2019). It is important clinicians provide clear
249 and consistent information regarding prognosis to facilitate realistic expectations for patients
250 (Lim et al., 2019).

251 Study limitations

252 The limited number of self-selected responders means the results might not be generalisable
253 to the wider population of clinicians involved in the management of recent onset tendon-related
254 pain in a primary care setting. Recruitment for this survey was conducted, in part, via Twitter
255 which may have excluded healthcare professionals who do not use this platform or other
256 platforms instead. Survey studies are also susceptible to response, social and acquiescence
257 bias as individuals with an interest are more likely to respond. However, the use of additional
258 recruitment strategies including email and professional networks may have gone some way to
259 mitigate this.

260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281

CONCLUSION

Data from this survey highlights some consistency between clinicians in their management of recent onset tendon-related pain; the majority of clinicians recommend against further investigations, steroid injections and recommending for exercise as an intervention. Understanding whether these approaches are clinically effective requires further investigation.

- 283 Addley, K., Burke, C., & McQuillan, P. (2010). Impact of a direct access occupational
284 physiotherapy treatment service. *Occupational Medicine*, 60(8), 651–653.
285 <https://doi.org/10.1093/occmed/kqq160>
- 286 Babatunde, O. O., Jordan, J. L., Van Der Windt, D. A., Hill, J. C., Foster, N. E., & Protheroe,
287 J. (2017). Effective treatment options for musculoskeletal pain in primary care: A
288 systematic overview of current evidence. In *PLoS ONE* (Vol. 12, Issue 6).
289 <https://doi.org/10.1371/journal.pone.0178621>
- 290 Burne, G., Mansfield, M., Gaida, J. E., & Lewis, J. S. (2019). Is there an association between
291 metabolic syndrome and rotator cuff-related shoulder pain? A systematic review. *BMJ*
292 *Open Sport and Exercise Medicine*, 5(1). <https://doi.org/10.1136/bmjsem-2019-000544>
- 293 Caneiro, J. P., Alaiti, R. K., Fukusawa, L., Hespanhol, L., Brukner, P., Sullivan, P. P. B. O.,
294 An, L., & Pain, A. S. (2020). There is more to pain than tissue damage : eight principles
295 to guide care of acute non- - traumatic pain in sport. *British Journal of Sports Medicine*,
296 0(0), 1–2. <https://doi.org/10.1136/bjsports-2019-101705>
- 297 Cardoso, T. B., Pizzari, T., Kinsella, R., Hope, D., & Cook, J. L. (2019). Current trends in
298 tendinopathy management. *Best Practice and Research: Clinical Rheumatology*, 33(1),
299 122–140. <https://doi.org/10.1016/j.berh.2019.02.001>
- 300 Carroll, L. J., Lis, A., Weiser, S., & Torti, J. (2016). How Well Do You Expect to Recover, and
301 What Does Recovery Mean, Anyway? Qualitative Study of Expectations After a
302 Musculoskeletal Injury. *Physical Therapy*, 96(6), 797–807.
303 <https://doi.org/10.2522/ptj.20150229>
- 304 Cuff, A., & Littlewood, C. (2018). Subacromial impingement syndrome – What does this
305 mean to and for the patient? A qualitative study. *Musculoskeletal Science and Practice*,
306 33(August 2017), 24–28. <https://doi.org/10.1016/j.msksp.2017.10.008>

307 Doiron-Cadrin, P., Lafrance, S., Saulnier, M., Cournoyer, É., Roy, J. S., Dyer, J. O.,
308 Frémont, P., Dionne, C., MacDermid, J. C., Tousignant, M., Rochette, A., Lowry, V.,
309 Bureau, N. J., Lamontagne, M., Coutu, M. F., Lavigne, P., & Desmeules, F. (2020).
310 Shoulder Rotator Cuff Disorders: A Systematic Review of Clinical Practice Guidelines
311 and Semantic Analyses of Recommendations. *Archives of Physical Medicine and*
312 *Rehabilitation*, 101(7), 1233–1242. <https://doi.org/10.1016/j.apmr.2019.12.017>

313 Foster, N. E., Anema, J. R., Cherkin, D., Chou, R., Cohen, S. P., Gross, D. P., Ferreira, P.
314 H., Fritz, J. M., Koes, B. W., Peul, W., Turner, J. A., Maher, C. G., Buchbinder, R.,
315 Hartvigsen, J., Cherkin, D., Foster, N. E., Maher, C. G., Underwood, M., van Tulder, M.,
316 ... Woolf, A. (2018). Prevention and treatment of low back pain: evidence, challenges,
317 and promising directions. *The Lancet*, 391(10137), 2368–2383.
318 [https://doi.org/10.1016/S0140-6736\(18\)30489-6](https://doi.org/10.1016/S0140-6736(18)30489-6)

319 Grimaldi, A., Mellor, R., Hodges, P., Bennell, K., Wajswelner, H., & Vicenzino, B. (2015).
320 Gluteal Tendinopathy: A Review of Mechanisms, Assessment and Management. *Sports*
321 *Medicine*, 45(8), 1107–1119. <https://doi.org/10.1007/s40279-015-0336-5>

322 Head, J., Mallows, A., Debenham, J., Travers, M. J., & Allen, L. (2019). The efficacy of
323 loading programmes for improving patient-reported outcomes in chronic midportion
324 Achilles tendinopathy: A systematic review. *Musculoskeletal Care*, 17(4), 283–299.
325 <https://doi.org/10.1002/msc.1428>

326 Lagas, I. F., Fokkema, T., Bierma-Zeinstra, S. M. A., Verhaar, J. A. N., van Middelkoop, M.,
327 & de Vos, R. (2020). How many runners with new-onset Achilles tendinopathy develop
328 persisting symptoms? A large prospective cohort study. *Scandinavian Journal of*
329 *Medicine & Science in Sports*, sms.13760. <https://doi.org/10.1111/sms.13760>

330 Lim, Y. Z., Chou, L., Au, R. T., Seneviwickrama, K. M. D., Cicuttini, F. M., Briggs, A. M.,
331 Sullivan, K., Urquhart, D. M., & Wluka, A. E. (2019). People with low back pain want
332 clear, consistent and personalised information on prognosis, treatment options and self-

333 management strategies: a systematic review. *Journal of Physiotherapy*, 65(3), 124–
334 135. <https://doi.org/10.1016/j.jphys.2019.05.010>

335 Littlewood, C., May, S., & Walters, S. (2013). Epidemiology of Rotator Cuff Tendinopathy: A
336 Systematic Review. *Shoulder & Elbow*, 5(4), 256–265.
337 <https://doi.org/10.1111/sae.12028>

338 Malliaras, P., Cook, J., Purdam, C., & Rio, E. (2015). Patellar Tendinopathy: Clinical
339 Diagnosis, Load Management, and Advice for Challenging Case Presentations. *Journal*
340 *of Orthopaedic & Sports Physical Therapy*, 45(11), 887–898.
341 <https://doi.org/10.2519/jospt.2015.5987>

342 Maquirriain, J., & Kokalj, A. (2014). Acute Achilles tendinopathy: Effect of pain control on leg
343 stiffness. *Journal of Musculoskeletal Neuronal Interactions*, 14(1), 131–136.

344 Millar, N. L., Silbernagel, K. G., Thorborg, K., Kirwan, P. D., Galatz, L. M., Abrams, G. D.,
345 Murrell, G. A. C., McInnes, I. B., & Rodeo, S. A. (2021). Tendinopathy. *Nature Reviews*
346 *Disease Primers*, 7(1), 1. <https://doi.org/10.1038/s41572-020-00234-1>

347 Mitham, K., Mallows, A., Seneviratne, G., Debenham, J., & Malliaras, P. (2020).
348 Conservative Management of Acute Lower Limb Tendinopathies: A Systematic Review.
349 *Musculoskeletal Care*.

350 Murphy, M., Travers, M., Gibson, W., Chivers, P., Debenham, J., Docking, S., & Rio, E.
351 (2018). Rate of Improvement of Pain and Function in Mid-Portion Achilles Tendinopathy
352 with Loading Protocols: A Systematic Review and Longitudinal Meta-Analysis. *Sports*
353 *Medicine*, 48(8), 1875–1891.
354 <http://search.ebscohost.com/login.aspx?direct=true&db=s3h&AN=130551969&site=eho>
355 st-live

356 Naunton, J., Harrison, C., Britt, H., Haines, T., & Malliaras, P. (2020). General practice
357 management of rotator cuff related shoulder pain: A reliance on ultrasound and

358 injection guided care. *PLoS ONE*, 15(1), 1–14.
359 <https://doi.org/10.1371/journal.pone.0227688>

360 NHS. (2019). *NSAIDs*. <https://www.nhs.uk/conditions/nsaids/>

361 NHS England. (2019). *Elective Care High Impact Interventions: First Contact Practitioner for*
362 *MSK Services*.

363 NICE. (n.d.). *Scenario: Rotator cuff disorders | Management | Shoulder pain | CKS | NICE*.
364 2017. Retrieved January 8, 2021, from [https://cks.nice.org.uk/topics/shoulder-](https://cks.nice.org.uk/topics/shoulder-pain/management/rotator-cuff-disorders/)
365 [pain/management/rotator-cuff-disorders/](https://cks.nice.org.uk/topics/shoulder-pain/management/rotator-cuff-disorders/)

366 NICE. (2020). *Scenario: Management | Management | Achilles tendinopathy | CKS | NICE*.
367 <https://cks.nice.org.uk/topics/achilles-tendinopathy/management/management/>

368 O'Neill, S., Watson, P. J., & Barry, S. (2016). a Delphi Study of Risk Factors for Achilles
369 Tendinopathy- Opinions of World Tendon Experts. *International Journal of Sports*
370 *Physical Therapy*, 11(5), 684–697.
371 <http://www.ncbi.nlm.nih.gov/pubmed/27757281>[http://www.pubmedcentral.nih.gov/a](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5046962)
372 [rticlerender.fcgi?artid=PMC5046962](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5046962)

373 Özkuk, K., & Ateş, Z. (2020). The effect of obesity on pain and disability in chronic shoulder
374 pain patients. *Journal of Back and Musculoskeletal Rehabilitation*, 33(1), 73–79.
375 <https://doi.org/10.3233/BMR-181384>

376 Public Health England. (2019). *Musculoskeletal conditions profile: short commentary,*
377 *December 2019 - GOV.UK*.
378 [https://www.gov.uk/government/publications/musculoskeletal-conditions-profile-](https://www.gov.uk/government/publications/musculoskeletal-conditions-profile-december-2019-update/musculoskeletal-conditions-profile-short-commentary-december-2019)
379 [december-2019-update/musculoskeletal-conditions-profile-short-commentary-](https://www.gov.uk/government/publications/musculoskeletal-conditions-profile-short-commentary-december-2019)
380 [december-2019](https://www.gov.uk/government/publications/musculoskeletal-conditions-profile-short-commentary-december-2019)

381 Rechart, M., Shiri, R., Karppinen, J., Jula, A., Heliövaara, M., & Viikari-Juntura, E. (2010).
382 Lifestyle and metabolic factors in relation to shoulder pain and rotator cuff tendinitis: A

383 population-based study. *BMC Musculoskeletal Disorders*, 11.
384 <https://doi.org/10.1186/1471-2474-11-165>

385 Riel, H., Lindstrøm, C. F., Rathleff, M. S., Jensen, M. B., & Olesen, J. L. (2019). Prevalence
386 and incidence rate of lower-extremity tendinopathies in a Danish general practice: A
387 registry-based study. *BMC Musculoskeletal Disorders*, 20(1), 4–9.
388 <https://doi.org/10.1186/s12891-019-2629-6>

389 Rio, E., Moseley, L., Purdam, C., Samiric, T., Kidgell, D., Pearce, A. J., Jaberzadeh, S., &
390 Cook, J. (2014). The pain of tendinopathy: Physiological or pathophysiological? *Sports*
391 *Medicine*, 44(1), 9–23. <https://doi.org/10.1007/s40279-013-0096-z>

392 Scott, A., Squier, K., Alfredson, H., Bahr, R., Cook, J. L., Coombes, B., De Vos, R. J., Fu, S.
393 N., Grimaldi, A., Lewis, J. S., Maffulli, N., Magnusson, S. P., Malliaras, P., Mc Auliffe,
394 S., Oei, E. H. G., Purdam, C. R., Rees, J. D., Rio, E. K., Gravare Silbernagel, K., ...
395 Zwerver, J. (2020). ICON 2019: International Scientific Tendinopathy Symposium
396 Consensus: Clinical Terminology. *British Journal of Sports Medicine*, 54(5), 260–262.
397 <https://doi.org/10.1136/bjsports-2019-100885>

398 Steuri, R., Sattelmayer, M., Elsig, S., Kolly, C., Tal, A., Taeymans, J., & Hilfiker, R. (2017).
399 Effectiveness of conservative interventions including exercise, manual therapy and
400 medical management in adults with shoulder impingement: a systematic review and
401 meta-analysis of RCTs. *British Journal of Sports Medicine*, bjsports-2016-096515.
402 <https://doi.org/10.1136/bjsports-2016-096515>

403 Van Der Vlist, A. C., Breda, S. J., Oei, E. H. G., Verhaar, J. A. N., & De Vos, R. J. (2019).
404 Clinical risk factors for Achilles tendinopathy: A systematic review. *British Journal of*
405 *Sports Medicine*, 53(21), 1352–1361. <https://doi.org/10.1136/bjsports-2018-099991>

406 van der Vlist, A. C., van Veldhoven, P. L. J., van Oosterom, R. F., Verhaar, J. A. N., & de
407 Vos, R. J. (2020). Isometric exercises do not provide immediate pain relief in Achilles
408 tendinopathy: A quasi-randomized clinical trial. *Scandinavian Journal of Medicine and*

409 *Science in Sports*, 30(9), 1712–1721. <https://doi.org/10.1111/sms.13728>

410 Van Der Vlist, A. C., Winters, M., Weir, A., Ardern, C. L., Welton, N. J., Caldwell, D. M.,
411 Verhaar, J. A. N., & De Vos, R.-J. (2020). Which treatment is most effective for patients
412 with Achilles tendinopathy? A living systematic review with network meta-analysis of 29
413 randomised controlled trials. *Br J Sports Med*, 0, 1–8. [https://doi.org/10.1136/bjsports-](https://doi.org/10.1136/bjsports-2019-101872)
414 [2019-101872](https://doi.org/10.1136/bjsports-2019-101872)

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431 **Table 1.** Presenting condition, subjective and objective findings for the shoulder and Achilles scenarios

Patient Characteristics	Referral	Duration	Aggravating Factors	Easing Factors	Physical Examination
50 year old person	Non-traumatic right shoulder pain	3-week history	<ul style="list-style-type: none"> Lifting kettle Taking a coat on and off Moving the arm away from the body 	<ul style="list-style-type: none"> Resting arm by their side Avoiding provocative positions 	<ul style="list-style-type: none"> Resisted shoulder abduction and external rotation are painful. Full passive shoulder external rotation. Examination of neck does not reproduce shoulder pain. No redness, bruising or swelling present.
50 year old person	Non-traumatic right Achilles pain	3-week history	<ul style="list-style-type: none"> At the start of light jogging Stiff with prolonged rest 	<ul style="list-style-type: none"> Eases after a few minutes of jogging. 	<ul style="list-style-type: none"> Pain with palpation of mid-portion Achilles tendon. Pain on double leg calf raise Full passive dorsiflexion and plantarflexion Examination of knee and does not reproduce any heel pain No redness, bruising or swelling present.

432

433 **Table 2.** Respondents by professional background

Scenario	Professional Background (n)	
Shoulder (118)	Physiotherapist (97)	Physiotherapist (69)
		First Contact Practitioner (23)
		Advanced Practice Physiotherapist (5)
	General Practitioner (6)	
	Chiropractor (12)	
	Sports Rehabilitator (1)	
	Orthopaedic Surgeon (1)	
Sport & Exercise Medicine Physician (1)		
Achilles (103)	Physiotherapist (87)	Physiotherapist (62)
		First Contact Practitioner (20)
		Advanced Practice Physiotherapist (5)
	General Practitioner (4)	
	Chiropractor (9)	
	Sports Rehabilitator (1)	
	Orthopaedic Surgeon (1)	
Sport & Exercise Medicine Physician (1)		

434

435


436

437 **Table 3.** Respondent's data for medication for shoulder and Achilles scenario, and FCPs and OCs

Medication	FCP Shoulder	FCP Achilles	OC Shoulder	OC Achilles
None	9/23 (39.1%)	11/20 (55.0%)	44/95 (46.3%)	55/83 (66.3%)
Non-Opioid	10/23 (43.5%)	4/20 (20.0%)	43/95 (45.3%)	24/83 (28.9%)
NSAID	3/23 (13.0%)	4/20 (20.0%)	5/95 (5.3%)	4/83 (4.8%)
Mild-Opioid	0/23 (0.0%)	0/20 (0.0%)	2/95 (2.1%)	0/83 (0.0%)
It Depends	1/23 (4.3%)	1/20 (5.0%)	1/95 (1.1%)	0/83 (0.0%)

438


439



440

Figure 1. Active treatment modalities selected for the shoulder and Achilles scenarios

441




442

443

Figure 2. Lifestyle factors discussed by respondents for the shoulder and Achilles


444



445

446

Figure 3. Prognosis advised by respondents for the scenarios



447

448

Figure 4. Prognosis advised by FCPs and OCs for the shoulder scenario

449

Would you choose to refer for further investigation at this stage?

	<i>n</i> (%)	No	X-Ray	MRI	USS	Bloods
Shoulder						
	All	112 (94.9)	0	1 (0.8)	4 (3.4)	1 (0.8)
	FCP (23)	23 (100)	0 (0)	0 (0)	0 (0)	0 (0)
	Other (95)	89 (93.7)	0 (0)	1 (1.1)	4 (4.2)	1 (1.1)
Achilles						
	All	98 (96.1)	0 (0)	0 (0)	4 (3.9)	1 (1)
	FCP (20)	20 (100)	0 (0)	0 (0)	0 (0)	0 (0)
	Other (83)	78 (94)	0 (0)	0 (0)	4 (4.8)	1 (1.2)

Would you recommend rest for this person, if so, how long?

	No	Relative Rest	<1 week	1-2 weeks	>2-4 weeks	>4-6 weeks	>6 weeks
Shoulder							
	All (118)	96 (81.4)	6 (5.1)	9 (7.6)	5 (4.2)	1 (0.9)	1 (0.9)
	FCP (23)	18 (78.3)	0 (0)	1 (4.4)	1 (4.4)	1 (4.4)	0 (0)
	Other (95)	77 (81.1)	5 (5.3)	8 (8.4)	4 (4.2)	0 (0)	1 (1.1)
Achilles							
	All (103)	73 (70.9)	9 (8.7)	9 (8.7)	5 (4.9)	5 (4.9)	0 (0)
	FCP (20)	13 (65)	4 (20)	1 (5)	0 (0)	2 (10)	0 (0)
	Other (83)	60 (72.3)	7 (8.4)	8 (9.6)	5 (6)	1 (1.2)	0 (0)

Would you recommend any medication for this person?

	No	Non-Opioid	NSAID	Mild Opioid	It Depends
Shoulder					
	All (118)	53 (44.9)	53 (44.9)	8 (6.8)	2 (1.7)
	FCP (23)	9 (39.1)	10 (43.5)	3 (13)	0 (0)
	Other (95)	44 (46.3)	43 (45.3)	5 (5.3)	2 (2.1)
Achilles					
	All (103)	66 (64.1)	28 (27.2)	8 (7.8)	0 (0)
	FCP (20)	11 (55)	4 (20)	4 (20)	0 (0)
	Other (83)	55 (66.3)	24 (28.9)	4 (4.8)	0 (0)

If this person were a manual or office worker, would you advise them to continue to work?

	No (Manual)	No (Office)	Amended Duties (Manual)	Amended Duties (Office)	Full Duties (Manual)	Full Duties (Office)	It Depends (Manual)	It Depends (Office)
Shoulder								
All (118)	2 (1.7)	0 (0)	94 (79.7)	37 (31.4)	12 (10.2)	73 (61.9)	10 (8.5)	8 (6.8)
FCP (23)	0 (0)	0 (0)	20 (87)	3 (13)	1 (4.3)	17 (73.9)	2 (8.7)	3 (13)
Other (95)	2 (2.1)	0 (0)	74 (79.6)	34 (25.8)	9 (9.7)	56 (58.9)	8 (8.6)	5 (5.3)
Achilles								
All (103)	3 (2.9)	1 (1)	54 (52.4)	14 (13.6)	39 (37.9)	83 (80.6)	7 (6.8)	5 (4.9)
FCP (20)	1 (5)	1 (5)	10 (50)	1 (5)	7 (35)	18 (90)	2 (10)	0 (0)
Other (83)	2 (2.4)	0 (0)	44 (53)	13 (15.7)	32 (38.6)	65 (78.3)	5 (6)	5 (6)

Would you discuss any of the following if they were evident? Some respondents checked more than one option

	No	Obesity	Smoking	Physical Activity	Alcohol	Metabolic Disorders
Shoulder						
All (118)	4 (3.5)	82 (70.7)	88 (75.9)	107 (92.2)	60 (51.7)	92 (79.3)
FCP (23)	1 (4.4)	18 (78.3)	17 (73.9)	20 (87)	13 (56.5)	18 (78.3)
Other (95)	3 (3.2)	64 (67.4)	71 (74.7)	87 (91.6)	47 (49.5)	74 (77.9)
Achilles						
All (103)	0 (0)	95 (94.1)	85 (84.2)	99 (98.0)	55 (54.5)	80 (79.2)
FCP (20)	0 (0)	18 (90)	17 (85)	19 (95)	10 (50)	17 (85)
Other (83)	0 (0)	77 (92.8)	68 (81.9)	80 (96.4)	45 (54.2)	63 (75.9)

Would you discuss any of the following if they were evident? Some respondents checked more than one option

	No	Occupation	Friends & Family Support	Social Participation
Shoulder				
All (118)	3 (2.5)	111 (94.1)	70 (59.3)	73 (61.9)
FCP (23)	0 (0)	21 (91.3)	14 (60.9)	14 (60.9)
Other (95)	3 (3.2)	90 (94.7)	56 (58.9)	59 (62.1)
Achilles				
All (103)	9 (8.7)	87 (84.5)	59 (57.3)	75 (72.8)
FCP (20)	1 (5)	16 (80)	10 (50)	15 (75)
Other (83)	8 (9.6)	71 (85.5)	49 (59)	60 (72.3)

Would you discuss any of the following if they were evident? Some respondents checked more than one option

	No	Emotional Factors	Cognitive Factors	Behavioural Factors
Shoulder				
All (118)	2 (1.7)	99 (83.9)	100 (84.7)	103 (87.3)
FCP (23)	1 (4.3)	17 (73.9)	20 (87)	20 (87)
Other (95)	1 (1.1)	82 (86.3)	80 (84.2)	83 (87.4)
Achilles				
All (103)	3 (2.5)	85 (72)	94 (79.7)	89 (75.4)
FCP (20)	1 (5)	14 (70)	17 (85)	17 (85)
Other (83)	2 (2.4)	71 (85.5)	77 (92.8)	72 (86.7)

How long would you expect it to take for this person to experience meaningful recovery from the first time they see you?

	<1 month	1-2 months	>2-4 months	>4-6 months	>6 months	I would not expect meaningful recovery
Shoulder						
All (118)	18 (15.3)	51 (43.2)	40 (33.9)	7 (5.9)	2 (1.7)	0 (0)
FCP (23)	1 (4.3)	9 (39.1)	11 (47.8)	1 (4.3)	1 (4.3)	0 (0)
Other (95)	17 (17.9)	42 (44.2)	29 (30.5)	6 (6.3)	1 (1.1)	0 (0)
Achilles						
All (103)	8 (7.8)	29 (28.2)	50 (48.5)	14 (13.6)	2 (1.9)	0 (0)
FCP (20)	1 (5)	4 (20)	14 (70)	1 (5)	0 (0)	0 (0)
Other (83)	7 (6.8)	25 (24.3)	36 (35)	12 (12.6)	2 (1.9)	0 (0)

Would they benefit from a future appointment with a musculoskeletal health care professional at this stage i.e., physiotherapist, osteopath, chiropractor etc?

	Yes	No
Shoulder		
All (118)	102 (86.4)	16 (13.6)
FCP (23)	18 (78.3)	5 (21.7)
Other (95)	84 (88.4)	11 (11.6)
Achilles		
All (103)	91 (88.3)	12 (11.7)
FCP (20)	16 (80)	4 (20)
Other (83)	75 (90.4)	8 (9.6)

