

Does Model Complexity add Value to Asset Allocation? Evidence from Machine Learning Forecasting Models*

Iason Kynigakis[†] and Ekaterini Panopoulou[‡]

Abstract

This study evaluates the benefits of integrating return forecasts from a variety of machine learning and forecast combination methods into an out-of-sample asset allocation framework. The economic evaluation of the forecasts shows that model complexity translates to improved results in the majority of cases considered, with shrinkage methods and shallow neural networks generating the highest individual performance. Overall, an investor would consistently realize superior out-of-sample gains by incorporating forecast combinations of machine learning models in the portfolio formation process.

Keywords:

Forecast Combination, Machine Learning, Portfolio Optimization, Return Predictability

*We would like to thank three anonymous referees and the Editor, Eric Ghysels, for helpful advice and suggestions. We also thank participants at the 2018 International Conference on Computational and Financial Econometrics, the 2019 International Conference on Applied Theory, Macro and Empirical Finance, the 2019 Annual Conference of the International Association for Applied Econometrics, the 2019 International Accounting and Finance Doctoral Symposium. Iason Kynigakis gratefully acknowledges the support of Science Foundation Ireland under grant number 16/SPP/3347 and 17/SPP/5447.

[†]Smurfit Graduate Business School, University College Dublin, Dublin, Ireland, Email: iason.kynigakis@ucd.ie

[‡]Corresponding author. Essex Business School, University of Essex, Colchester, UK, Email: a.panopoulou@essex.ac.uk, Phone: +44 (0) 1206 872194

1. Introduction

The allocation of wealth among risky assets is one of the most important problems faced by investors. Investors' objectives and constraints along with estimating expected returns plays a crucial role in constructing optimal portfolios. Since forecasting returns is quite challenging, the historical average is often used as an input in portfolio optimization. However, existing literature shows that out-of-sample return predictability adds economic value to asset allocation. This study sets out to examine whether return forecasts generated by linear and nonlinear machine learning methods and their combinations benefit portfolios consisting of stocks, bonds and commodities, when compared to simple forecast combinations, the equal-weighted portfolio or portfolios based on the historical average.

Our study contributes primarily to two strands of literature. First, our study adds to the literature of asset allocation and portfolio formation that exploits the predictability of asset returns. There exists a rich literature in finance, such as DeMiguel, Garlappi and Uppal (2009), Duchin and Levy (2009), Kritzman, Page and Turkington (2010), Kirby and Ostdiek (2012), Bianchi and Guidolin (2014) and Gao and Nardari (2018), who evaluate the out-of-sample performance of asset portfolios relative to simple benchmarks such as the equal-weighted portfolio. Specifically, we relate to the literature that incorporates machine learning in an asset allocation framework (see Callot, Caner, Onder, and Ulasan, 2019; D'Hondt, De Winne, Ghysels and Raymond, 2020). Our contribution to this strand of literature arises from investigating the benefits of integrating return forecasts from machine learning methodologies into an out-of-sample portfolio optimization framework, by comparing the alternative portfolios to the widely used benchmarks of the equal-weighted portfolio and portfolios based on the historical average forecast.

Second, it contributes to the growing literature that uses machine learning methodologies to forecast economic and financial variables. The methodologies we employ have been applied in the context of macroeconomic forecasting using a large number of predictors. Specifically, studies using shrinkage methods to examine the predictability of key macroeconomic indicators include Bai and Ng (2008), De Mol, Giannone and Reichlin (2008) and Stock and Watson (2012), Carrasco and Rossi (2016), Kotchoni, Leroux and Stevanovic (2019) and Babii, Ghysels and Striaukas (2019). The advantages of machine learning in the context of asset pricing and return predictability have been explored among

others by Rapach, Strauss and Zhou (2013), Neely, Rapach, Tu and Zhou (2014), Lima and Meng (2017), Kelly, Pruitt and Su (2019), Rapach, Strauss, Tu and Zhou (2019) and Kozak (2019). Gu, Kelly and Xiu (2020), Bianchi, Büchner and Tamoni (2020) and Kim and Swanson (2014) provide a comprehensive comparison of the predictive accuracy of machine learning methodologies for the equity premium, bond risk premia and key macroeconomic variables, respectively. Our contribution to this literature stems from exploring the economic value of a wide range of machine learning methods when used to model stock, bond and commodity returns used as inputs to asset allocation.

We employ a variety of machine learning methods along with forecast combination schemes to generate the return forecasts for individual stocks, bonds and commodities. Specifically, we consider shrinkage methods with a wide range of convex and non-convex penalties, supervised and unsupervised dimensionality reduction techniques, ensembles of regression trees, support vector machines, artificial neural networks and methods that combine forecasts not only from single predictor models, but also those from the machine learning approaches.

To explore the potential benefits of using the machine learning methods in an asset allocation setting, we construct portfolios based on the return forecasts generated from the multivariate prediction models. We compare the out-of-sample performance of portfolios utilizing machine learning forecasts to that of the equal-weighted portfolio and a mean-variance portfolio based on the historical average forecast. The analysis is conducted for a conservative and an aggressive investor and for different levels of leverage. Additionally, we conduct robustness analysis and investigate how alternative estimates of the covariance matrix affect the performance of the portfolios. We explore the performance of the portfolios in terms of the average characteristics of the weight vectors, variable importance and evaluate the models for the full sample and around NBER-dated recessions and expansions. Finally, we examine the performance of the portfolios incorporating transaction costs based on a penalized mean-variance objective function that dampens the effects of transaction costs to portfolio returns.

Overall, our study shows that using machine learning techniques can be beneficial for the out-of-sample portfolio performance. Our asset allocation results show that the majority of the portfolios outperform the equal-weighted and historical average portfolio benchmarks. When comparing portfolios across different combinations of weight constraints, our findings indicate that allocations that allow leverage

further improve the performance of portfolios based on machine learning methods and that machine learning methods benefit more the portfolios of an aggressive investor. However, we observe that mean-variance portfolios exhibit similar out-of-sample performance across different specifications of the covariance matrix. Furthermore, we find that portfolios based on machine learning consist primarily of stocks and commodities, with bonds being a small part of portfolios belonging to a conservative investor. For the equal-weighted and historical average and most of the simple forecast combination portfolios we find higher Sharpe ratios during expansionary periods. The pattern is similar for most of the dimensionality reduction and non-linear machine learning methods, while for shrinkage methods and machine learning forecast combinations the pattern is reversed. Finally, when introducing transaction costs, the performance of the portfolios deteriorates due to the high degree of turnover, even when using the modified mean-variance objective function. However, portfolios based on forecast combinations of machine learning models tend to yield the best performance.

The remainder of this study is organized as follows. Section 2 describes the methods used to generate the out-of-sample return forecasts used in portfolio optimization. Section 3 provides details on the data, sample splitting and hyperparameter tuning. Section 4 presents the asset allocation framework and examines the economic value of the forecasts, Section 5 investigates further the portfolio performance, while Section 6 concludes.

2. Return Prediction Models

Let r_t denote the return of an asset at time t and $\mathbf{r} = (r_1, r_2, \dots, r_T)$ the $T \times 1$ vector of asset returns.

We express by $x_{i,t-1}$ the i th predictor at time $t - 1$, $\mathbf{x}_{t-1} = (x_{1,t-1}, x_{2,t-1}, \dots, x_{p,t-1})'$ is the $1 \times p$ vector of p candidate predictors and $\mathbf{X} = (\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{T-1})$ the $T \times p$ matrix of predictors. The equation we use to model an asset's return takes the following general form

$$r_t = f(\mathbf{x}_{t-1}) + \varepsilon_t \quad (1)$$

Once the model \hat{f} has been estimated, the expected return of an asset at time $t + 1$ using data available through t is $E_t(r_{t+1}) = \hat{f}(\mathbf{x}_t)$. This prediction can then be used in asset allocation.

A commonly used method is the classic normal linear regression model, estimated by ordinary least squares (OLS). In this case the form to approximate f is the following linear function

$$f_{\boldsymbol{\theta}}(\mathbf{X}) = \alpha + \mathbf{X}\boldsymbol{\beta} \quad (2)$$

where α is the intercept and $\boldsymbol{\beta} = (\beta_1, \beta_2, \dots, \beta_p)$ is the coefficient vector. The estimates of the parameters $\boldsymbol{\theta} = (\alpha, \boldsymbol{\beta})$ are obtained by minimizing the residual sum of squares:

$$\underset{\boldsymbol{\theta}}{\operatorname{argmin}} \mathcal{L}(\boldsymbol{\theta}) = \underset{\boldsymbol{\theta}}{\operatorname{argmin}} \|\mathbf{r} - f_{\boldsymbol{\theta}}(\mathbf{X})\|^2 \quad (3)$$

where $\mathcal{L}(\cdot)$ indicates the least squares loss and $\|\cdot\|$ denotes the l_2 norm. First, we consider the kitchen sink (KS) model, which is a multivariate prediction model utilizing all p predictors. The forecast at $t + 1$ of a regression with p predictors is given by $\hat{r}_{t+1} = \hat{\alpha} + \mathbf{x}_t \hat{\boldsymbol{\beta}}$.

It is well known that this model tends to have poor forecasting performance, as the estimated parameters have low bias but high variance. This problem becomes more acute as the number of predictors increases. Another simple model we consider is the historical average (HA), where the forecast is estimated as $\hat{r}_{t+1} = (1/t) \sum_{s=1}^t r_s$.

In this study we focus on the evaluation of machine learning methods for estimating f in the context of mean-variance portfolio optimization. To this end, we consider alternative models that belong to shrinkage, dimensionality reduction, non-linear machine learning methods and forecast combinations that combine the estimates of simple bivariate prediction models or those of machine learning models.

2.1. Shrinkage Methods

In general, shrinkage methods regularize the coefficient estimates and involve fitting the model in all p predictors. These procedures shrink the coefficients towards zero relative to the OLS estimates and aim at significantly reducing the respective coefficient variances. Shrinkage methods can also perform variable selection depending on the type of regularization. A shrinkage method is similar to the simple linear model, in that it considers only the baseline, untransformed predictors, however, it modifies the least squares problem by adding one additional term in the loss function. In the most general form, a shrinkage method includes a penalty term in the loss function:

$$\operatorname{argmin}_{\boldsymbol{\theta}}[\mathcal{L}(\boldsymbol{\theta}) + \mathcal{P}(\boldsymbol{\beta}; \cdot)], \text{ where } \mathcal{P}(\boldsymbol{\beta}; \cdot) = \sum_{i=1}^p \mathcal{P}(\beta_i; \cdot) \quad (4)$$

There are several choices for the penalty function $\mathcal{P}(\cdot)$.¹ We consider the shrinkage methods with the following penalties: ridge, lasso, elastic net, adaptive lasso, bridge, smoothly clipped absolute deviation, minimax concave penalty and smooth integration of counting and absolute deviation.

The first set of shrinkage methods are all based on convex penalties that can be derived by the following penalty function:

$$\mathcal{P}(\beta_i; \lambda; \gamma; a) = \lambda \hat{w}_i [\gamma |\beta_i| + (1 - \gamma) \beta_i^2], \text{ with } \hat{w}_i = \frac{1}{|\hat{\beta}_i|^a} \quad (5)$$

where $\lambda > 0$ is a tuning parameter, which is determined separately and controls the amount of shrinkage and γ is a hyperparameter that controls the trade-off between l_1 and l_2 regularization. The parameter \hat{w}_i is the weight corresponding to coefficient $|\beta_i|$, $\hat{\beta}_i$ is the OLS estimate and $a \geq 0$ is a hyperparameter that controls the strength of the weight.

When $a = 0$ and $\gamma = 0$ then the above function becomes *ridge regression* (Hoerl and Kennard, 1970) that shrinks the coefficients towards zero and when $a = 0$ and $\gamma = 1$ it yields the *least absolute shrinkage and selection operator* (lasso), introduced by Tibshirani (1996), which allows for both shrinkage and variable selection, by setting some of the coefficients equal to zero. The *elastic net* (EN), proposed by Zou and Hastie (2005), combines both l_1 and l_2 terms in the penalty, thus simultaneously performing continuous shrinkage, automatic variable selection and can also select groups of correlated variables. The penalty produces the EN solution when $a = 0$ and $\gamma \in (0,1)$. The *adaptive lasso* (ALasso), developed by Zou (2006), solves the drawback of the original lasso, which is that it does not necessarily satisfy the oracle properties (Fan and Li, 2001). This is achieved by modifying the lasso to include adaptive weights that penalize individual coefficients less severely. The adaptive lasso solution is derived by setting $a > 0$ and $\gamma = 1$.

¹ Note that the intercept, α , is not included in the penalty term. The penalty is applied to the coefficient vector $\boldsymbol{\beta}$ that measures the association of each predictor with the asset returns and not the intercept, which is a measure of the mean value of the asset returns when, $\mathbf{x}_1 = \mathbf{x}_2 = \dots = \mathbf{x}_p = 0$. Penalization on the intercept is not typically considered, since it would make the optimization procedure dependent on the origin chosen for the asset returns; i.e. adding a constant to each observation of the asset returns, would not simply result in a shift of the predictions by the same amount.

The convex penalties described above are combinations of l_1 and l_2 terms that often select models that are overly dense (Fan and Li, 2001). It is typical in such situations to turn to greedier methods such as those that employ non-convex penalties that include an l_0 term, which is typically associated to best-subsets selection and penalizes the number of non-zero coefficients in the model. These penalties achieve sparser solutions for the same or improved prediction accuracy and enjoy superior variable-selection properties.

Bridge regression developed by Frank and Friedman (1993) and Friedman (2012), has a penalty term based on the l_γ norm and is given by

$$\mathcal{P}(\beta_i; \lambda; \gamma) = \lambda |\beta_i|^\gamma \quad (6)$$

where $\lambda > 0$ and $\gamma \geq 0$ are the two tuning parameters. The bridge penalty term for $0 \leq \gamma \leq 2$ represents all the penalties between ridge regression and best-subsets selection. When using the squared error loss it includes ridge ($\gamma = 2$), the lasso ($\gamma = 1$) and best-subsets ($\gamma = 0$), which produces the sparsest solutions by forcing many coefficients to be equal to zero and applies no shrinkage to the non-zero coefficients. For $\gamma > 1$ all coefficients are strictly non-zero and all penalties in the power family are convex, while for $\gamma < 1$ the penalties are non-convex.

The *smoothly clipped absolute deviation* (SCAD), proposed by Fan and Li (2001), is a non-convex penalty function given by

$$\mathcal{P}(\beta_i; \lambda; \gamma) = \begin{cases} \lambda |\beta_i|, & \text{if } |\beta_i| \leq \lambda \\ \frac{2\gamma\lambda|\beta_i| - |\beta_i|^2 - \lambda^2}{2(\gamma - 1)}, & \text{if } \lambda < |\beta_i| \leq \gamma\lambda \\ \frac{\lambda^2(\gamma + 1)}{2}, & \text{if } |\beta_i| > \gamma\lambda \end{cases} \quad (7)$$

for $\gamma > 2$. SCAD coincides with the lasso until $|\beta_i| = \lambda$, then smoothly transitions to a quadratic function until $|\beta_i| = \gamma\lambda$ and then it remains constant for all $|\beta_i| > \gamma\lambda$. For small coefficients, the SCAD penalty has similar penalization rate as the lasso but leaves large coefficients not excessively penalized.

The *minimax concave penalty* (MCP), developed by Zhang (2010), is defined by

$$\mathcal{P}(\beta_i; \lambda; \gamma) = \begin{cases} \lambda |\beta_i| - \frac{|\beta_i|^2}{2\gamma}, & \text{if } |\beta_i| \leq \lambda\gamma \\ \frac{\gamma\lambda^2}{2}, & \text{if } |\beta_i| > \lambda\gamma \end{cases} \quad (8)$$

for each value of $\lambda > 0$ and $\gamma > 1$, there is a continuum of penalties and threshold operators varying from hard thresholding ($\gamma \rightarrow 1 +$) to soft thresholding ($\gamma \rightarrow \infty$). MCP starts with the same rate of penalization as the lasso but smoothly relaxes the penalization rate to zero as the absolute value of the coefficient increases. Furthermore, MCP relaxes the penalization rate immediately, compared to SCAD, where the rate remains flat for a while before decreasing.

Finally, the *smooth integration of counting and absolute deviation* (SICA) penalty (Lv and Fan, 2009) takes the form

$$\mathcal{P}(\beta_i; \lambda; \gamma) = \lambda \frac{(\gamma + 1)|\beta_i|}{\gamma + |\beta_i|} \quad (9)$$

with $\lambda > 0$ and a small shape parameter $\gamma > 0$. SICA is another non-convex regularization method which is a combination between the l_0 and l_1 penalties and therefore gives sparse solutions. For smaller values of γ , SICA yields results closer to the best-subsets selection, while for larger values of γ it is closer to the lasso.²

2.2. Dimensionality Reduction Methods

The next set of models incorporates the information of a large set of economic variables in a predictive regression framework using latent factors, which are estimated either in a supervised way (using information in both \mathbf{r} and \mathbf{X}) or an unsupervised way (using information only in \mathbf{X}). The function f takes a similar form to equation (2)

$$f(\mathbf{X}) = \alpha + (\mathbf{X}\mathbf{A})\boldsymbol{\beta} = \alpha + \mathbf{Z}\boldsymbol{\beta} \quad (10)$$

where $\mathbf{A} = (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2, \dots, \boldsymbol{\alpha}_K)$ is a $p \times K$ matrix of weights, $\boldsymbol{\beta} = (\beta_1, \beta_2, \dots, \beta_K)$ is the coefficient vector, $\mathbf{Z} = (\mathbf{z}_0, \mathbf{z}_1, \dots, \mathbf{z}_{T-1})$ is a $T \times K$ matrix of latent factors, with $\mathbf{z}_{t-1} = (z_{1,t-1}, z_{2,t-1}, \dots, z_{K,t-1})'$ and $K \ll p$. The latent factor models we use differ based on how the matrix \mathbf{A} is derived.

Partial least squares (PLS), introduced by Wold (1966) and more recently by Kelly and Pruitt (2015), identifies the features in a supervised way. Specifically, PLS decomposes the matrix of predictors \mathbf{X}

² For all penalized methods, we choose $\lambda \in [10^{-3}, 10^3]$. For the elastic net, $\gamma \in [0.1, 0.9]$ so that the solutions differ from those of ridge or the lasso, while for the adaptive lasso $a \in [0.1, 2]$. For the bridge penalty $\gamma \in [0.1, 1.9]$, excluding $\gamma = 1$, so that the ridge and lasso solutions are not included. For SCAD, $\gamma \in [3.7, 10]$, 3.7 being the recommended value in Fan and Li (2001). For MCP, $\gamma \in [1.5, 100]$, while for SICA we used the values, $\gamma = \{10^{-2}, 10^{-1}\}$.

and the zero-mean vector of asset returns \mathbf{r} into the form: $\mathbf{X} = \mathbf{Z}\mathbf{P}' + \mathbf{E}$ and $\mathbf{r} = \mathbf{Z}\mathbf{q}' + \mathbf{e}$, where the matrix \mathbf{P} and the vector \mathbf{q} are the loadings, while \mathbf{E} and \mathbf{e} are the residuals. In order to find the PLS component matrix \mathbf{Z} , the columns of the weight matrix \mathbf{A} need to be obtained through successive optimization problems. The criterion to find the k^{th} estimated weight vector \mathbf{a}_k is

$$\underset{\mathbf{a}}{\operatorname{argmax}}[\boldsymbol{\alpha}'(\mathbf{X}'\mathbf{r}\mathbf{r}'\mathbf{X})\mathbf{a}] \quad \text{s.t.} \quad \boldsymbol{\alpha}'\boldsymbol{\alpha} = 1, \quad \boldsymbol{\alpha}'\boldsymbol{\Sigma}_{\mathbf{X}\mathbf{X}}\boldsymbol{\alpha} = 0 \quad (11)$$

where $\boldsymbol{\Sigma}_{\mathbf{X}\mathbf{X}}$ is the sample covariance of \mathbf{X} . The version of PLS we employ is SIMPLS proposed by de Jong (1993). If $K = p$ then PLS would give a solution equivalent to the OLS estimates.

Sparse partial least squares (SPLS) is an extension of PLS that imposes the l_1 penalty to promote sparsity onto a surrogate weight vector \mathbf{c} instead of the original weight vector \mathbf{a} , while keeping \mathbf{a} and \mathbf{c} close to each other (Chun and Keles, 2010). The first SPLS weight vector solves

$$\underset{\mathbf{a}, \mathbf{c}}{\operatorname{argmin}}[(\mathbf{c} - \mathbf{a})'(\mathbf{X}'\mathbf{r}\mathbf{r}'\mathbf{X})(\mathbf{c} - \mathbf{a}) + \lambda_1 \|\mathbf{c}\|_1 + \lambda_2 \|\mathbf{c}\|^2] \quad \text{s.t.} \quad \boldsymbol{\alpha}'\boldsymbol{\alpha} = 1 \quad (12)$$

where λ_1 and λ_2 are non-negative tuning parameters. To solve SPLS a large λ_2 value is usually required and setting $\lambda_2 = \infty$ yields a solution that has the form of the soft threshold estimator by Zou and Hastie (2005). This reduces the number of tuning parameters to two, the tuning parameter λ_1 and the number of latent factors K .

In the dimensionality reduction methods described above the directions that best represent the predictors \mathbf{X} are derived in a supervised way since the vector of asset returns, \mathbf{r} , is used to determine the component directions. The next set of models derives the latent factors in an unsupervised way.

Principal component analysis (PCA) is the most widely used method to obtain estimates of the latent factors. PCA can be viewed as a regression-type problem (see e.g. Friedman, Hastie and Tibshirani, 2009) where the goal is to find the first K principal component weight vectors by minimizing:

$$\underset{\mathbf{A}}{\operatorname{argmin}}\|\mathbf{X} - \mathbf{X}\mathbf{A}\mathbf{A}'\|^2, \quad \text{s.t.} \quad \mathbf{A}'\mathbf{A} = \mathbf{I}_K \quad (13)$$

The solution to this problem is most often obtained via singular value decomposition: $\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}'$, by setting $\mathbf{A} = \mathbf{V}$. The columns of $\mathbf{V} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_K)$ are the principal components weights. Each \mathbf{v}_k is used to derive the k th principal component, $\mathbf{z}_k = \mathbf{X}\mathbf{v}_k$, thus, $\mathbf{Z}\mathbf{V}$ is the dimension reduced version of the

original predictors. The derived variable \mathbf{z}_1 is the first principal component of \mathbf{X} and has the largest sample variance amongst all normalized linear combinations of the columns of \mathbf{X} .

Sparse principal component analysis (SPCA), developed by Zou, Hastie and Tibshirani (2006), is based on the regression/reconstruction property of PCA and produces modified principal components with sparse weights, such that each principal component is a linear combination of only a few of the original predictors. They show how PCA can be viewed in terms of a ridge regression problem and by adding the l_1 penalty, they convert it to an elastic net regression, which allows for the estimation of sparse principal components. The following regression criterion is proposed to derive the sparse principal component weights:

$$\operatorname{argmin}_{\mathbf{A}, \mathbf{C}} [\|\mathbf{X} - \mathbf{XAC}'\|^2 + \lambda_1 \|\mathbf{c}\|_1 + \lambda_2 \|\mathbf{c}\|^2] \quad \text{s.t.} \quad \mathbf{A}'\mathbf{A} = \mathbf{I}_K \quad (14)$$

where \mathbf{C} is $p \times K$. If $\lambda_1 = \lambda_2 = 0$, $T > p$ and restrict $\mathbf{C} = \mathbf{A}$, then the minimizer of the objective function is exactly the first K weight vectors of ordinary PCA. When $p \gg T$, in order to obtain a unique solution, $\lambda_2 > 0$ is required. The l_1 penalty on \mathbf{c}_k induces sparseness of the weights, with larger values of λ_1 leading to sparser solutions. The algorithm by Zou and Hastie (2005) is used to compute the sparse approximations of each principal component.

Independent component analysis (ICA), developed by Comon (1994), aims at finding a linear representation of non-Gaussian data so that the components are statistically independent. The ICA objective is:

$$\operatorname{argmin}_{\mathbf{A}} \|\mathbf{XA}\|_1, \quad \text{s.t.} \quad \mathbf{A}'\mathbf{A} = \mathbf{I}_K \quad (15)$$

Solving the ICA problem amounts to finding an orthogonal \mathbf{A} such that the components of the vector random variable $\mathbf{Z} = \mathbf{XA}$ are independent and non-Gaussian. More in detail, the independent components are estimated by iterative estimation of the matrix \mathbf{A} , systematically increasing the degree of independence of the components. However, since there is no direct measure of independence, non-Gaussianity is used instead.³

³ Popular approaches for measuring independence or non-Gaussianity in ICA are based on entropy. We use the FastICA algorithm developed by Hyvärinen and Oja (2000), which uses negentropy as a measure of Non-Gaussianity.

Ordinary ICA has two drawbacks; it requires constrained optimization which can become difficult in high dimensional settings and is sensitive to whitening, a pre-processing step that decorrelates the input data, which cannot always be computed exactly when $p \gg T$. Le, Karpenko, Ngiam and Ng (2011), propose *reconstruction independent component analysis* (RICA), which overcomes the drawbacks of ICA, by replacing ICA's orthonormality constraint with a reconstruction penalty. This produces the unconstrained problem:

$$\underset{\mathbf{A}}{\operatorname{argmin}}[\|\mathbf{XA}\|_1 + \lambda\|\mathbf{X} - \mathbf{XAA}'\|^2] \quad (16)$$

where $\lambda > 0$ is a regularization parameter. RICA is equivalent to ICA when $K < p$, data is whitened and λ approaches infinity.⁴

2.3. Nonlinear Machine Learning Methods

So far the models we described assume a linear relationship between return and predictors. In this section we consider three types of nonlinear models from the machine learning literature; ensembles of regression trees, support vector machines and artificial neural networks.

A regression tree is a non-parametric model that is constructed using a recursive binary splitting approach (Breiman, Friedman, Stone and Olshen, 1984). At first, starting from the top of the tree, we divide the predictor space into two distinct and non-overlapping rectangular regions or leaves and then model the asset returns as the simple average of \mathbf{r} within that region. Then one or both of those regions are split into two more regions and this process continues until certain stopping criteria are met. The predictor variable upon which a branch is based, and the value where the branch is split, is chosen to minimize the forecast error. Specifically, the prediction of a tree, \mathcal{T} , with M leaves R_1, R_2, \dots, R_M and maximum depth D , is defined as $\mathcal{T}(\mathbf{X}, c, M, D) = \sum_{m=1}^M c_m \mathbf{I}_{\{\mathbf{X} \in R_m(D)\}}$, where $R_m(D)$ represents one of the partitions of the predictor space. The score c associated with partition m , c_m , is the simple average of returns within region R_m , written as $c_m = 1/T_m \sum_{\mathbf{X} \in R_m} \mathbf{r}$, where T_m denotes the number of observations in region m . At each branch we choose the sorting variable among the set of predictors and the split value that minimize the following loss function:

⁴ For all dimensionality reduction approaches the number of latent factors is $K \in [1,6]$. The parameter that controls for the l_1 penalty in the sparse methods is $\lambda_1 \in [10^{-2}, 10^3]$. Then for reconstruction ICA $\lambda \in [10^{-2}, 10]$.

$$Q_m(c, T_m) = \frac{1}{T_m} \sum_{\mathbf{x} \in R_m} (\mathbf{r} - c_m)^2 \quad (17)$$

Branching halts when the depth, D , of the tree reaches a pre-specified threshold that is tuned adaptively using a validation sample. Regression trees are among the methods that tend to overfit, resulting in inferior out-of-sample performance. To improve the prediction accuracy of regression trees we consider several ensemble approaches that combine a large number of trees, $\mathcal{T}_1, \mathcal{T}_2, \dots, \mathcal{T}_K$, to yield a single consensus prediction.

The first set of methods are based on boosting, first proposed by Schapire (1990) and Freund (1995) for classification problems, which recursively combines a large number of shallow trees, known as “weak learners”, to form an ensemble of trees with greater stability than a single more complex tree. The first ensemble method we consider is the *gradient boosting machine* (GBM), proposed by Friedman (2001), who extends boosting to regression frameworks. Gradient boosting is initialized by fitting a shallow tree, then a second tree with the same depth is used to fit the residuals of the previous model and the forecasts of these two trees are added together to form a single ensemble prediction. This procedure is repeated sequentially until the total number of iterations K is reached. The goal is to minimize the objective function, based on the square loss

$$\operatorname{argmin}_f \mathcal{L}(\mathbf{r}, f_K(\mathbf{X})) = \operatorname{argmin}_f \|\mathbf{r} - f_K(\mathbf{X})\|^2 \quad (18)$$

Gradient boosting is an additive model that can be expressed as

$$f_K(\mathbf{X}) = \sum_{k=1}^K f_k(\mathbf{X}, c_k, M_k, D, v), \quad (19)$$

where K is the total number of trees and f_k is given by $f_k(\mathbf{X}, v) = f_{k-1}(\mathbf{X}) + v\mathcal{T}_k(\mathbf{X}, c_k, M_k, D)$.

The parameter v controls the learning rate of the boosting procedure that scales the contribution of each tree to the ensemble by a factor of $0 < v < 1$ and prevents the model from overfitting the residuals.

We also consider a *regularized gradient boosting machine* (RGBM), which is an extension to gradient boosting that includes a regularization term in the loss function to control for the complexity of the model and avoid overfitting (Chen and Guestrin, 2016). The updated objective function is

$$\operatorname{argmin}_f \mathcal{L}(\mathbf{r}, f_K(\mathbf{X})) + \Omega(f_K(\mathbf{X})) \quad (20)$$

Refining the definition of a tree as $\mathcal{T}(\mathbf{X}) = w_{q(\mathbf{X})}$, where w is a vector of scores for each region and q is a function assigning each observation to the corresponding leaf, the regularization term Ω is defined as

$$\Omega(f_K(\mathbf{X})) = \gamma M_f + \frac{1}{2} \lambda \|w\|^2 + a \|w\|_1, \quad (21)$$

where the regularization parameter γ is the complexity cost by introducing an additional leaf to a tree and the parameters λ and a control the l_2 and l_1 regularization of the weights. The set of hyperparameters for gradient boosting machine is $\{K, v, D\}$ and $\{K, v, D, \gamma, \lambda, a\}$ for extreme gradient boosting.

The second set of ensemble methods are based on bootstrap aggregating or bagging (Breiman, 1996) that combines many noisy but approximately unbiased models to reduce the variance of the estimates. The baseline bagging procedure estimates $\mathcal{T}_1, \mathcal{T}_2, \dots, \mathcal{T}_B$ based on B different bootstrap samples of the data and then averages their forecasts to obtain a single low-variance model, given by

$$f_B(\mathbf{X}) = \frac{1}{B} \sum_{b=1}^B \mathcal{T}_b(\mathbf{X}, c_b, M_b, D), \quad (22)$$

In bagging, most commonly the regression trees are simply i.d. (identically distributed, but not independent), with the variance of the average estimates, as the number of bootstrapped trees increases, depending on the product of the variance of each tree and the correlation among trees (see Hastie, Tibshirani and Friedman, 2009). We opt to use stationary bootstrap (see Politis and Romano, 1992), instead of standard bootstrap, due to the underlying time series structure of the data.⁵

The first variation of bagging we consider is *random forests* (RF), proposed by Breiman (2001), which aims to reduce the variance of the average estimate by minimizing the correlation among the regression trees in the ensemble. The random forest procedure builds a set of de-correlated trees by considering only a randomly drawn subset of predictors for splitting at each potential branch. This lowers the average correlation among predictions and further reduces the variance relative to bagging. We also

⁵ We would like to thank an anonymous referee for this suggestion.

consider *extremely randomized trees* (ERT), proposed by Geurts, Ernst and Wehenkel (2006), which is an extension to the random forest algorithm. This model has two distinguishing features compared to random forests: the first is that each regression tree is trained using the full sample instead of a bootstrap sample and the second is that the top-down splitting of a tree is randomized. This is achieved by selecting the value where the branch is split randomly, instead of choosing the optimal split value locally for each input variable under consideration. Specifically, the split values are selected from a uniform distribution within the empirical range of the input variable in the training set and the randomized split that gives the highest performance is chosen to split the region. The hyperparameters to be determined using the validation set approach within each iteration of the expanding window, are the number of trees in the ensemble, the depth of the trees and the size of the randomly selected subset of predictors.⁶

Support vector machine⁷ (SVM) regression is a non-parametric technique that learns a non-linear function by mapping linear functions into high dimensional kernel induced feature space. We first consider the *epsilon-insensitive SVM* (e-SVM), as seen in Vapnik (1995), where the objective is to find a function $f(\mathbf{X})$ that deviates from \mathbf{r} by a value no greater than ϵ for each training observation of \mathbf{X} and at the same time is as flat as possible. The e-SVM can be formulated as a Lagrange dual problem by introducing \mathbf{a} and \mathbf{a}^* vectors of non-negative multipliers for each observation of \mathbf{X} . The dual optimization problem for ϵ -SVM becomes:

$$\underset{\mathbf{a}}{\operatorname{argmin}} \frac{1}{2} (\mathbf{a} - \mathbf{a}^*)' \mathbf{Q} (\mathbf{a} - \mathbf{a}^*) + \mathbf{r}' (\mathbf{a} - \mathbf{a}^*) + \epsilon \mathbf{i}' (\mathbf{a} + \mathbf{a}^*) \quad (23)$$

$$\text{s.t. } \mathbf{i}' (\mathbf{a} - \mathbf{a}^*) = 0$$

$$0 \leq a_i, a_i^* \leq C, i = 1, 2, \dots, t$$

where \mathbf{i} is the unity vector, \mathbf{Q} is a $t \times t$ positive semidefinite matrix, $Q_{i,j} \equiv K(x_i, x_j) = \phi(x_i)' \phi(x_j)$ is a non-linear kernel⁸ function and $\phi(x)$ is a transformation that maps x to a high-dimensional space.

⁶ For the boosting-type methods, the number of trees is set to $K = 1 \sim 1000$, the depth to $D = \{1, 2\}$ and the learning rate to $v = \{0.01, 0.1\}$. The regularization parameters for RGBM are set to $\gamma \in [10^{-5}, 10^{-3}]$ and $\lambda, \alpha \in [0.1, 10]$. For bagging-type approaches, the number of trees is set to $K = 1 \sim 500$, the depth to $D = \{1, 2\}$ and the size of the selected subset of predictors to $[2, 44]$.

⁷ The equivalence of support vector machines with the lasso has been examined by Jaggi (2014) and with the elastic net by Zhou, Chen, Song, Gardner, Weinberger and Chen (2015).

⁸ Kozak (2019) applies the kernel trick to the cross section of returns.

Based on the Karush-Kuhn-Tucker conditions, only a certain number of Lagrange multipliers \mathbf{a} and \mathbf{a}^* will assume non-zero values. The observations associated with them have approximation errors equal to or larger than $\epsilon > 0$ and are referred to as support vectors. The parameter ϵ is the trade-off between the sparseness of the representation and closeness to the data, with large values of ϵ resulting to fewer support vectors and thus sparser representation of the solution.⁹ The cost parameter, $C > 0$, controls the penalty imposed on observations that lie outside the epsilon margin, ϵ , and helps to prevent overfitting as it represents the balance between the flatness of $f(\mathbf{X})$ and the extent to which violations to ϵ are tolerated.

One of the problems of e-SVM is choosing the appropriate value for the parameter ϵ . The *nu-support vector machine* (nu-SVM), proposed by Schölkopf, Bartlett, Smola and Williamson (1999) and Schölkopf, Smola, Williamson and Bartlett (2000), is a method that automatically adjusts ϵ , by considering it as part of the optimization problem. The dual optimization problem for nu-SVM becomes

$$\begin{aligned} \underset{\mathbf{a}}{\operatorname{argmin}} \quad & \frac{1}{2}(\mathbf{a} - \mathbf{a}^*)' \mathbf{Q}(\mathbf{a} - \mathbf{a}^*) + \mathbf{r}'(\mathbf{a} - \mathbf{a}^*) & (24) \\ \text{s.t.} \quad & \mathbf{i}'(\mathbf{a} - \mathbf{a}^*) = 0 \\ & 0 \leq a_i, a_i^* \leq C, i = 1, 2, \dots, t \\ & \mathbf{i}'(\mathbf{a} + \mathbf{a}^*) = C\nu \end{aligned}$$

where the parameter $0 \leq \nu \leq 1$ determines the proportion of the number of support vectors kept in the solution with respect to the total number of observations in the dataset. The regression function is then given by

$$f(\mathbf{X}) = \sum_{i=1}^t (a_i^* - a_i)K(x_i, x) \quad (25)$$

We use the radial basis function (RBF) kernel, with parameter $\gamma > 0$, given by $K(x_i, x) = \exp(-\gamma|x_i - x|^2)$. The set of hyperparameters to be selected are $\{C, \gamma, \epsilon\}$ for e-SVM and $\{C, \gamma, \nu\}$ for nu-SVM.¹⁰

⁹ As support vectors are usually only a small subset of the training observations, this characteristic is referred to as the sparsity of the solution.

¹⁰ The cost parameter is set to $C \in [10^{-5}, 10^{-2}]$ and the parameter of the kernel function is set to $\gamma \in [10^{-3}, 10^3]$. The parameter specific to e-SVM, is set to $\epsilon \in [10^{-5}, 10^{-2}]$, while the parameter of nu-SVM is set to $\nu \in [0.1, 1]$.

The final nonlinear method we consider is artificial neural networks, which have been shown to be universal approximators for any continuous function f (Hornik, Stinchcombe and White, 1989; Cybenko, 1989). We focus on the *multilayer perceptron* (MLP), a type of feed-forward neural network where information flows through the function being estimated from \mathbf{X} , through intermediate computations used to approximate f , to the output \mathbf{r} . These models are comprised of a number of layers with multiple nodes in each layer. They consist of an input layer of the predictors, one or more hidden layers, with nodes that transform the predictors using nonlinear activation functions and an output layer that allows a final transformation of the outcome of the hidden layers to form a prediction. For the training and design of the neural networks we follow a similar approach to Gu, Kelly and Xiu (2020). Feed-forward neural networks can be defined as a composition of $h^{(1)}, h^{(2)}, \dots, h^{(L)}$ nonlinear activation functions for each of the L hidden layers of the network $\mathbf{Z}^{(L)} = h^{(L)} \circ \dots \circ h^{(2)} \circ h^{(1)}(\mathbf{X})$, with $\mathbf{Z}^{(l)} = h\left(b_0^{(l)} + \mathbf{W}^{(l)'}\mathbf{Z}^{(l-1)}\right)$, for $l = 1, 2, \dots, L$, where $\mathbf{Z}^{(l)}$ is the l th layer of the network with $m = 1, 2, \dots, M$ nodes, $\mathbf{W}^{(l)}$ is the matrix of weights and $b_0^{(l)}$ is the bias. For the first hidden layer the input is the matrix of predictors, $\mathbf{Z}^{(0)} = \mathbf{X}$, such that $\mathbf{Z}^{(1)} = h\left(b_0^{(1)} + \mathbf{W}^{(1)'}\mathbf{X}\right)$. The results from each hidden layer are aggregated in the output layer; $f(\mathbf{X}) = b_0^{(L+1)} + \mathbf{w}^{(L+1)'}\mathbf{Z}^{(L)}$. We consider three different network architectures based on whether the network has one, two or three hidden layers (MLP1, MLP2 and MLP3). The number of hidden nodes in each layer is selected according to the geometric pyramid rule by Masters (1993). The activation function applied to each node can take various forms. We follow the existing literature and use the rectified linear unit (ReLU) defined as $h(x) = \max(0, x)$, which encourages sparsity in the number of active nodes and avoids the vanishing gradient problem (Nair and Hinton, 2010 and Glorot, Bordes and Bengio, 2011).

The weights and biases of the neural network are estimated by minimizing the following objective function

$$\underset{\mathbf{w}, \mathbf{b}_0}{\operatorname{argmin}} \mathcal{L}(\mathbf{r}, f(\mathbf{X})) = \underset{\mathbf{w}, \mathbf{b}_0}{\operatorname{argmin}} \|\mathbf{r} - f(\mathbf{X})\|^2 \quad (26)$$

The estimates of the parameters of a neural network are solutions of a non-convex optimization problem. The neural network is trained using stochastic gradient descent (SGD). Unlike standard

gradient descent that uses the entire training sample to evaluate the gradient at each iteration of the optimization, SGD evaluates the gradient from a random subset of the data and iteratively minimizes the objective function through back propagation.¹¹ Training a neural network is challenging due to the large number of parameters to be estimated and the nonconvexity of the objective function. To alleviate those concerns we modify the loss function by adding a parameter norm penalty. We consider an elastic net type of penalty (Goodfellow, Bengio and Courville, 2016), based on the l_1 and l_2 norm of the weights. The regularized objective then becomes

$$\operatorname{argmin}_{\mathbf{W}, \mathbf{b}_0} \left[\mathcal{L}(\mathbf{r}, f(\mathbf{X})) + \lambda_1 \|\mathbf{W}\|_1 + \lambda_2 \frac{1}{2} \|\mathbf{W}\|^2 \right], \quad (27)$$

where higher values of the hyperparameters $\lambda_1 > 0$ and $\lambda_2 > 0$ correspond to more regularization. Similarly to the elastic net regression, the penalty term induces sparsity to the weights. We choose to penalize only the weights of the affine transformation at each layer and leave the biases unregularized. Following Gu, Kelly and Xiu (2020), we consider several other forms of regularization. To improve the performance of the model we implement early stopping, batch normalization and forecast averaging. In each iteration of the optimization algorithm the parameter estimates are updated so as to reduce prediction errors in the training sample and then the predictive performance of the model for that iteration is evaluated using data from the validation sample. Early stopping is implemented by stopping the training process prematurely when the validation error no longer decreases. Specifically, the optimization process halts when the maximum number of iterations (100) is reached or if the validation error has not improved for 5 consecutive iterations, preventing overfitting and significantly speeds up the training process. In both cases the parameter estimates of the best performing model are retrieved. Batch normalization (Ioffe and Szegedy, 2015) reduces the variability of the predictors by scaling the input of activations. It was proposed to solve the problem of internal covariate shift in which the distribution of the inputs of the hidden layer change during training, as the parameters of the previous layers change. For each node in each training step, the algorithm cross-sectionally standardizes the output of a previous activation to restore the representation power of the node. This allows to increase

¹¹ The version of SGD we implement is the adaptive moment estimation algorithm (Adam), introduced by Kingma and Ba (2015). Adam computes individual adaptive learning rates for the model parameters using estimates of first and second moments of the gradients.

the stability of the neural network and increase the speed of training. Finally, to reduce the prediction variance of the neural network, due to the stochastic nature of the optimization, we adopt an ensemble approach (Hansen and Salamon, 1990 and Dietterich, 2000) by forming the final prediction from the average of the forecasts from ten models initialized from different random seeds.¹²

2.4. Forecast Combination Methods

The forecast combination approach was originally proposed by Bates and Granger (1969) and can be used as an alternative approach to individual forecasting methods (see Timmermann, 2006 for a comprehensive review). Forecast combinations may be preferred over using forecasts based on individual models, since the latter could suffer from model uncertainty and instability, while combining different models can increase accuracy by including valuable information from each model.

The forecast combinations, denoted by \hat{r}_{t+1}^C , are the weighted averages of return forecasts from m individual models and can be expressed as:

$$\hat{r}_{t+1}^C = \hat{\mathbf{r}} \boldsymbol{\omega}_t \quad (28)$$

where $\hat{\mathbf{r}} = (\hat{r}_{1,t+1}, \hat{r}_{2,t+1}, \dots, \hat{r}_{m,t+1})'$ is the vector of m individual forecasts and $\boldsymbol{\omega}_t = (\omega_{1,t}, \omega_{2,t}, \dots, \omega_{m,t})$ are the combining weights of the individual forecasts at time t . We consider two forecast combination approaches based on the type of models that are averaged. In the first approach we combine $m = p$ number of simple bivariate prediction models based on individual predictors. This simple linear model is based on a single predictor x_i , and the forecast for $t + 1$ using the i th predictor is given by:

$$\hat{r}_{i,t+1} = \hat{a}_i + x_{i,t} \hat{\beta}_i, \text{ for } i = 1, 2, \dots, p.$$

This model is estimated using OLS. In the second approach we combine the forecasts generated by the $m = 23$ machine learning models.¹³

We further differentiate the forecast combinations based on the way $\boldsymbol{\omega}_t$ is computed. Specifically, the *mean combination* (MC) sets the weight $\omega_{i,t} = 1/m$ for $i = 1, 2, \dots, m$ and the *median combination*

¹² The tuning parameters for the neural networks are set to $\lambda_1, \lambda_2 \in [10^{-5}, 10^{-3}]$ the learning rate is set to 0.01 and the parameters for Adam left to the defaults. The maximum epochs are set to 100, patience is set to 5 and the forecasts are an ensemble of 10 networks.

¹³ We would like to thank an anonymous referee for suggesting the forecast combination of machine learning models.

(MDC) is the median of \hat{r}_{t+1} . The *trimmed mean combination* (TMC) sets $\omega_{i,t} = 0$ for 5% of the forecasts with the lowest and highest values and $\omega_{i,t} = 1/(0.9 \times m)$ for the remaining forecasts. These simple forecast averaging schemes do not require a holdout period, q_0 .

For the second type of forecasting methods, the combining weights are computed based on the historical forecasting performance of the individual models over the holdout period, q_0 . Aiolfi and Timmermann (2006) consider a method based on the rank of each model according to the Mean Squared Forecast Error (MSFE). This weighing scheme lets the weights be inversely proportional to the forecast models' rank, $RANK_i$, i.e. $\omega_{i,t} = RANK_{i,t}^{-1} / \sum_{i=1}^m RANK_{i,t}^{-1}$, where the model with the lowest MSFE value gets a rank of 1, the model with the second lowest MSFE value gets a rank of 2 and so forth. Aiolfi and Timmermann (2006) also consider a clustering approach to combine forecasts. Specifically, the forecasts from the individual models are grouped into L equal-sized clusters based on their past MSFE performance, with the first cluster containing the models with the lowest MSFE. Each combination forecast is the average of the individual forecasts contained in the first cluster. This procedure starts with the initial holdout period and then goes through the end of the available OOS period using a rolling window. We consider forecast combinations with two (CL2) and three (CL3) clusters.

The third type of combining methods considered is also based on past performance of the individual models and uses time-varying combination weights. Stock and Watson (2004), proposed the *discounted mean square forecast error* combining method, which uses the following weights $\omega_{i,t} = \varphi_{i,t}^{-1} / \sum_{i=1}^m \varphi_{i,t}^{-1}$, where $\varphi_{i,t} = \sum_{s=R}^{t-1} \psi^{t-1-q_0} (r_{s+1} - \hat{r}_{i,s+1})^2$, for $t = R + q_0, \dots, T$, where R is the in-sample period and ψ is a discount factor, with $0 < \psi \leq 1$. In the case of $\psi < 1$ this method assigns greater weights to recent individual predictive regression forecasts. When $\psi = 1$, then there is no discounting and the equation above produces the optimal combination forecast derived by Bates and Granger (1969) for the case where the individual forecasts are uncorrelated. The values for ψ considered are 1 and 0.9 (DMSFE1 and DMSFE09).

To facilitate the discussion of our findings, Table 1 lists all the models used in this study grouped in six categories.

[Insert Table 1 Here]

3. Data and Sample Splitting

3.1. Data

On the asset side, our dataset consists of 50 individual stocks, 12 bond portfolios and 52 commodities, for a total of $N = 114$ assets. On the predictor side, we consider a common set of $p = 44$ economic indicators. The sample period starts from January 1980 to December 2019, for a period of 40 years (or $T = 480$ monthly observations).

Specifically, we obtain the total returns for the 50 stocks with the highest market capitalization from the Center for Research in Security Prices (CRSP) that have no missing return observations over the full sample period.¹⁴ For the bond portfolios we use the returns of the Fama maturity portfolios, also retrieved from CRSP, where the first 10 portfolios are defined according to maturity in 6-month intervals for up to 60 months, while the remaining are portfolios for maturities between 60 and 120 months and for maturities greater than 120 months. The returns of the 52 commodities are calculated from the prices acquired from the Primary Commodity Price System dataset of the International Monetary Fund (IMF) and are chosen based on having a full price history over the sample period. The majority of the commodities are classified as food or beverages, for a total of 27 assets, while the remaining belong to the following general categories: 11 are metals, 9 are agricultural raw materials, 3 are categorized as fertilizer and 2 are energy commodities. Further details regarding the assets along with their identifiers can be found in Tables A1 to A3 in the appendix, for stocks, bond and commodities respectively.

The choice of predictors draws upon the literature of return predictability, such as Rapach, Wohar and Rangvid (2005), Welch and Goyal (2008), Rapach, Strauss and Zhou (2010), Neely, Rapach, Tu and Zhou (2014), Lima and Meng (2017) for stocks, Ludvigson and Ng (2009), Lin, Wu and Zhou (2018) and Gargano, Pettenuzzo and Timmermann (2019) for bonds and Gargano and Timmermann (2014) and Gao and Nardari (2018) for commodities. The first set of predictors are the factors used in the Fama-French 3- and 5-factor models (Fama and French 1993, 2016) and the 4-factor model of Carhart

¹⁴ We restrict our selection to stocks listed to the NYSE, AMEX, and NASDAQ stock exchanges (exchange codes 1, 2 or 3) and to ordinary common shares (share codes 10 or 11).

(1997). We include the market (MKT), size (SMB), value (HML), profitability (RMW), investment (CMA) and momentum (MOM) factors. As a proxy to market liquidity, we use the Pastor and Stambaugh (2003) aggregate innovation in liquidity measure (LIQ). We also consider variables from Welch and Goyal (2008), which are the dividend-price ratio (DP), dividend yield (DY), earnings-price ratio (EP), dividend-payout ratio (DE), stock variance (SVAR), book-to-market ratio (BM), net equity expansion (NTIS), Treasury bill rate (TBL), long-term yield (LTY), long-term return (LTR), term spread (TMS), default yield spread (DFY) and default return spread (DFR). Additional predictors related to the bond market are the ICE BofA US corporate total return index (BAML) and the Cochrane and Piazzesi (2005) factor (CP). In the set of predictors, we include economic indicators such as the industrial production (INDPRO), the money stock M1 (M1), the consumer price index (CPI), the producer price index (PPI), capacity utilization (CAP), the unemployment rate (UNRATE), the inventory-sales ratio (IR), housing starts (HOUST) and total consumer credit (Credit). As a proxy for the overall commodity market, we use the S&P Goldman Sachs commodity total return index (GSCI) and include four commodity currencies (Chen, Rogoff and Rossi, 2010), the Australian dollar-US dollar (USAU), the Canadian dollar-US dollar (USCA), the Indian rupee-US dollar (USIN) and the New Zealand dollar-US dollar (USNZ). Finally, we consider eight additional variables, namely the Chicago Board Options Exchange volatility index (VXO), the University of Michigan consumer sentiment index (UMSENT), the Chicago Fed national activity index (CFNAI), Kilian’s (2009) real economic activity index (REA), the Philadelphia Fed business outlook survey current and future activity indices (GAC and GAF) and the macroeconomic and financial uncertainty indices (UMacro and UFin respectively), proposed by Jurado, Ludvigson and Ng (2015) and Ludvigson, Ma and Ng (2015). The candidate variables and their sources are given in Table A4 in the appendix, while descriptive statistics of the predictors are reported in Table A5.

3.2. Sample Splitting and Hyperparameter Tuning

We generate out-of-sample forecasts of asset returns by employing a recursive forecasting scheme. The total sample, T , is divided into the in-sample part, R and the out-of-sample part, $Q = T - R$. The first q_0 forecasts of the out-of-sample period, Q , serve as the hold-out period for the forecast combination methods that require it. The initial size of the recursive window, used to estimate the individual

forecasting models, is set to $R = 180$ monthly observations (or 15 years, from January 1980 to December 1994) and the hold-out period, for the forecast combination methods that require it, is set to $q_0 = 60$ observations (or 5 years, from January 1995 to December 1999). The window expands by one observation at a time, leading to an out-of-sample size of 240 monthly observations from January 2000 to December 2019.

The machine learning models used to generate the return forecasts rely on hyperparameter tuning. The choice of hyperparameters controls the amount of model complexity and is critical for the performance of the model. Specifically, we adopt a validation sample approach similar to Gu, Kelly and Xiu (2020), in which the optimal set of values for the tuning parameters is selected in the validation sample. One of the advantages of using this approach over k -fold cross validation is that we maintain the temporal ordering of the data. Specifically, in each iteration of the expanding window, the in-sample is split into two disjoint periods, the training subsample, consisting of 80% of the observations, with the remaining observations belonging to the validation subsample. In the training subsample the model is estimated for several sets of hyperparameters. The second subsample is used to select the optimal set of tuning parameters, by constructing forecasts, using the model estimates from the training sample for the respective hyperparameter set, for the observations in the validation sample. The optimal set of hyperparameters is chosen so as to minimize the Mean Squared Error over the validation subsample. Once the optimal set of hyperparameters is chosen, the model is refitted using all data from the in-sample period and the estimates of the model parameters are kept to construct the forecasts. The predictors are standardized for all methods; first separately for observations within each of the training and validation subsamples, during the selection of the optimal set of hyperparameters and then using all observations of the in-sample window, when estimating the parameters of the model. For a detailed description of cross-validation see Friedman, Hastie and Tibshirani (2009). Due to the computational cost of training these methods, we avoid recursively fitting the models in each iteration. Instead, we estimate the parameters once a year and retain those estimates to derive the remaining forecasts for that year.

4. Optimal Asset Allocation

Consider an investor who allocates her wealth among N individual assets with a $N \times 1$ portfolio weight vector: $\mathbf{w} = (w_1, w_2, \dots, w_N)$. The initial wealth is normalized to 1. The benchmark strategy is the naive diversification rule of an equal-weighted portfolio, where $w_j = 1/N$, for $j = 1, 2, \dots, N$. The objective of the main framework is to optimize the trade-off between risk and return. As a basic measure of portfolio risk the standard deviation of the portfolio (Markowitz, 1952), is used. The mean-variance (MV) optimization problem is:

$$\underset{\mathbf{w}}{\operatorname{argmin}}[\gamma \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - \mathbf{w}' \hat{\mathbf{r}}] \quad (29)$$

where $\boldsymbol{\Sigma}$ is the $N \times N$ covariance matrix of asset returns, $\hat{\mathbf{r}} = (\hat{r}_{1,t+1}, \hat{r}_{2,t+1}, \dots, \hat{r}_{N,t+1})$ is the $N \times 1$ vector of return forecasts for each asset and γ is the coefficient of relative risk aversion. As an alternative to the $1/N$ benchmark, portfolios using historical average forecasts are considered. The two benchmarks are compared to portfolios based on forecasts generated by multivariate prediction models. All portfolio models include short-selling and leverage constraints to avoid implausible positions. The first constraint sets an upper bound to the sum of the portfolio weights, $\mathbf{w}' \mathbf{i}_N = h$, where \mathbf{i}_N is an N -vector of ones and h denotes the maximum leverage, for example $h = 1$ ensures that the portfolio weights sum up to one, while $h = 1.5$ indicates that the investor cannot borrow more than 50% of total wealth. The second constraint sets a lower bound to the weight of each asset as $w_j \geq 0$, with $j = 1, \dots, N$, which leads to portfolios without short selling (Jagannathan and Ma, 2003). The portfolio return at $t + 1$ can then be computed as:

$$r_{P,t+1} = \hat{\mathbf{w}}_t' \mathbf{r} + (1 - \hat{\mathbf{w}}_t' \mathbf{i}_N) r_{f,t+1} \quad (30)$$

where $\mathbf{r} = (r_{1,t+1}, r_{2,t+1}, \dots, r_{N,t+1})$ is an N -vector of risky asset returns. In the case of $h = 1$, the portfolio return is equivalent to $r_{P,t+1} = \hat{\mathbf{w}}_t' \mathbf{r}$. For the mean-variance optimization framework, two approaches are used to estimate the covariance matrix. The first is the graphical lasso, which is a static estimator that derives a sparse version of the covariance matrix. The second is a dynamic estimator based on the DCC GARCH. The estimates of the covariance matrix, $\hat{\boldsymbol{\Sigma}}$, for assets $i = 1, 2, \dots, N$, are based on a rolling window of 120 observations.

The graphical lasso algorithm, proposed by Friedman, Hastie and Tibshirani (2008), estimates the sparse precision matrix (inverse of the covariance matrix), using the l_1 (lasso) penalty to enforce sparsity. The graphical lasso problem is to maximize the following penalized log likelihood:

$$\log(\det\mathbf{\Theta}_t) - \text{tr}(\mathbf{S}_t\mathbf{\Theta}_t) - \rho\|\mathbf{\Theta}_t\|_1 \quad (31)$$

where \mathbf{S}_t is the sample covariance matrix and $\rho \geq 0$ is a tuning parameter controlling the amount of regularization. Here, $\mathbf{\Theta}_t = \mathbf{\Sigma}_t^{-1}$, with entries $\theta_{i,j}$, is the $N \times N$ inverse of the covariance matrix and $\|\mathbf{\Theta}_t\|_1$ is the l_1 norm of $\mathbf{\Theta}_t$ – the sum of the absolute value of the elements $\theta_{i,j}$. The penalty parameter ρ is chosen by the validation sample approach, to make the value of $\log(\det\mathbf{\Sigma}_{1,t}^{-1}) - \text{tr}(\mathbf{\Sigma}_{2,t}\mathbf{\Sigma}_{1,t}^{-1})$ large, where $\mathbf{\Sigma}_{1,t}$ is the covariance matrix estimated using the training set and $\mathbf{\Sigma}_{2,t}$ is the covariance estimated over the validation set.

For the dynamic conditional correlation GARCH model, proposed by Engle (2002), the one-period ahead covariance based on the DCC GARCH model evolves according to:

$$\mathbf{\Sigma}_{t+1} = \mathbf{D}_{t+1}\mathbf{R}_{t+1}\mathbf{D}_{t+1} \quad (32)$$

where \mathbf{D}_{t+1} is an $N \times N$ diagonal matrix with conditional standard deviation $\hat{\sigma}_{i,t+1}$ on the i th diagonal element and \mathbf{R}_{t+1} is the $N \times N$ correlation matrix, with ones on the diagonal and conditional correlations in the off-diagonal. The estimation of the DCC GARCH has two steps. The first step involves estimating the diagonal elements of the conditional standard deviation matrix, \mathbf{D}_{t+1} , where we estimate the conditional standard deviation, $\hat{\sigma}_{i,t+1}$, of the i th asset is using a GARCH(1,1) model. The second step involves the estimation of the conditional correlation matrix, \mathbf{R}_{t+1} . Removing the conditional mean from the N series of asset returns yields the residuals, $\boldsymbol{\epsilon}_{t+1}$ and the standardized residuals, \mathbf{u}_{t+1} , can be obtained using the conditional standard deviation matrix, \mathbf{D}_{t+1} : $\mathbf{u}_{t+1} = \mathbf{D}_{t+1}^{-1}\boldsymbol{\epsilon}_{t+1}$. The conditional correlation structure then is:

$$\mathbf{Q}_{t+1} = (1 - a - b)\bar{\mathbf{Q}} + a\mathbf{u}_t\mathbf{u}_t' + b\mathbf{Q}_t \quad (33)$$

$$\mathbf{R}_{t+1} = \mathbf{Q}_{t+1}^{*-1}\mathbf{Q}_{t+1}\mathbf{Q}_{t+1}^{*-1}$$

where $\bar{\mathbf{Q}}$ is the unconditional covariance of the standardized residuals and \mathbf{Q}_{t+1}^* is a diagonal matrix composed of the square root of the diagonal elements of \mathbf{Q}_{t+1} .

4.1. Portfolio Performance

In this section, we assess the economic value of using return forecasts in asset allocation. The portfolios are constructed recursively using the related return and covariance estimates in each iteration, starting in January 2000. The buy-and-hold portfolio returns are calculated for the period of one month and the portfolio is rebalanced monthly until the end of the evaluation period (December 2019). Each portfolio is computed for different combination of weight constraints: unleveraged long-only portfolios ($0 \leq w_j \leq 1$) and leverage restricted to 50% of wealth ($0 \leq w_j \leq 1.5$). Two types of investors are considered based on different values of the coefficient of risk aversion, $\gamma = 2$ for an aggressive investor and $\gamma = 10$ for a conservative investor. Similar values of risk aversion are used in Callot, Kock and Medeiros (2017) and DeMiguel, Garlappi and Uppal (2009).

The performance of the portfolios is evaluated over the out-of-sample period using the average return of the portfolio and the out-of-sample Sharpe ratio. The Sharpe ratio (SR) is calculated as the fraction of the out-of-sample excess return (average realized return less the risk-free rate) divided by the standard deviation of the out-of-sample portfolio returns: $SR = (\bar{r}_p - r_f) / \hat{\sigma}_p$, where $\bar{r}_p = 1/(Q - q_0) \sum_{t=1}^{Q-q_0} r_{p,t}$ is the average realized return of the portfolio over the out-of-sample period, r_f is the risk free rate and $\hat{\sigma}_p$ is the standard deviation of the portfolio excess returns over the out-of-sample period. Additionally, we report portfolio turnover, defined as a measure of the amount of trading required to implement a particular strategy. Following DeMiguel, Garlappi and Uppal (2009), the portfolio turnover is defined as the average absolute change of the portfolio weights over the $Q - q_0$ rebalancing periods across the N assets and is given as follows:

$$PT_P = \frac{1}{Q - q_0 - 1} \sum_{t=1}^{Q-q_0-1} \sum_{j=1}^N |w_{j,t+1} - w_{j,t}| \quad (34)$$

where $w_{j,t+1}$ is the weight in asset j at time $t + 1$ and $w_{j,t}$ is the weight in asset j at time before rebalancing at $t + 1$. When a portfolio is rebalanced at $t + 1$, $|w_{j,t+1} - w_{j,t}|$ denotes the magnitude of trading asset j .

Table 2 presents the results of the portfolio evaluation based on average return. The first row gives the average return of the $1/N$ portfolio, which is 5.73% across all panels since derivation of the weights

for this strategy does not involve any optimization or estimation. The remaining rows of the table report the average return of the mean-variance portfolio based on the historical average, KS and alternative forecasting models. These results vary based on the estimator of the covariance matrix, the type of investor and combination of weight constraints.

[Insert Table 2 Here]

Focusing on the results of the first panel, which are based on the graphical lasso estimates of the covariance matrix, the average return of the HA portfolio for an aggressive investor ($\gamma = 2$), is 4% for $w_j \in [0,1]$ and 7.12% for $w_j \in [0,1.5]$. On the other hand, the average return of a conservative investor ($\gamma = 10$), for weight constraints $0 \leq w_j \leq 1$ is 6.06% and when $0 \leq w_j \leq 1.5$ it is 9.16%. For the portfolios based on the KS forecasts, the average returns range from 24.96% to 37.38% for an aggressive investor and from 20.43% to 27.29% for a conservative investor. Turning to the results of the second panel, HA portfolios of a conservative investor benefit more when using the DCC estimator instead of the graphical lasso, while for the KS portfolios, the sparse estimator yields higher average returns across different combinations of weight constraints and levels of risk aversion.

Overall, forecast combinations of machine learning models tend to achieve the best performance in terms of average return, especially for an aggressive investor. Individual models that perform well are those based on shrinkage methods, partial least squares, sparse partial least squares and a neural network with a single hidden layer. Relaxing the leverage constraint consistently improves the performance of the models. The average return of the mean-variance portfolios, based on the graphical lasso, across all alternative predictive regressions, for an aggressive investor, is between 4.41% (CL2) and 27.08% (DMSFE09_{ML}) for the case when no short sales or leverage is allowed. For a 50% leverage constraint, the respective returns are 7.05% and 38.93%. In the case of a conservative investor, for weight constraints $0 \leq w_j \leq 1$ the average return is between 6.40% (CL2) and 20.21% (MCP) and when leverage is allowed, returns range from 9.81% (MDC and CL2) to 27.48% (MCP). Comparing the results between the two specifications for the covariance matrix, the graphical lasso outperforms the DCC, except in the case of portfolios based on certain forecast combinations of bivariate prediction models and for portfolios based on nu-SVM. The majority of the machine learning models and their

combinations outperform the equal-weighted allocation and the HA benchmarks, while portfolios based on the KS model prove to be harder benchmarks to overcome. The model that fails to outperform the EW portfolio is the neural network with three hidden layers, for portfolios based on the dynamic covariance estimator and $\gamma = 10$, while the HA portfolio yields higher average return than RGBM and a neural network with three hidden layers for portfolios of a conservative investor based on the DCC covariance.

Table 3 reports the results for the annualized Sharpe ratio. Our findings indicate that between 18 and 19 models for an aggressive investor and 20 to 25 models for a conservative investor outperform the EW benchmark, with a ratio of 0.78, depending on the weight constraints and estimates of the covariance matrix. The simple forecast combination methods, along with ensemble methods and support vector machines fail to outperform the EW in the majority of specifications. The Sharpe ratio for the HA portfolio with $\gamma = 2$ varies between 0.25 and 0.37 and is outperformed by 37 to 39 of the models, while for a conservative investor the ratio is between 0.59 and 0.74, with 32 to 37 models generating better performance. More importantly, the models that consistently outperform both benchmarks are the forecast combinations of machine learning models. For example, for an aggressive investor with a 50% leverage constraint, the Sharpe ratio ranges from 0.95 (Rank_{ML}) to 1.28 ($\text{DMSFE09}_{\text{ML}}$).

[Insert Table 3 Here]

The KS portfolio generates ratios from 0.96 to 1.16 for an aggressive investor and between 1.14 and 1.25 for a conservative investor. The Sharpe ratio for portfolios with relative risk aversion of 2, is between 0.29 and 1.22 for unleveraged and long only portfolios and 0.33 to 1.28 for a 50% leverage constraint. Among the best-performing portfolios are those with return forecasts generated by machine learning model combinations, PLS, Ridge and MLP1. Based on different weight constraints, the Sharpe ratios for a conservative investor are higher compared to those of a more aggressive investor, with values from 0.37 to 1.25 when $w_j \in [0,1]$ and 0.38 to 1.27 for $w_j \in [0,1.5]$. Shrinkage methods, PLS, MLP1 and ML forecast combinations are among the models that yield the highest ratios across all

weight constraints. Finally, portfolios based on graphical lasso tend to generate slightly higher Sharpe ratios than the DCC GARCH.

The first part of Figure 1 presents the cumulative return series of the EW and HA benchmarks and the eight portfolios based on forecast combinations of machine learning models of an aggressive investor with 50% leverage. The eight strategies significantly outperform the EW and HA benchmarks throughout the out-of-sample period, especially after 2002-2003. The cumulative return series of the forecast combinations exhibit overall an upward trend and experience a slight downward trend during the global financial crisis. The discounted mean square forecast error scheme with $\psi = 0.9$ displays superior gains compared to the other models after about 2013 and achieves the highest end-of-period value. The worst performing forecast combination is based on the rank scheme, while the cumulative returns of the remaining models evolve in a similar way over time.

[Insert Figure 1 Here]

The second part of Figure 1 depicts the cumulative return of the EW and HA benchmarks and the eight portfolios based on forecast combinations of machine learning models of a conservative investor without any leverage. Overall, the performance of the eight portfolios of the conservative investor is more stable over time compared to those of an aggressive investor, however, the end of period value is approximately half that of the portfolios with $\gamma = 2$. The eight portfolios considerably outperform the HA and EW benchmarks, especially after the global financial crisis, with the portfolio based on a cluster combination scheme of two clusters yielding the highest end-of-period-value.

We report the annualized average turnover of the portfolios as a percentage in Table 4. The portfolio with the lowest turnover (except for the $1/N$) is the HA, with values between 6.55% and 10.58%, in the case of graphical lasso and from 16.12% to 39.31% for the DCC. The majority of the portfolios of an aggressive investor have a higher turnover than those of a conservative investor. Additionally, most of the portfolios based on the DCC have higher turnover, the exceptions being shrinkage methods, PLS and certain machine learning combinations in the case of $\gamma = 2$ and neural networks for most of the combinations of portfolio parameters.

[Insert Table 4 Here]

Among the multivariate prediction models, the simple forecast combinations generate the lowest values, with turnover between 12.67% (MDC) and 164.42% (Rank) for portfolios based on sparse covariance estimates and from 27.19% (MDC) to 191.68% (Rank) for the dynamic covariance estimator. Machine learning models have a turnover in the range of 57.35% and 276.97%, while combinations of ML models generate turnover from 147.25% to 264.40%. Portfolios based on extremely randomized trees have the lowest turnover, while a neural network with three hidden layers consistently produces the highest average turnover across all models.

4.2. Asset Selection and Portfolio Weights

The percentage of wealth invested in each asset class throughout the out-of-sample period is reported in Table 5 for portfolios based on the graphical lasso. In the case of the EW portfolio 43.86% is invested in stocks, 10.53% in bonds and 45.61% in commodities. Inspecting the Table, we observe that the remaining models allocate little to no wealth to bonds. Focusing on the results for unleveraged portfolios, the HA and simple forecast combination methods tend to invest primarily in stocks with only a small percentage (approximately 5% to 10% depending on the degree of risk aversion) invested in commodities. The KS model and shrinkage methods show similar results, placing approximately equal weights between stocks and commodities. Portfolios based on dimensionality reduction and non-linear methods invest on average 70% in stocks and 30% on commodities, while in the case of forecast combinations of machine learning models the split is 60% to stocks and 40% to commodities.

[Insert Table 5 Here]

Figure 2 presents the average contribution of each asset to the portfolio across the OOS period. The 30 assets with the highest contribution to the portfolio of an aggressive and a conservative investor are reported. The assets that contribute most to the portfolio of an aggressive investor are 20 stocks and 10 commodities, while for the conservative investor there are 18 stocks and 12 commodities. The commodities with the highest contributions according to both investors are bananas, lead, nickel, oats, oil, pork, silver, sugar (free market), uranium and urea, while the portfolio of a conservative investor places significant weights also on iron and orange juice. Portfolios based on the HA, simple forecast combinations, unsupervised dimensionality reduction methods, bagging-type ensembles and support

vector machines tend to focus heavily on certain stocks such as Walmart (WMT) and Altria Group (MO), while the remaining methods allocate wealth across assets more evenly.

[Insert Figure 2 Here]

4.3. Portfolio Performance During Business Cycles

In this section we examine the performance of the portfolios during NBER dated recessions and expansions. Table 6 reports the results for the average return. For the EW, HA and the majority of the simple forecast combinations the average return is positive during expansions but negative during recessions, the exception being the rank combination scheme for an aggressive investor that generates higher and positive returns during recessions. For the forecast combinations of machine learning models, the portfolios of an aggressive investor and those of a conservative investor without any leverage also yield better performance during expansions, however, compared to the simple forecast combinations the returns during both subperiods are positive. This pattern persists for most of the dimensionality reduction or non-linear methods and is reversed for shrinkage methods, where the models yield higher return during recessionary subperiods.

[Insert Table 6 Here]

We report the results in terms of Sharpe ratio in Table 7. Our findings suggest that the portfolios of a conservative investor are consistently higher during expansions than in recessions, with most models generating positive Sharpe ratios during both subperiods, except for EW, HA and forecast combinations of bivariate regression models that have negative ratios during recessions. For an aggressive investor the performance based on the Sharpe ratio is qualitatively similar to that based on average return, with the exception of forecast combinations of machine learning models that tend to have higher ratios during recessions than expansions.

[Insert Table 7 Here]

4.4. Variable Importance

In this section we present the contribution of each feature to the learning process. Gu, Kelly and Xiu (2020) highlight the importance of quantifying the influence of each predictor as a way to interpret machine learning models. The measure of variable importance is calculated as the change in the out-of-sample R^2 , as defined in Campbell and Thompson (2007), from setting the values of a predictor to zero

within each iteration of the out-of-sample period and then averaging these values to obtain a single variable importance measure for each predictor of a particular model. To gain further insight as to which features contribute most for each asset class, we further summarize the results by averaging the variable importance measure, of a specific feature and model, for all assets depending on whether they are a stock, bond, or a commodity. Figure 3 reports the variable importance, for the three asset classes and the 20 most influential predictors of each model. Variables are ranked so that those with the highest variable importance are on top and the lowest are at the bottom, with the ranking starting from the first model (KS).

[Insert Figure 3 Here]

For stocks and the KS model, the most important features are the industrial production, market return, dividend-price ratio, capacity utilization, earnings-price ratio and term spread. The important variables for shrinkage methods are similar to those of the KS model, however, the MKT variable becomes less important, while financial uncertainty is more influential. For dimensionality reduction methods, influential variables for PCA- and PLS-type of methods include financial uncertainty, the payout ratio, stock variance and real economic activity index, while for ICA and RICA the dividend-price ratio and the market are the most important features. Non-linear machine learning methods tend to use a broader set of predictors, the exception being the neural network with a single hidden layer that focuses on dividend-price ratio and the market proxy. The variable importance results for the bond market indicate that most models use the information from the full set of predictors, except for RICA and MLP1 that are skewed towards the dividend-price ratio. For the commodities, influential predictors for most shrinkage methods include the market, dividend-price ratio, industrial production, capacity utilization, CFNAI and GSCI. Turning to the dimensionality reduction methods, ICA and RICA focus especially on the market, DP and TMS. On the other hand, the remaining methods draw information from a wider set of variables. Non-linear methods again appear to not focus on a particular predictor, except for the market and dividend-price ratio variables in the case of MLP1.

5. Further Performance Evaluation and Robustness Checks

In this section we examine portfolio performance when transactions costs are introduced, for different rebalancing frequency and for time-varying risk aversion parameter. We also perform several robustness checks, such as employing an alternative shrinkage estimator of the covariance matrix, including power series and interactions in the predictor set and using the Huber loss function as an objective function.

5.1. Effects of Transaction Costs

The estimation of transaction costs is based on portfolio turnover. Given a transaction cost of c , the trading cost of the entire portfolio is $c \sum_{j=1}^N |w_{j,t+1} - w_{j,t}|$. The return of the portfolio after transaction costs is as follows:

$$r_{P,t+1}^{TC} = (1 + r_{P,t+1}) \left(1 - c \sum_{j=1}^N |w_{j,t+1} - w_{j,t}| \right) - 1 \quad (35)$$

We follow Olivares-Nadal and DeMiguel (2018) who show that incorporating a l_p transaction cost term in the mean-variance portfolio problem may help to reduce the impact of estimation error. Here we consider the case of proportional transaction costs and modify the mean-variance optimization problem by adding a l_1 transaction cost term, which is equivalent to assuming that transaction costs are proportional to the amount traded. The new constrained optimization problem becomes

$$\begin{aligned} & \underset{\mathbf{w}}{\operatorname{argmin}} [\gamma \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - \mathbf{w}' \hat{\mathbf{r}} + c \|\mathbf{w} - \mathbf{w}_0\|_1] \\ & \text{s.t. } \mathbf{w}' \mathbf{i}_N = h \text{ and } w_j \geq 0, \text{ with } j = 1, \dots, N \end{aligned} \quad (36)$$

where c is the transaction cost parameter and \mathbf{w}_0 are the weights of the portfolio from the previous period before rebalancing. For the first period of the expanding window the weights \mathbf{w}_0 are initialized based on the original mean-variance allocation. The transaction costs are set to $c = 50$ bps for each asset. When $c = 0$ the above optimization problem becomes equivalent to the one in (29).

Panel A of Table 8 reports the portfolio performance in terms of Sharpe ratio, for transaction costs of 50 bps using the penalized portfolio objective function with a covariance matrix estimated by the graphical lasso. When transaction costs are introduced, the portfolios of a conservative investor are

affected more than those of an aggressive investor, with several models failing to outperform the EW portfolio and the number of models with Sharpe ratios lower than the EW and HA portfolios increasing.

[Insert Table 8 Here]

More in detail, the Sharpe ratio for the HA portfolio of an aggressive (conservative) investor is 0.36 (0.57) and 0.42 (0.66) for unleveraged and leveraged positions, while the respective values for the KS portfolios range from 0.64 to 0.68. The Sharpe ratio for portfolios of an aggressive investor based on alternative predictive models, is between -0.15 (MLP3) and 0.83 (DMSFE09_{ML}) for unleveraged portfolios and -0.24 (MLP3) to 0.82 (DMSFE09_{ML}) for a 50% leverage constraint. Turning to the results for the conservative investor, for $w_j \in [0,1]$ the Sharpe ratio is between -0.23 (MLP3) and 0.80 (Rank_{ML}) or from -0.25 (MLP3) to 0.83 (Rank_{ML}) when leverage is allowed. Overall, models based on machine learning forecast combinations yield the highest ratios, while those based on RICA and a neural network with three hidden layers generate the lowest Sharpe ratios. Turning to the results for average turnover (Panel B), the added penalty to the mean-variance objective has the greatest effect on the HA and simple forecast combination portfolios, which produce lower turnover compared to the remaining strategies. The turnover for portfolios with $\gamma = 2$ ranges from 15.30% (ERT) to 260.69% (Ridge) and for $\gamma = 10$, the respective values are 11.01% (ERT) and 239.16% (GBM).

The cumulative returns of the EW, HA and forecast combinations of machine learning models for an aggressive investor are plotted in the first part of Figure 4. The cumulative returns of all series fluctuate much more compared to the case without transaction costs (Figure 1); this volatility becomes more pronounced around the two crises periods. After the global financial crisis, the portfolios based on machine learning forecast combinations show significant gains over the two benchmarks, however, all models exhibit similar behavior over time with the discounted mean square forecast error scheme with $\psi = 0.9$ having the highest end-of-period value. The cumulative return graphs of the eight forecast combinations along with the two benchmarks for portfolios with $\gamma = 10$ and transaction costs are presented in the second part of Figure 4. Overall, the difference in behavior of the cumulative return series across models is more pronounced, compared to when $\gamma = 2$. The portfolios based on cluster combination forecasts outperform all other models (especially after the dot-com bubble) and achieve

the highest end-of-period values. The performance of the remaining forecast combinations is superior to the EW portfolio after the global financial crisis.

[Insert Figure 4 Here]

5.2. Alternative Rebalancing Frequencies

In the analysis detailed above, we demonstrate that for monthly rebalancing, portfolios based on machine learning forecast combinations generate the highest performance. In this Section we examine the effects to portfolio performance when the rebalancing frequency of the portfolios is reduced from monthly to quarterly or annual. We report the results for the two alternative rebalancing frequencies, for the average return and the Sharpe ratio in Tables A6 and A7, respectively. Overall, the results indicate that machine learning models favor higher rebalancing frequencies. Reducing the frequency with which a portfolio is rebalanced leads to considerably lower performance for all models, in terms of both measures. Specifically, the results based on average return show that the best performing portfolio in the case of quarterly rebalancing is based on a shallow neural network. However, when rebalancing frequency is reduced to annual, the best performing portfolios become sparse PCA and a neural network with two hidden layers for the aggressive and conservative investors, respectively. On the contrary, the results based on the Sharpe ratio show that all alternative models fail to outperform the equally weighted allocation.

5.3. Time-Varying Parameter of Risk-Aversion

In the previous section we used values for the parameter of risk aversion that remain fixed across the portfolio formation period. Here we employ a new measure of time-varying risk aversion, named RAbex, proposed by Bekaert, Engstrom, Xu (2021), which is derived from observable financial variables, such as earnings yield, corporate return spread (Baa-Aaa), term spread (10yr-3mth), equity return realized variance, corporate bond return realized variance, and equity risk-neutral variance.¹⁵ The results for the average return, Sharpe ratio and turnover are reported in Table A8 in the Appendix. In terms of average return and Sharpe ratio, the best performing models are those based on forecast combinations of machine learning models, followed by the KS model, a shallow neural network (MLP1)

¹⁵ We would like to thank an anonymous referee for their suggestion.

and shrinkage methods. The best performing forecast combination is DMSFE09_{ML} with AR (SR) 26.56% (1.30) for unleveraged positions and 37.04% (1.33) when leverage is allowed. The KS model, the neural network with three hidden layers and shrinkage methods exhibit the highest average turnover. Overall, the performance of portfolios with time-varying risk-aversion is between that of the portfolios with $\gamma = 2$ and $\gamma = 10$ from the previous section.

5.4. Shrinkage Covariance Estimator

As an alternative to the graphical lasso used in the previous section, we consider portfolios based on the linear shrinkage estimator of the covariance matrix proposed by Ledoit and Wolf (2004). Specifically, the sample covariance matrix is shrunk towards a one-parameter matrix, where all the variances are the same and all covariances are zero. The results, presented in Table A9 in the appendix, remain qualitatively similar to the graphical lasso estimator. The quantitative difference in performance between the two estimators is small and any small benefits from using the shrinkage over the sparse estimator are mixed and depend on the model used to generate the return forecasts.

5.5. Power-Series and Interactions

Our findings so far have shown that shrinkage methods tend to outperform the majority of the non-linear methods, with the exception of a shallow neural network. In this respect, we also examine how the performance of the portfolios is affected when the predictor set of the shrinkage methods is replaced with one that includes higher polynomials and two-way interactions of the original variables. The results are reported in Appendix Table A10. Our findings suggest that the performance of the portfolios is significantly reduced across combinations of weight constraints and values for the parameter of risk aversion. Moreover, shrinkage methods with non-convex penalties are more affected than those based on convex penalties.

5.6. Alternative Loss Function

The presence of extreme observations in financial and economic variables can undermine the stability of the models due to the emphasis placed by least squares on large errors. In some cases, it is possible to improve model performance by using an alternative to the least squares objective function in equation (3). Similarly to Gu, Kelly and Xiu (2020), we consider the Huber loss function which is a hybrid of the least squares loss for relatively small errors and the absolute loss for relatively large errors.

Specifically, we replace the least square loss for the shrinkage methods with convex penalties (Ridge, Lasso and Elastic Net) and boosting ensembles (GBM and RGBM). The optimal parameter of the Huber loss function, which interpolates between the mean and the median regression, is chosen using the validation sample approach described in Section 3.2. The results are presented in Table A11 in the Appendix. Our findings suggest that shrinkage methods tend to generate higher average returns and Sharpe ratios when using the least square loss. However, the Huber loss function provides small economic benefits to portfolios based on forecasts from boosting methods for the case of a conservative investor ($\gamma = 10$).

6. Conclusion

This study sets out to explore whether return forecasts generated by machine learning add value to portfolios consisting of stocks, bonds and commodities. The portfolios are constructed based on the proposed models and their performance is compared to that of the equal-weighted portfolio and a mean-variance portfolio based on the historical average. The majority of the portfolios utilizing return forecasts outperform the $1/N$ and historical average benchmarks in terms of average returns and Sharpe ratio, with forecast combinations of machine learning models yielding the highest performance. Portfolios with leverage generate higher average return than the unleveraged allocations. There are no major changes when comparing either performance measure across different specifications of the covariance matrix. When transaction costs are introduced and a mean-variance objective function with a transaction cost penalty is utilized, the results for the monthly-rebalanced portfolios continue to favor forecast combinations of machine learning methods even though the performance of the portfolios is considerably reduced.

References

- Aiolfi, M. and A. Timmermann. "Persistence in Forecasting Performance and Conditional Combination Strategies." *Journal of Econometrics*, 135 (2006), 31-53.
- Babii, A.; E. Ghysels and J. Striaukas. "Estimation and HAC-based Inference for Machine Learning Time Series Regressions", (2019), Available at SSRN: <https://ssrn.com/abstract=3503191>.

- Bai, J. and S. Ng. "Forecasting Economic Time Series using Targeted Predictors." *Journal of Econometrics*, 146 (2008), 304-317.
- Bates, J. M. and C. W. Granger. "The Combination of Forecasts." *Journal of the Operational Research Society*, 20 (1969), 451-468.
- Bekaert, G.; E. C. Engstrom and N. R. Xu. "The Time Variation in Risk Appetite and Uncertainty", (2021), *Management Science*, forthcoming.
- Bianchi, D. and M. Guidolin. "Can Long-Run Dynamic Optimal Strategies Outperform Fixed-Mix Portfolios? Evidence from Multiple Data Sets." *European Journal of Operational Research*, 236 (2014), 160-176.
- Bianchi, D.; M., Büchner and A. Tamoni. "Bond Risk Premia with Machine Learning" *The Review of Financial Studies*, 34 (2020), 1046-1089.
- Breiman, L. "Bagging Predictors." *Machine Learning*, 24 (1996), 123-140.
- Breiman, L. "Random Forests." *Machine Learning*, 45 (2001), 5-32.
- Breiman, L.; J. Friedman; R. Olshen and C. Stone. "Classification and Regression Trees." *Wadsworth Inc*, (1984).
- Callot, L. A.; A. B. Kock and M. C. Medeiros. "Modeling and forecasting large, realized covariance matrices and portfolio choice." *Journal of Applied Econometrics*, 32 (2017), 140-158.
- Callot, L.; M. Caner; A. O. Önder and E Ulaşan. "A nodewise regression approach to estimating large portfolios." *Journal of Business & Economic Statistics*, (2019), 1-12.
- Campbell, J. Y. and S. B. Thompson. "Predicting Excess Stock Returns Out of Sample: Can Anything Beat the Historical Average?" *The Review of Financial Studies*, 21 (2007), 1509-1531.
- Carhart, M. M. "On persistence in mutual fund performance." *The Journal of Finance*, 52 (1997), 57-82.
- Carrasco, M. and B. Rossi. "In-sample inference and forecasting in misspecified factor models." *Journal of Business & Economic Statistics*, 34 (2016), 313-338.
- Chen, T. and C. Guestrin. "Xgboost: A Scalable Tree Boosting System." *Proceedings of the 22Nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, (2016), 785-794.
- Chen, Y.; K. S. Rogoff and B. Rossi. "Can Exchange Rates Forecast Commodity Prices?" *The Quarterly Journal of Economics*, 125 (2010), 1145-1194.
- Chun, H. and S. Keleş. "Sparse Partial Least Squares Regression for Simultaneous Dimension Reduction and Variable Selection." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72 (2010), 3-25.
- Cochrane, J. H. and M. Piazzesi. "Bond risk premia." *American Economic Review*, 95 (2005), 138-160.
- Comon, P. "Independent Component Analysis, a New Concept?" *Signal Processing*, 36 (1994), 287-314.
- Cybenko, G. "Approximation by Superpositions of a Sigmoidal Function." *Mathematics of Control, Signals and Systems*, 2 (1989), 303-314.
- D'Hondt, C.; R. De Winne; E. Ghysels and S. Raymond. "Artificial Intelligence Alter Egos: Who might benefit from robo-investing?" *Journal of Empirical Finance*, 59 (2020), 278-299.
- De Jong, S. "SIMPLS: An Alternative Approach to Partial Least Squares Regression." *Chemometrics and Intelligent Laboratory Systems*, 18 (1993), 251-263.

- De Mol, C.; D. Giannone and L. Reichlin. "Forecasting using a Large Number of Predictors: Is Bayesian Shrinkage a Valid Alternative to Principal Components?" *Journal of Econometrics*, 146 (2008), 318-328.
- DeMiguel, V.; L. Garlappi and R. Uppal. "Optimal Versus Naive Diversification: How Inefficient is the 1/N Portfolio Strategy?" *The Review of Financial Studies*, 22 (2009), 1915-1953.
- Dietterich, T. G. "Ensemble Methods in Machine Learning." *International Workshop on Multiple Classifier Systems*, (2000), 1-15.
- Duchin, R. and H. Levy. "Markowitz Versus the Talmudic Portfolio Diversification Strategies." *Journal of Portfolio Management*, 35 (2009), 71.
- Engle, R. "Dynamic Conditional Correlation: A Simple Class of Multivariate Generalized Autoregressive Conditional Heteroskedasticity Models." *Journal of Business & Economic Statistics*, 20 (2002), 339-350.
- Fama, E. F. and K. R. French. "Common risk factors in the returns on stocks and bonds." *Journal of Financial Economics*, 33 (1993), 3-56.
- Fama, E. F. and K. R. French. "Dissecting anomalies with a five-factor model." *The Review of Financial Studies*, 29 (2016), 69-103.
- Fan, J. and R. Li. "Variable Selection Via Nonconcave Penalized Likelihood and its Oracle Properties." *Journal of the American Statistical Association*, 96 (2001), 1348-1360.
- Frank, L. E. and J. H. Friedman. "A Statistical View of some Chemometrics Regression Tools." *Technometrics*, 35 (1993), 109-135.
- Freund, Y. "Boosting a Weak Learning Algorithm by Majority." *Information and Computation*, 121 (1995), 256-285.
- Friedman, J. H. "Fast Sparse Regression and Classification." *International Journal of Forecasting*, 28 (2012), 722-738.
- Friedman, J. H. "Greedy Function Approximation: A Gradient Boosting Machine." *Annals of Statistics*, (2001), 1189-1232.
- Friedman, J.; T. Hastie and R. Tibshirani. "Sparse Inverse Covariance Estimation with the Graphical Lasso." *Biostatistics*, 9 (2008), 432-441.
- Gao, X. and F. Nardari. "Do Commodities Add Economic Value in Asset Allocation? New Evidence from Time-Varying Moments." *Journal of Financial and Quantitative Analysis*, 53 (2018), 365-393.
- Gargano, A. and A. Timmermann. "Forecasting Commodity Price Indexes using Macroeconomic and Financial Predictors." *International Journal of Forecasting*, 30 (2014), 825-843.
- Gargano, A.; D. Pettenuzzo and A. Timmermann. "Bond Return Predictability: Economic Value and Links to the Macroeconomy." *Management Science*, 65 (2017), 508-540.
- Geurts, P.; D. Ernst and L. Wehenkel. "Extremely Randomized Trees." *Machine Learning*, 63 (2006), 3-42.
- Glorot, X.; A. Bordes and Y. Bengio. "Deep Sparse Rectifier Neural Networks." *Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics*, (2011), 315-323.
- Goodfellow, I.; Y. Bengio and A. Courville. "Deep Learning." *MIT Press*, (2016).
- Gu, S.; B. T. Kelly and D. Xiu. "Empirical Asset Pricing Via Machine Learning" *The Review of Financial Studies*, 33 (2020), 2223-2273.

Hansen, L. K. and P. Salamon. "Neural Network Ensembles." *IEEE Transactions on Pattern Analysis & Machine Intelligence*, (1990), 993-1001.

Hastie, T.; R. Tibshirani and J. Friedman. "The Elements of Statistical Learning: Prediction, Inference and Data Mining." *Springer-Verlag, New York*, (2009).

Hoerl, A. E. and R. W. Kennard. "Ridge Regression: Biased Estimation for Nonorthogonal Problems." *Technometrics*, 12 (1970), 55-67.

Hornik, K.; M. Stinchcombe and H. White. "Multilayer Feedforward Networks are Universal Approximators." *Neural Networks*, 2 (1989), 359-366.

Hyvärinen, A. and E. Oja. "Independent Component Analysis: Algorithms and Applications." *Neural Networks*, 13 (2000), 411-430.

Ioffe, S. and C. Szegedy. "Batch Normalization: Accelerating Deep Network Training by Reducing Internal Covariate Shift." *Proceedings of the 32nd International Conference on Machine Learning*, 37 (2015), 448-456.

Jagannathan, R. and T. Ma. "Risk reduction in large portfolios: Why imposing the wrong constraints helps." *The Journal of Finance*, 58 (2003), 1651-1683.

Jaggi, M. "An Equivalence between the Lasso and Support Vector Machines." *Chapman and Hall/CRC*, (2014).

Jurado, K.; S. C. Ludvigson and S. Ng. "Measuring Uncertainty." *American Economic Review*, 105 (2015), 1177-1216.

Kelly, B. and S. Pruitt. "The Three-Pass Regression Filter: A New Approach to Forecasting using Many Predictors." *Journal of Econometrics*, 186 (2015), 294-316.

Kelly, B. T.; S. Pruitt and Y. Su. "Characteristics are Covariances: A Unified Model of Risk and Return." *Journal of Financial Economics*, 134 (2019), 501-524.

Kilian, L. "Not all Oil Price Shocks are Alike: Disentangling Demand and Supply Shocks in the Crude Oil Market." *American Economic Review*, 99 (2009), 1053-1069.

Kim, H. H. and N. R. Swanson. "Forecasting Financial and Macroeconomic Variables using Data Reduction Methods: New Empirical Evidence." *Journal of Econometrics*, 178 (2014), 352-367.

Kingma, D. P. and J. Ba. "Adam: A Method for Stochastic Optimization." *International Conference on Learning Representations*, (2015).

Kirby, C. and B. Ostdiek. "It's all in the Timing: Simple Active Portfolio Strategies that Outperform Naive Diversification." *Journal of Financial and Quantitative Analysis*, 47 (2012), 437-467.

Kotchoni, R.; M. Leroux and D. Stevanovic. "Macroeconomic Forecast Accuracy in a data-rich Environment." *Journal of Applied Econometrics*, 34 (2019), 1050-1072.

Kozak, S. "Kernel trick for the cross-section.", (2019), Available at SSRN: <https://ssrn.com/abstract=3307895>.

Kritzman, M.; S. Page and D. Turkington. "In Defense of Optimization: The Fallacy of 1/N." *Financial Analysts Journal*, 66 (2010), 31-39.

Le, Q. V.; A. Karpenko; J. Ngiam and A. Y. Ng. "ICA with Reconstruction Cost for Efficient Overcomplete Feature Learning." *Advances in Neural Information Processing Systems*, (2011), 1017-1025.

Ledoit, O. and M. Wolf. "A Well-Conditioned Estimator for Large-Dimensional Covariance Matrices." *Journal of Multivariate Analysis*, 88 (2004), 365-411.

- Lima, L. R. and F. Meng. "Out-of-Sample Return Predictability: A Quantile Combination Approach." *Journal of Applied Econometrics*, 32 (2017), 877-895.
- Lin, H.; C. Wu and G. Zhou. "Forecasting Corporate Bond Returns with a Large Set of Predictors: An Iterated Combination Approach." *Management Science*, 64 (2017), 4218-4238.
- Ludvigson, S. C. and S. Ng. "Macro Factors in Bond Risk Premia." *The Review of Financial Studies*, 22 (2009), 5027-5067.
- Ludvigson, S. C.; S. Ma and S. Ng. "Uncertainty and business cycles: exogenous impulse or endogenous response?" NBER Working Paper No. w21803, (2015). Available at SSRN: <https://ssrn.com/abstract=2703208>
- Lv, J. and Y. Fan. "A Unified Approach to Model Selection and Sparse Recovery using Regularized Least Squares." *The Annals of Statistics*, 37 (2009), 3498-3528.
- Markowitz, H. M. "Portfolio Selection." *The Journal of Finance*, 7 (1952), 77-91.
- Masters, T. "Practical neural network recipes in C." *Morgan Kaufmann*, (1993).
- Nair, V. and G. E. Hinton. "Rectified Linear Units Improve Restricted Boltzmann Machines." *Proceedings of the 27th International Conference on Machine Learning*, (2010), 807-814.
- Neely, C. J.; D. E. Rapach; J. Tu and G. Zhou. "Forecasting the Equity Risk Premium: The Role of Technical Indicators." *Management Science*, 60 (2014), 1772-1791.
- Olivares-Nadal, A. V. and V. DeMiguel. "A Robust Perspective on Transaction Costs in Portfolio Optimization." *Operations Research*, 66 (2018), 733-739.
- Pástor, L. and R. F. Stambaugh. "Liquidity risk and expected stock returns." *Journal of Political Economy*, 111 (2003), 642-685.
- Politis, D. N. and J. P. Romano. "A general resampling scheme for triangular arrays of α -mixing random variables with application to the problem of spectral density estimation." *The Annals of Statistics*, (1992), 1985-2007.
- Rapach, D. E.; J. K. Strauss and G. Zhou. "International Stock Return Predictability: What is the Role of the United States?" *The Journal of Finance*, 68 (2013), 1633-1662.
- Rapach, D. E.; J. K. Strauss and G. Zhou. "Out-of-Sample Equity Premium Prediction: Combination Forecasts and Links to the Real Economy." *The Review of Financial Studies*, 23 (2010), 821-862.
- Rapach, D. E.; J. K. Strauss; J. Tu and G. Zhou. "Industry Return Predictability: A Machine Learning Approach." *The Journal of Financial Data Science*, 1 (2019), 9-28.
- Rapach, D. E.; M. E. Wohar and J. Rangvid. "Macro Variables and International Stock Return Predictability." *International Journal of Forecasting*, 21 (2005), 137-166.
- Schapire, R. E. "The Strength of Weak Learnability." *Machine Learning*, 5 (1990), 197-227.
- Schölkopf, B.; A. J. Smola; R. C. Williamson and P. L. Bartlett. "New Support Vector Algorithms." *Neural Computation*, 12 (2000), 1207-1245.
- Schölkopf, B.; P. L. Bartlett; A. J. Smola and R. C. Williamson. "Shrinking the Tube: A New Support Vector Regression Algorithm." *Advances in Neural Information Processing Systems*, (1999), 330-336.
- Stock, J. H. and M. W. Watson. "Combination Forecasts of Output Growth in a seven-country Data Set." *Journal of Forecasting*, 23 (2004), 405-430.
- Stock, J. H. and M. W. Watson. "Generalized Shrinkage Methods for Forecasting using Many Predictors." *Journal of Business & Economic Statistics*, 30 (2012), 481-493.

- Tibshirani, R. "Regression Shrinkage and Selection Via the Lasso." *Journal of the Royal Statistical Society: Series B (Methodological)*, 58 (1996), 267-288.
- Timmermann, A. "Forecast Combinations." *Handbook of Economic Forecasting*, 1 (2006), 135-196.
- Vapnik, V. N. "The nature of statistical learning theory" *Wiley, New York*, (1995).
- Welch, I. and A. Goyal. "A Comprehensive Look at the Empirical Performance of Equity Premium Prediction." *The Review of Financial Studies*, 21 (2008), 1455-1508.
- Wold, H. "Estimation of Principal Components and Related Models by Iterative Least Squares." *Multivariate Analysis*, (1966), 391-420.
- Zhang, C. "Nearly Unbiased Variable Selection Under Minimax Concave Penalty." *The Annals of Statistics*, 38 (2010), 894-942.
- Zhou, Q.; W. Chen; S. Song; J. Gardner, K. Weinberger and Y. Chen. "A reduction of the elastic net to support vector machines with an application to GPU computing." *Proceedings of the AAAI Conference on Artificial Intelligence*, 29 (2015).
- Zou, H. "The Adaptive Lasso and its Oracle Properties." *Journal of the American Statistical Association*, 101 (2006), 1418-1429.
- Zou, H. and T. Hastie. "Regularization and Variable Selection Via the Elastic Net." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67 (2005), 301-320.
- Zou, H.; T. Hastie and R. Tibshirani. "Sparse Principal Component Analysis." *Journal of Computational and Graphical Statistics*, 15 (2006), 265-286.

Tables

TABLE 1 List of Models

EW	Equally weighted portfolio
HA	Historical average
KS	Kitchen sink model (OLS)
Forecast Combinations of Bivariate Prediction Models	
MC	Equally-weighted average of forecasts (Mean Combination)
MDC	Median Combination forecast
TMC	Trimmed Mean Combination forecast
Rank	Weighted average of forecasts based on MSE ranking
CL2	Average of forecasts contained in the first of two clusters
CL3	Average of forecasts contained in the first of three clusters
DMSFE1	Discounted mean square forecast error, $\psi=1$
DMSFE09	Discounted mean square forecast error, $\psi=0.9$
Shrinkage Methods	
Ridge	Ridge regression
Lasso	Least absolute shrinkage and selection operator
EN	Elastic net
aLasso	Adaptive least absolute shrinkage and selection operator
Bridge	Bridge regression
SCAD	Smoothly clipped absolute deviation
MCP	Minimax concave penalty
SICA	Smooth integration of counting and absolute deviation
Dimensionality Reduction Methods	
PCA	Principal component analysis
SPCA	Sparse principal component analysis
PLS	Partial least squares
SPLS	Sparse partial least squares
ICA	Independent component analysis
RICA	Reconstruction independent component analysis
Non-Linear Machine Learning Methods	
RF	Random forests
ERT	Extremely randomized trees
GBM	Gradient boosting machine
RGBM	Regularized gradient boosting machine
e-SVM	Epsilon support vector machine
nu-SVM	Nu support vector machine
MLP1	Multilayer Perceptron, one hidden layer
MLP2	Multilayer Perceptron, two hidden layers
MLP3	Multilayer Perceptron, three hidden layers
Forecast Combinations of Machine Learning Models	
MC _{ML}	Equally-weighted average of ML forecasts (Mean Combination)
MDC _{ML}	Median Combination of ML forecasts
TMC _{ML}	Trimmed Mean Combination of ML forecasts
Rank _{ML}	Weighted average of ML forecasts based on MSE ranking
CL2 _{ML}	Average of ML forecasts contained in the first of two clusters
CL3 _{ML}	Average of ML forecasts contained in the first of three clusters
DMSFE1 _{ML}	Discounted mean square forecast error, $\psi=1$, of ML forecasts
DMSFE09 _{ML}	Discounted mean square forecast error, $\psi=0.9$, of ML forecasts

TABLE 2 Portfolio Performance Based on Average Return

Model	A. Sparse Covariance				B. Dynamic Covariance			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	5.73	5.73	5.73	5.73	5.73	5.73	5.73	5.73
HA	4.00	7.12	6.06	9.16	3.69	6.77	6.52	10.28
KS	24.96	37.38	<u>20.43</u>	27.29	23.74	33.74	<u>18.84</u>	25.79
Forecast Combinations of Bivariate Prediction Models								
MC	5.29	8.30	6.65	10.22	5.83	8.79	6.88	10.58
MDC	5.53	9.05	6.62	9.81	5.15	8.48	7.06	10.65
TMC	5.39	8.28	6.51	9.91	5.58	8.53	6.90	10.54
Rank	8.11	12.56	7.17	10.81	8.32	12.06	7.68	11.86
CL2	4.41	7.05	6.40	9.81	5.32	7.78	6.27	10.24
CL3	4.78	7.33	6.48	10.12	5.68	7.92	5.99	10.16
DMSFE1	5.29	8.30	6.66	10.24	5.82	8.72	6.86	10.60
DMSFE09	5.34	8.30	6.64	10.28	5.82	8.56	6.85	10.71
Shrinkage Methods								
Ridge	25.31	36.55	20.20	27.40	23.89	33.15	17.97	24.76
Lasso	23.49	32.02	18.13	24.35	22.81	31.95	15.37	20.62
EN	23.30	31.59	18.03	24.13	22.65	31.48	15.23	20.20
aLasso	22.43	31.67	18.11	24.79	21.10	29.56	15.15	20.65
Bridge	22.09	30.50	18.68	26.23	21.07	29.44	16.31	23.41
SCAD	21.17	29.63	18.92	25.92	20.50	27.84	15.94	23.94
MCP	25.19	35.76	20.21	<u>27.48</u>	24.60	34.26	18.52	<u>26.04</u>
SICA	21.31	30.36	17.27	23.86	19.78	28.09	15.25	20.98
Dimensionality Reduction Methods								
PCA	9.44	14.11	9.43	12.67	9.99	14.66	9.41	12.69
SPCA	9.64	16.14	9.52	12.44	9.31	16.14	10.25	13.11
PLS	24.06	34.96	16.94	21.27	21.86	31.56	14.52	17.79
SPLS	21.14	29.52	13.58	17.53	20.45	29.33	13.80	17.07
ICA	11.23	15.53	7.51	9.95	13.56	18.26	9.16	12.18
RICA	8.68	12.28	8.11	11.68	7.26	9.71	6.62	9.23
Non-Linear Machine Learning Methods								
RF	12.19	17.12	9.64	13.48	11.93	16.54	8.77	11.79
ERT	5.88	9.76	7.60	11.45	6.78	10.09	7.59	11.56
GBM	10.88	16.54	11.50	17.16	11.22	16.72	10.53	15.50
RGBM	9.98	15.33	9.55	13.57	6.68	10.92	6.12	8.43
e-SVM	9.48	13.70	8.77	12.63	6.82	9.78	7.91	11.67
nu-SVM	7.05	11.78	6.86	10.59	7.63	13.03	8.66	12.55
MLP1	24.66	34.66	18.05	24.75	22.52	31.98	16.03	20.87
MLP2	13.12	18.73	10.35	13.51	11.96	17.06	7.60	10.71
MLP3	6.79	10.16	6.83	10.20	7.10	8.72	5.63	8.62
Forecast Combinations of Machine Learning Models								
MC _{ML}	25.17	34.12	16.77	21.11	24.05	32.11	13.98	18.33
MDC _{ML}	24.52	36.15	17.25	20.82	22.64	31.98	15.32	19.23
TMC _{ML}	25.25	34.99	16.87	21.63	24.24	32.68	13.94	18.75
Rank _{ML}	19.60	30.09	16.54	21.37	18.89	28.18	14.50	19.08
CL2 _{ML}	23.45	35.10	19.07	24.84	24.28	36.41	17.46	21.47
CL3 _{ML}	24.01	35.62	18.24	23.88	26.12	<u>38.53</u>	18.17	22.71
DMSFE1 _{ML}	25.49	35.66	17.07	21.55	24.33	33.70	14.41	18.61
DMSFE09 _{ML}	<u>27.08</u>	<u>38.93</u>	17.93	23.21	<u>26.60</u>	37.28	15.49	19.23

Notes: This table reports the annualized average return of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. Panel A reports the performance based on a sparse covariance estimator estimated using the graphical lasso, while the results of Panel B are based on a dynamic covariance estimated using the DCC GARCH. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0, 1]$) and portfolios with leverage ($w_j \in [0, 1.5]$). The models with the highest average return are underlined.

TABLE 3 Portfolio Performance Based on Sharpe Ratio

Model	A. Sparse Covariance				B. Dynamic Covariance			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	0.78	0.78	0.78	0.78	0.78	0.78	0.78	0.78
HA	0.30	0.37	0.59	0.67	0.25	0.34	0.63	0.74
KS	1.05	1.16	<u>1.25</u>	1.25	0.96	1.00	1.14	1.16
Forecast Combinations of Bivariate Prediction Models								
MC	0.37	0.42	0.65	0.76	0.39	0.42	0.65	0.75
MDC	0.40	0.47	0.64	0.72	0.35	0.41	0.69	0.77
TMC	0.38	0.42	0.63	0.74	0.37	0.41	0.66	0.75
Rank	0.46	0.54	0.68	0.79	0.44	0.49	0.69	0.81
CL2	0.30	0.35	0.62	0.72	0.35	0.38	0.60	0.73
CL3	0.31	0.35	0.63	0.75	0.37	0.39	0.57	0.72
DMSFE1	0.37	0.42	0.65	0.76	0.39	0.42	0.65	0.75
DMSFE09	0.37	0.42	0.64	0.76	0.39	0.41	0.65	0.76
Shrinkage Methods								
Ridge	1.07	1.14	<u>1.25</u>	<u>1.27</u>	0.96	0.99	1.11	1.16
Lasso	0.96	0.95	1.10	1.12	0.86	0.89	0.90	0.93
EN	0.94	0.93	1.09	1.12	0.85	0.87	0.89	0.91
aLasso	0.91	0.93	1.09	1.12	0.80	0.83	0.89	0.93
Bridge	0.90	0.91	1.17	1.21	0.80	0.84	0.99	1.07
SCAD	0.86	0.89	1.19	1.19	0.79	0.81	0.98	1.10
MCP	1.05	1.10	<u>1.25</u>	<u>1.27</u>	0.97	1.01	1.12	<u>1.19</u>
SICA	0.86	0.90	1.09	1.12	0.73	0.78	0.92	0.97
Dimensionality Reduction Methods								
PCA	0.48	0.54	0.78	0.79	0.50	0.56	0.75	0.76
SPCA	0.39	0.50	0.76	0.80	0.39	0.50	0.82	0.81
PLS	1.08	1.12	1.05	0.98	0.95	1.00	0.87	0.79
SPLS	0.92	0.92	0.84	0.81	0.85	0.86	0.83	0.78
ICA	0.48	0.48	0.56	0.56	0.58	0.56	0.63	0.64
RICA	0.41	0.44	0.62	0.67	0.35	0.35	0.49	0.51
Non-Linear Machine Learning Methods								
RF	0.59	0.61	0.72	0.76	0.57	0.59	0.65	0.64
ERT	0.37	0.46	0.74	0.83	0.40	0.45	0.70	0.79
GBM	0.45	0.48	0.65	0.74	0.46	0.47	0.59	0.66
RGBM	0.43	0.47	0.53	0.57	0.29	0.33	0.37	0.38
e-SVM	0.55	0.57	0.79	0.90	0.39	0.40	0.68	0.76
nu-SVM	0.44	0.53	0.62	0.74	0.46	0.57	0.76	0.83
MLP1	1.01	1.02	1.09	1.14	0.91	0.90	0.90	0.87
MLP2	0.59	0.61	0.77	0.79	0.53	0.58	0.55	0.60
MLP3	0.36	0.40	0.59	0.68	0.36	0.33	0.44	0.52
Forecast Combinations of Machine Learning Models								
MC _{ML}	1.16	1.16	1.16	1.13	1.07	1.04	0.94	0.94
MDC _{ML}	1.04	1.11	1.09	1.00	0.95	0.96	0.96	0.92
TMC _{ML}	1.16	1.18	1.17	1.16	1.07	1.06	0.95	0.98
Rank _{ML}	0.86	0.95	1.15	1.17	0.84	0.88	1.02	1.06
CL2 _{ML}	1.00	1.04	1.17	1.19	1.07	1.13	<u>1.15</u>	1.11
CL3 _{ML}	0.99	1.04	1.08	1.08	1.13	1.18	<u>1.15</u>	1.14
DMSFE1 _{ML}	1.17	1.19	1.18	1.15	1.06	1.07	0.98	0.97
DMSFE09 _{ML}	<u>1.22</u>	<u>1.28</u>	1.23	1.23	<u>1.18</u>	<u>1.21</u>	1.08	1.03

Notes: This table reports the annualized Sharpe ratio of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. Panel A reports the performance based on a sparse covariance estimator estimated using the graphical lasso, while the results of Panel B are based on a dynamic covariance estimated using the DCC GARCH. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0, 1]$) and portfolios with leverage ($w_j \in [0, 1.5]$). The models with the highest Sharpe ratio are underlined.

TABLE 4 Portfolio Performance Based on Average Turnover

Model	A. Sparse Covariance				B. Dynamic Covariance			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
HA	8.04	9.79	6.55	10.58	16.12	24.82	23.35	39.31
KS	182.42	273.82	177.69	260.47	180.35	270.92	178.03	263.34
Forecast Combinations of Bivariate Prediction Models								
MC	40.67	48.02	24.61	35.98	51.88	67.30	36.82	59.07
MDC	21.04	24.48	12.67	19.02	29.62	40.78	27.19	45.93
TMC	37.17	44.13	22.71	33.30	47.37	61.99	34.49	56.34
Rank	122.36	164.42	82.44	113.65	136.22	191.68	102.25	145.23
CL2	59.60	74.43	38.22	55.29	69.89	92.74	48.42	79.69
CL3	78.89	103.17	52.67	74.59	90.25	122.22	63.01	98.70
DMSFE1	40.83	48.31	24.77	36.18	52.05	67.56	36.92	59.20
DMSFE09	40.73	48.51	24.73	36.19	52.09	67.83	36.71	58.71
Shrinkage Methods								
Ridge	182.75	274.05	177.30	258.44	180.47	270.85	178.39	263.68
Lasso	177.26	267.33	170.80	247.97	174.01	262.86	175.47	259.47
EN	178.04	268.31	171.49	248.62	175.21	264.24	175.77	259.54
aLasso	177.83	267.44	170.35	246.90	175.67	263.87	175.10	258.56
Bridge	182.63	273.42	175.50	257.16	180.47	270.65	177.79	264.61
SCAD	182.63	274.20	174.96	255.89	180.92	270.54	177.20	263.06
MCP	183.03	274.55	177.54	260.03	181.34	271.11	178.87	265.27
SICA	181.75	271.61	174.18	254.07	180.34	269.76	176.32	261.25
Dimensionality Reduction Methods								
PCA	151.17	215.06	118.22	167.30	157.19	230.38	130.80	187.16
SPCA	155.49	224.36	128.32	179.96	163.11	239.16	142.87	205.50
PLS	177.51	266.15	169.86	246.51	174.65	263.06	172.22	253.15
SPLS	177.85	265.59	168.19	243.13	178.80	266.75	174.92	258.38
ICA	152.13	224.07	128.19	179.37	155.72	230.42	138.87	198.97
RICA	153.47	223.24	131.73	188.33	154.96	228.85	140.82	204.02
Non-Linear Machine Learning Methods								
RF	151.46	219.92	123.11	170.48	154.25	229.41	138.38	196.37
ERT	82.87	110.56	57.37	80.06	101.30	138.07	72.43	106.63
GBM	178.06	266.65	172.91	255.86	178.92	268.34	174.85	259.51
RGBM	181.99	271.01	173.65	254.98	182.68	272.69	177.34	260.04
e-SVM	121.57	176.16	105.33	151.77	126.40	189.28	119.60	175.38
nu-SVM	117.27	172.07	96.80	135.37	124.46	184.33	114.53	168.32
MLP1	163.82	247.16	166.66	248.59	160.98	242.38	163.10	244.87
MLP2	177.05	266.60	175.71	259.82	174.24	262.03	172.86	258.34
MLP3	<u>185.95</u>	<u>276.97</u>	<u>181.64</u>	<u>269.58</u>	<u>185.44</u>	<u>276.57</u>	<u>180.35</u>	<u>269.61</u>
Forecast Combinations of Machine Learning Models								
MC _{ML}	176.62	263.17	166.56	241.71	175.56	261.14	169.32	250.21
MDC _{ML}	176.91	263.59	167.65	243.65	177.58	264.40	169.98	250.88
TMC _{ML}	177.22	263.41	165.86	239.91	176.16	262.47	169.94	250.85
Rank _{ML}	168.14	250.13	150.69	211.93	169.14	253.28	158.75	228.07
CL2 _{ML}	166.36	246.79	151.69	216.84	166.34	248.31	158.09	230.43
CL3 _{ML}	165.37	244.75	147.25	208.76	166.25	248.27	156.09	225.24
DMSFE1 _{ML}	175.92	261.89	165.69	239.49	174.48	260.09	168.90	250.31
DMSFE09 _{ML}	175.81	261.93	165.96	239.93	175.35	263.88	171.18	253.19

Notes: This table reports the annualized average turnover of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. Panel A reports the performance based on a sparse covariance estimator estimated using the graphical lasso, while the results of Panel B are based on a dynamic covariance estimated using the DCC GARCH. The performance is reported for different levels of risk aversion ($\gamma = 2,10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$). The models with the highest turnover are underlined.

TABLE 5 Average Weight in Each Asset Class for Portfolios Based on the Sparse Covariance Estimator

Model	Aggressive ($\gamma=2$)						Conservative ($\gamma=10$)					
	[0,1]			[0,1.5]			[0,1]			[0,1.5]		
	S	B	C	S	B	C	S	B	C	S	B	C
EW	43.86	10.53	45.61	43.86	10.53	45.61	43.86	10.53	45.61	43.86	10.53	45.61
HA	97.46	0.00	2.54	142.69	0.00	7.31	90.71	1.29	8.00	120.23	17.09	12.68
KS	50.25	0.00	49.75	75.33	0.00	74.67	48.49	0.00	51.51	70.38	0.00	79.62
Forecast Combinations of Bivariate Prediction Models												
MC	96.21	0.00	3.79	141.59	0.00	8.41	90.31	1.15	8.54	120.68	15.45	13.87
MDC	97.47	0.00	2.53	142.91	0.00	7.09	90.98	1.12	7.90	121.16	16.22	12.62
TMC	96.44	0.00	3.56	141.95	0.00	8.05	90.47	1.15	8.38	120.87	15.51	13.62
Rank	91.72	0.00	8.28	135.55	0.09	14.36	84.40	2.61	12.99	114.70	13.76	21.54
CL2	95.23	0.00	4.77	140.44	0.00	9.56	88.98	1.53	9.49	119.48	15.16	15.36
CL3	94.30	0.00	5.70	139.25	0.06	10.69	87.49	2.07	10.44	118.09	14.77	17.14
DMSFE1	96.19	0.00	3.81	141.57	0.00	8.43	90.28	1.14	8.58	120.68	15.40	13.92
DMSFE09	96.27	0.00	3.73	141.68	0.00	8.32	90.25	1.16	8.58	120.60	15.44	13.96
Shrinkage Methods												
Ridge	51.01	0.00	48.99	76.76	0.00	73.24	49.32	0.00	50.68	71.47	0.00	78.53
Lasso	51.06	0.00	48.94	77.44	0.00	72.56	49.76	0.01	50.22	71.61	0.22	78.17
EN	51.03	0.00	48.97	77.35	0.00	72.65	50.04	0.01	49.95	72.08	0.20	77.72
aLasso	50.82	0.00	49.18	76.91	0.00	73.09	48.93	0.00	51.07	70.38	0.34	79.28
Bridge	51.68	0.00	48.32	77.79	0.00	72.21	49.86	0.00	50.14	72.65	0.07	77.28
SCAD	51.23	0.00	48.77	78.33	0.00	71.67	50.36	0.00	49.64	72.85	0.02	77.13
MCP	51.38	0.00	48.62	76.91	0.00	73.09	49.21	0.00	50.79	71.69	0.00	78.31
SICA	51.31	0.00	48.69	77.43	0.00	72.57	50.31	0.00	49.69	72.22	0.04	77.75
Dimensionality Reduction Methods												
PCA	74.36	2.60	23.04	111.13	4.53	34.34	68.38	5.83	25.79	95.56	13.54	40.91
SPCA	74.02	1.58	24.40	109.68	3.06	37.25	68.01	4.86	27.13	95.23	11.72	43.05
PLS	57.22	0.00	42.78	86.80	0.00	63.20	56.12	0.18	43.70	81.04	0.84	68.12
SPLS	63.61	0.00	36.39	95.33	0.00	54.67	59.49	0.61	39.89	85.28	2.32	62.40
ICA	79.66	0.13	20.20	118.22	0.51	31.27	71.71	2.22	26.07	99.58	7.06	43.37
RICA	73.15	0.67	26.18	108.35	1.34	40.31	66.17	2.33	31.51	92.72	6.52	50.76
Non-Linear Machine Learning Methods												
RF	64.60	0.10	35.30	98.25	0.28	51.48	68.30	0.81	30.89	100.72	4.24	45.05
ERT	92.97	0.01	7.02	138.11	0.29	11.60	86.41	1.28	12.31	118.98	10.85	20.17
GBM	50.09	0.00	49.91	75.63	0.00	74.37	50.34	0.05	49.61	73.41	0.14	76.45
RGBM	59.12	0.00	40.88	87.64	0.00	62.36	55.25	0.08	44.66	80.05	0.35	69.59
e-SVM	93.38	0.17	6.45	139.02	0.68	10.29	84.27	4.65	11.07	111.32	18.61	20.07
nu-SVM	84.91	0.00	15.09	126.30	0.13	23.57	77.05	2.05	20.89	104.28	9.41	36.31
MLP1	53.18	0.00	46.82	81.59	0.00	68.41	55.71	0.10	44.19	80.95	0.33	68.72
MLP2	55.15	0.30	44.55	82.56	0.61	66.82	49.51	2.75	47.74	68.41	6.31	75.29
MLP3	54.16	1.92	43.92	80.54	3.48	65.98	45.63	7.18	47.19	61.65	15.96	72.39
Forecast Combinations of Machine Learning Models												
MC _{ML}	59.53	0.00	40.47	90.03	0.00	59.97	57.63	0.27	42.10	81.44	1.22	67.34
MDC _{ML}	58.40	0.00	41.60	87.90	0.06	62.03	56.32	0.55	43.13	80.78	2.08	67.14
TMC _{ML}	59.51	0.00	40.49	90.46	0.00	59.54	58.15	0.33	41.52	82.29	1.38	66.32
Rank _{ML}	59.41	0.00	40.59	89.33	0.00	60.67	61.39	0.86	37.75	89.78	3.69	56.54
CL2 _{ML}	53.18	0.00	46.82	79.41	0.00	70.59	54.46	1.00	44.54	81.02	3.39	65.59
CL3 _{ML}	52.55	0.00	47.45	79.03	0.13	70.84	53.57	1.51	44.92	80.12	4.71	65.17
DMSFE1 _{ML}	60.17	0.00	39.83	91.50	0.00	58.50	59.12	0.39	40.49	83.70	1.69	64.61
DMSFE09 _{ML}	61.71	0.00	38.29	93.48	0.00	56.52	59.34	0.33	40.33	83.44	1.68	64.87

Notes: This table reports the average weights in each of the three asset classes (Stock, Bond and Commodities), for the out-of-sample period of January 2000 to December 2019. The mean-variance portfolios are based on the graphical lasso estimator. The average weights are reported for different levels of risk aversion ($\gamma = 2,10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$).

TABLE 6 Portfolio Performance Based on Average Return During Business Cycles

Model	A. Expansions				B. Recessions			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	7.42	7.42	7.42	7.42	-8.15	-8.15	-8.15	-8.15
HA	5.87	9.96	8.13	12.11	-11.34	-16.27	-10.98	-15.05
KS	24.24	35.83	<u>19.91</u>	26.58	30.89	50.13	24.68	33.18
Forecast Combinations of Bivariate Prediction Models								
MC	6.18	10.16	8.53	12.77	-2.04	-6.96	-8.88	-10.77
MDC	6.82	11.24	8.50	12.52	-5.04	-9.00	-8.85	-12.52
TMC	6.34	10.31	8.44	12.56	-2.38	-8.41	-9.38	-11.88
Rank	6.67	11.39	8.36	12.69	19.96	22.11	-2.60	-4.67
CL2	5.86	9.64	8.21	12.38	-7.55	-14.20	-8.50	-11.32
CL3	5.75	9.31	8.14	12.53	-3.29	-8.96	-7.21	-9.70
DMSFE1	6.18	10.13	8.54	12.78	-2.06	-6.74	-8.84	-10.64
DMSFE09	6.29	10.15	8.52	12.78	-2.47	-6.89	-8.82	-10.24
Shrinkage Methods								
Ridge	23.12	33.30	19.20	26.64	43.39	63.31	<u>28.40</u>	<u>33.68</u>
Lasso	21.91	29.73	17.70	24.65	36.47	50.86	21.66	21.91
EN	21.69	29.16	17.56	24.41	36.58	51.54	21.89	21.81
aLasso	20.66	29.07	17.47	24.47	37.05	53.06	23.37	27.45
Bridge	20.39	28.09	18.26	26.19	36.09	50.34	22.16	26.52
SCAD	19.87	27.44	18.88	26.45	31.87	47.67	19.23	21.58
MCP	23.13	32.49	19.26	<u>26.76</u>	42.11	62.66	28.07	33.45
SICA	18.85	26.99	17.10	24.36	41.53	58.09	18.62	19.69
Dimensionality Reduction Methods								
PCA	9.08	13.65	9.77	13.83	12.36	17.90	6.65	3.14
SPCA	10.26	17.33	10.83	14.68	4.56	6.30	-1.24	-5.94
PLS	24.16	35.50	17.93	23.00	23.19	30.55	8.74	7.00
SPLS	20.12	29.50	14.66	19.29	29.58	29.68	4.74	2.99
ICA	12.52	18.69	10.74	15.04	0.61	-10.41	-19.10	-31.96
RICA	10.26	14.57	9.57	14.16	-4.30	-6.58	-3.89	-8.69
Non-Linear Machine Learning Methods								
RF	13.32	18.77	11.61	16.82	2.92	3.55	-6.52	-14.03
ERT	6.16	11.41	9.83	14.96	3.51	-3.84	-10.73	-17.44
GBM	15.41	23.26	14.43	20.56	-26.45	-38.73	-12.58	-10.82
RGBM	14.37	22.14	13.12	17.86	-26.15	-40.67	-19.78	-21.74
e-SVM	9.97	14.99	9.55	13.82	5.41	3.04	2.40	2.84
nu-SVM	8.36	13.89	8.33	12.58	-3.78	-5.57	-5.17	-5.76
MLP1	23.24	32.72	17.69	24.59	36.32	50.67	21.01	26.07
MLP2	12.29	18.37	11.50	15.43	19.94	21.73	0.82	-2.36
MLP3	8.75	12.86	7.40	10.97	-9.28	-12.03	2.17	3.86
Forecast Combinations of Machine Learning Models								
MC _{ML}	22.32	30.91	16.59	21.77	<u>48.65</u>	60.57	18.25	15.70
MDC _{ML}	22.94	34.37	17.22	21.36	37.57	50.75	17.51	16.40
TMC _{ML}	22.49	31.70	16.58	22.28	48.03	62.04	19.19	16.35
Rank _{ML}	16.83	26.11	15.92	21.77	42.39	62.80	21.67	18.03
CL2 _{ML}	21.50	31.98	18.99	25.63	39.42	60.82	19.75	18.33
CL3 _{ML}	21.40	32.23	18.14	24.79	45.49	63.53	19.04	16.45
DMSFE1 _{ML}	22.74	32.23	16.82	22.22	48.10	<u>63.89</u>	19.14	16.07
DMSFE09 _{ML}	<u>24.91</u>	<u>36.78</u>	17.91	24.03	44.97	56.63	18.04	16.48

Notes: This table reports the annualized average return of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019, during business cycles. The mean-variance portfolios are based on the graphical lasso estimator. Panel A reports the performance during NBER expansions, while Panel B reports the performance during NBER recessions. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$). The models with the highest average return are underlined.

TABLE 7 Portfolio Performance Based on Sharpe Ratio During Business Cycles

Model	A. Expansions				B. Recessions			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	<u>1.25</u>	<u>1.25</u>	1.25	1.25	-0.59	-0.59	-0.59	-0.59
HA	0.45	0.55	0.85	0.97	-0.70	-0.68	-0.74	-0.76
KS	1.09	1.20	1.35	1.34	0.90	1.04	0.94	0.95
Forecast Combinations of Bivariate Prediction Models								
MC	0.46	0.55	0.90	1.03	-0.11	-0.25	-0.59	-0.56
MDC	0.51	0.61	0.88	1.00	-0.28	-0.35	-0.59	-0.64
TMC	0.47	0.56	0.89	1.01	-0.12	-0.30	-0.61	-0.61
Rank	0.41	0.53	0.87	1.02	0.72	0.62	-0.16	-0.22
CL2	0.43	0.52	0.88	0.99	-0.34	-0.46	-0.55	-0.56
CL3	0.42	0.49	0.88	1.01	-0.13	-0.27	-0.45	-0.48
DMSFE1	0.46	0.55	0.90	1.03	-0.11	-0.24	-0.58	-0.55
DMSFE09	0.46	0.55	0.90	1.03	-0.13	-0.25	-0.58	-0.53
Shrinkage Methods								
Ridge	1.04	1.12	1.32	1.37	1.30	1.34	<u>1.08</u>	<u>0.96</u>
Lasso	0.96	0.95	1.21	1.28	1.02	1.01	0.78	0.59
EN	0.94	0.93	1.20	1.27	1.03	1.03	0.79	0.59
aLasso	0.91	0.93	1.19	1.25	1.02	1.03	0.83	0.73
Bridge	0.87	0.90	1.27	1.33	1.08	1.07	0.87	0.78
SCAD	0.85	0.87	1.30	1.34	0.98	1.03	0.77	0.61
MCP	1.03	1.07	1.33	1.38	1.26	1.32	1.06	0.94
SICA	0.82	0.88	1.20	1.27	1.13	1.12	0.72	0.56
Dimensionality Reduction Methods								
PCA	0.50	0.58	0.93	1.02	0.41	0.42	0.31	0.11
SPCA	0.45	0.59	1.02	1.14	0.13	0.12	-0.05	-0.20
PLS	1.17	1.23	1.28	1.25	0.71	0.65	0.31	0.17
SPLS	0.91	0.97	0.99	1.00	0.97	0.68	0.19	0.08
ICA	0.55	0.61	0.92	0.99	0.02	-0.25	-0.88	-1.11
RICA	0.59	0.65	0.95	1.07	-0.11	-0.12	-0.14	-0.23
Non-Linear Machine Learning Methods								
RF	0.70	0.73	0.96	1.06	0.09	0.08	-0.32	-0.52
ERT	0.43	0.59	1.05	1.20	0.13	-0.12	-0.70	-0.88
GBM	0.65	0.69	0.86	0.95	-0.98	-0.99	-0.53	-0.33
RGBM	0.64	0.70	0.76	0.79	-1.00	-1.07	-0.88	-0.70
e-SVM	0.59	0.65	0.88	1.03	0.25	0.10	0.17	0.16
nu-SVM	0.54	0.66	0.81	0.96	-0.19	-0.19	-0.31	-0.26
MLP1	1.06	1.08	1.23	1.31	0.93	0.89	0.73	0.67
MLP2	0.59	0.64	0.95	1.03	0.65	0.51	0.04	-0.08
MLP3	0.47	0.52	0.68	0.79	-0.46	-0.40	0.13	0.17
Forecast Combinations of Machine Learning Models								
MC _{ML}	1.13	1.15	1.32	1.35	<u>1.47</u>	1.32	0.71	0.47
MDC _{ML}	1.03	1.12	1.18	1.16	1.16	1.10	0.71	0.47
TMC _{ML}	1.12	1.17	1.32	1.38	<u>1.47</u>	1.35	0.77	0.49
Rank _{ML}	0.82	0.93	1.26	1.37	1.19	1.22	0.87	0.57
CL2 _{ML}	0.98	1.00	1.24	1.33	1.18	1.29	0.83	0.58
CL3 _{ML}	0.93	1.00	1.12	1.19	1.33	1.34	0.85	0.52
DMSFE1 _{ML}	1.14	1.18	1.33	1.38	1.46	<u>1.36</u>	0.76	0.48
DMSFE09 _{ML}	1.23	1.32	<u>1.40</u>	<u>1.45</u>	1.33	1.20	0.72	0.51

Notes: This table reports the annualized Sharpe ratio of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019, during business cycles. The mean-variance portfolios are based on the graphical lasso estimator. Panel A reports the performance during NBER expansions, while Panel B reports the performance during NBER recessions. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$). The models with the highest Sharpe ratio are underlined.


TABLE 8 Portfolio Performance after Transaction Costs of 50 bps

Model	A. Sharpe Ratio				B. Average Turnover			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	0.78	0.78	0.78	0.78	0.00	0.00	0.00	0.00
HA	0.36	0.42	0.57	0.66	0.21	0.27	0.53	0.75
KS	0.64	0.68	0.67	0.65	172.65	259.98	<u>165.22</u>	<u>236.91</u>
Forecast Combinations of Bivariate Prediction Models								
MC	0.32	0.41	0.60	0.65	0.40	0.62	0.71	1.22
MDC	0.34	0.41	0.57	0.63	0.26	0.32	0.60	0.96
TMC	0.33	0.42	0.60	0.64	0.45	0.60	0.71	1.17
Rank	0.50	0.50	0.65	0.72	13.16	18.72	10.32	13.28
CL2	0.25	0.30	0.57	0.64	1.09	1.87	1.48	2.24
CL3	0.36	0.41	0.59	0.65	2.23	3.85	2.64	3.87
DMSFE1	0.32	0.41	0.60	0.65	0.41	0.59	0.74	1.25
DMSFE09	0.31	0.41	0.60	0.66	0.34	0.49	0.72	1.24
Shrinkage Methods								
Ridge	0.64	0.64	0.66	0.65	<u>174.06</u>	<u>260.69</u>	162.27	234.07
Lasso	0.60	0.56	0.56	0.50	163.57	244.23	151.78	216.24
EN	0.58	0.55	0.56	0.50	163.93	246.26	152.48	217.76
aLasso	0.57	0.54	0.55	0.52	165.68	247.76	152.19	215.10
Bridge	0.50	0.48	0.58	0.57	170.09	253.71	160.23	232.95
SCAD	0.51	0.45	0.62	0.59	171.32	256.09	160.13	230.69
MCP	0.60	0.62	0.65	0.61	173.00	260.04	163.64	236.80
SICA	0.48	0.47	0.52	0.50	167.75	251.42	157.10	225.52
Dimensionality Reduction Methods								
PCA	0.16	0.18	0.45	0.51	64.89	91.64	47.84	64.66
SPCA	0.21	0.30	0.57	0.58	80.08	110.94	54.58	70.95
PLS	0.62	0.65	0.50	0.40	159.27	236.95	145.62	203.96
SPLS	0.54	0.53	0.32	0.29	157.38	233.93	141.72	197.18
ICA	0.30	0.24	0.24	0.29	79.86	111.50	57.27	75.14
RICA	-0.02	0.07	0.29	0.37	92.57	131.23	69.56	94.96
Non-Linear Machine Learning Methods								
RF	0.27	0.28	0.38	0.43	91.11	129.81	63.41	81.11
ERT	0.42	0.45	0.64	0.71	15.30	21.21	11.01	13.94
GBM	0.07	0.06	0.10	0.11	169.70	253.09	162.74	239.16
RGBM	0.02	0.02	0.05	0.04	167.95	250.45	157.67	226.08
e-SVM	0.27	0.27	0.57	0.70	28.85	44.20	28.59	37.44
nu-SVM	0.11	0.19	0.38	0.47	27.61	38.54	21.89	29.15
MLP1	0.62	0.63	0.53	0.51	151.05	229.23	150.86	220.42
MLP2	0.14	0.11	0.05	0.00	160.23	238.90	149.26	215.41
MLP3	-0.15	-0.24	-0.23	-0.25	167.83	246.51	156.72	228.84
Forecast Combinations of Machine Learning Models								
MC _{ML}	0.74	0.73	0.59	0.52	146.90	213.65	124.10	168.97
MDC _{ML}	0.71	0.68	0.60	0.50	150.71	220.99	129.56	174.37
TMC _{ML}	0.75	0.76	0.62	0.56	146.09	214.09	122.09	166.32
Rank _{ML}	0.58	0.68	<u>0.80</u>	<u>0.83</u>	119.23	170.80	85.02	107.90
CL2 _{ML}	0.59	0.62	0.78	0.83	124.73	181.71	94.61	124.17
CL3 _{ML}	0.62	0.66	0.73	0.74	129.24	186.32	95.54	125.94
DMSFE1 _{ML}	0.74	0.77	0.64	0.58	141.85	208.46	118.85	159.95
DMSFE09 _{ML}	<u>0.83</u>	<u>0.82</u>	0.72	0.68	142.91	209.06	117.23	159.25


Notes: This table reports the annualized Sharpe ratio and average turnover of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. The portfolios are based on the graphical lasso estimator and on the penalized mean-variance objective function with transaction costs of 50 bps. Panel A reports the performance based on the Sharpe ratio, Panel B reports the results based on average turnover. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0, 1]$) and portfolios with leverage ($w_j \in [0, 1.5]$). The models with the highest Sharpe ratio and turnover are underlined.

Figures

Aggressive Investor



Conservative Investor



— EW — HA — MC_{ML} — MDC_{ML} — TMC_{ML} — Rank_{ML} — CL2_{ML} — CL3_{ML} — DMSFE1_{ML} — DMSFE09_{ML}

FIGURE 1 Cumulative returns for portfolios without transaction costs. The mean-variance portfolios are based on the graphical lasso estimator, with combination of weight constraints $w_j \in [0,1.5]$, for the aggressive investor ($\gamma = 2$) and $w_j \in [0,1]$, for the conservative investor ($\gamma = 10$). The shaded regions depict the NBER-dated recessions and expansions.




FIGURE 2 Asset selection. Columns correspond to individual models and rows to the 30 assets that contribute most to the portfolio, according to the average weight across the out-of-sample period. The mean-variance portfolios are based on the graphical lasso estimator, with combination of weight constraints $w_j \in [0,1.5]$, for the aggressive investor ($\gamma = 2$) and $w_j \in [0,1]$, for the conservative investor ($\gamma = 10$). A darker colour indicates a greater influence of an asset to the portfolio utilising the return forecasts generated from the respective alternative model.



FIGURE 3 Variable importance. Columns correspond to individual models and rows to the 20 most influential predictors for each asset class. The colour of each model-predictor combination indicates the importance of the respective variable to the model, from most influential variable (blue) to least influential variables (white).




FIGURE 4 Cumulative returns for portfolios with transaction costs of 50 bps. The mean-variance portfolios are based on the graphical lasso estimator and on the penalized mean-variance objective function with transaction costs of 50 bps, with combination of weight constraints $w_j \in [0,1.5]$, for the aggressive investor ($\gamma = 2$) and $w_j \in [0,1]$, for the conservative investor ($\gamma = 10$). The shaded regions depict the NBER-dated recessions and expansions

Appendix

TABLE A1 List of Stocks

Ticker	PERMNO	Company Name
HON	10145	HONEYWELL INTERNATIONAL INC
KO	11308	COCA COLA CO
XOM	11850	EXXON MOBIL CORP
GE	12060	GENERAL ELECTRIC CO
IBM	12490	INTERNATIONAL BUSINESS MACHS COR
PEP	13856	PEPSICO INC
MO	13901	ALTRIA GROUP INC
COP	13928	CONOCOPHILLIPS
CVX	14541	CHEVRON CORP NEW
TXN	15579	TEXAS INSTRUMENTS INC
CVS	17005	C V S HEALTH CORP
KMB	17750	KIMBERLY CLARK CORP
BRK	17778	BERKSHIRE HATHAWAY INC DEL
UTX	17830	UNITED TECHNOLOGIES CORP
PG	18163	PROCTER & GAMBLE CO
SO	18411	SOUTHERN CO
CAT	18542	CATERPILLAR INC
CL	18729	COLGATE PALMOLIVE CO
BMJ	19393	BRISTOL MYERS SQUIBB CO
WBA	19502	WALGREENS BOOTS ALLIANCE INC
BA	19561	BOEING CO
ABT	20482	ABBOTT LABORATORIES
LMT	21178	LOCKHEED MARTIN CORP
PFE	21936	PFIZER INC
EMR	22103	EMERSON ELECTRIC CO
JNJ	22111	JOHNSON & JOHNSON
MMM	22592	3M CO
MRK	22752	MERCK & CO INC NEW
NEE	24205	NEXTERA ENERGY INC
DIS	26403	DISNEY WALT CO
HPQ	27828	H P INC
BAX	27887	BAXTER INTERNATIONAL INC
DUK	27959	DUKE ENERGY CORP NEW
OXY	34833	OCCIDENTAL PETROLEUM CORP
WFC	38703	WELLS FARGO & CO NEW
MCD	43449	MCDONALDS CORP
ADP	44644	AUTOMATIC DATA PROCESSING INC
TYC	45356	TYCO INTERNATIONAL LTD
JPM	47896	JPMORGAN CHASE & CO
UNP	48725	UNION PACIFIC CORP
TGT	49154	TARGET CORP
BK	49656	BANK OF NEW YORK MELLON CORP
LLY	50876	LILLY ELI & CO
WMT	55976	WALMART INC
AXP	59176	AMERICAN EXPRESS CO
INTC	59328	INTEL CORP
MDT	60097	MEDTRONIC INC
PNC	60442	P N C FINANCIAL SERVICES GRP INC
LOW	61399	LOWES COMPANIES INC
USB	66157	U S BANCORP DEL

TABLE A2 List of Bond Portfolios

Name	TREASNOX	Description
BP6	2000028	Bonds with maturities between 1 and 6 months
BP12	2000029	Bonds with maturities between 6 and 12 months
BP18	2000030	Bonds with maturities between 12 and 18 months
BP24	2000031	Bonds with maturities between 18 and 24 months
BP30	2000032	Bonds with maturities between 24 and 30 months
BP36	2000033	Bonds with maturities between 30 and 36 months
BP42	2000034	Bonds with maturities between 36 and 42 months
BP48	2000035	Bonds with maturities between 42 and 48 months
BP54	2000036	Bonds with maturities between 48 and 54 months
BP60	2000037	Bonds with maturities between 54 and 60 months
BP120	2000038	Bonds with maturities between 60 and 120 months
BP120b	2000039	Bonds with maturities greater than 120 months

TABLE A3 List of Commodities

Name	Description
Coal	Coal, Australian thermal coal, 12,000- btu/pound, less than 1% sulfur, 14% ash, FOB Newcastle/Port Kembla, US\$ per metric ton
Oil	Crude Oil (petroleum), West Texas Intermediate 40 API, Midland Texas, US\$ per barrel
Cotton	Cotton, Cotton Outlook 'A Index', Middling 1-3/32 inch staple, CIF Liverpool, US cents per pound
Hides	Hides, Heavy native steers, over 53 pounds, wholesale dealer's price, US, Chicago, fob Shipping Point, US cents per pound
Rubber	Rubber, Singapore Commodity Exchange, No. 3 Rubber Smoked Sheets, 1st contract, US cents per pound
Logsk	Hard Logs, Best quality Malaysian meranti, import price Japan, US\$ per cubic meter
Sawmal	Hard Sawnwood, Dark Red Meranti, select and better quality, C&F UK port, US\$ per cubic meter
Logore	Soft Logs, Average Export price from the US for Douglas Fir, US\$ per cubic meter
Sawore	Soft Sawnwood, average export price of Douglas Fir, US Price, US\$ per cubic meter
WoolC	Wool, coarse, 23 micron, Australian Wool Exchange spot quote, US cents per kilogram
WoolF	Wool, fine, 19 micron, Australian Wool Exchange spot quote, US cents per kilogram
Aluminum	Aluminum, 99.5% minimum purity, LME spot price, CIF UK ports, US\$ per metric ton
Copper	Copper, grade A cathode, LME spot price, CIF European ports, US\$ per metric ton
Iron	China import Iron Ore Fines 62% FE spot (CFR Tianjin port), US dollars per metric ton
Lead	Lead, 99.97% pure, LME spot price, CIF European Ports, US\$ per metric ton
Nickel	Nickel, melting grade, LME spot price, CIF European ports, US\$ per metric ton
Tin	Tin, standard grade, LME spot price, US\$ per metric ton
Uranium	Uranium, NUEXCO, Restricted Price, Nuexco exchange spot, US\$ per pound
Zinc	Zinc, high grade 98% pure, US\$ per metric ton
Gold	Gold, Fixing Committee of the London Bullion Market Association, London 3 PM fixed price, US\$ per troy ounce
Platinum	Platinum, LME spot price, USD/troy ounce
Silver	Silver, London Bullion Market Association, USD/troy ounce
Cocoa	Cocoa beans, International Cocoa Organization cash price, CIF US and European ports, US\$ per metric ton
CoffeOM	Coffee, Other Mild Arabicas, International Coffee Organization New York cash price, ex-dock New York, US cents per pound
CoffeeRob	Coffee, Robusta, International Coffee Organization New York cash price, ex-dock New York, US cents per pound
Tea	Tea, Mombasa, Kenya, Auction Price, US cents per Kg, From July 1998, Kenya auctions, Best Pekoe Fannings. Prior, London auctions, c.i.f. UK warehouses
Barley	Barley, Canadian no.1 Western Barley, spot price, US\$ per metric ton
Maize	Maize (corn), U.S. No.2 Yellow, FOB Gulf of Mexico, US price, US\$ per metric ton
Oats	Generic 1st 'O' Future, USD/bushel
Rice	Rice, 5 percent broken milled white rice, Thailand nominal price quote, US\$ per metric ton
Wheat	Wheat, No.1 Hard Red Winter, ordinary protein, Kansas City, US\$ per metric ton
Beef	Beef, Australian and New Zealand 85% lean fores, CIF US import price, US cents per pound
Lamb	Lamb, frozen carcass Smithfield London, US cents per pound
Poultry	Poultry (chicken), Whole bird spot price, Ready-to-cook, whole, iced, Georgia docks, US cents per pound
Pork	Swine (pork), 51-52% lean Hogs, US price, US cents per pound.
Bananas	Bananas, Central American and Ecuador, FOB US Ports, US\$ per metric ton
Fishmeal	Fishmeal, Peru Fish meal/pellets 65% protein, CIF, US\$ per metric ton
Groundnuts	Groundnuts (peanuts), 40/50 (40 to 50 count per ounce), cif Argentina, US\$ per metric ton
OrangeJuice	Generic 1st 'JO' Future, USD/lb
Salmon	Fish (salmon), Farm Bred Norwegian Salmon, export price, US\$ per kilogram
Shrimp	Thailand Whiteleg Shrimp 70 Shrimps/Kg Spot Price
SugarFM	Sugar, Free Market, Coffee Sugar and Cocoa Exchange (CSCE) contract no.11 nearest future position, US cents per pound
SugarUS	Sugar, US import price, contract no.14 nearest futures position, US cents per pound (Footnote: No. 14 revised to No. 16)
OliveOil	Olive Oil, extra virgin less than 1% free fatty acid, ex-tanker price UK, US\$ per metric ton
PalmOil	Palm oil, Malaysia Palm Oil Futures (first contract forward) 4-5 percent FFA, US\$ per metric ton
RapeseedOil	Rapeseed oil, crude, fob Rotterdam, US\$ per metric ton
SoybeanOil	Soybean Oil, Chicago Soybean Oil Futures (first contract forward) exchange approved grades, US\$ per metric ton
Soybeans	Soybeans, US soybeans, Chicago Soybean futures contract (first contract forward) No. 2 yellow and par, US\$ per metric ton
SunflowerOil	Sunflower oil, Sunflower Oil, US export price from Gulf of Mexico, US\$ per metric ton

DAP	US Gulf NOLA DAP Export Spot Price per MT, USD/metric tonne
Potassium	Potassium Chloride (Muriate of Potash) Standard Grade: FOB Vancouver Spot Price, USD/metric tonne
Urea	US Gulf NOLA Urea Granular Spot Price, USD/ST

Notes: The source of the data and their description is obtained from the IMF Primary Commodity Price System database.

TABLE A4 List of Predictors

Name	Description	Source
MKT	Fama-French Market Factor	Kenneth French's Website
SMB	Fama-French Size Factor	Kenneth French's Website
HML	Fama-French Value Factor	Kenneth French's Website
RMW	Fama-French Profitability Factor	Kenneth French's Website
CMA	Fama-French Investment Factor	Kenneth French's Website
MOM	Carhart Momentum Factor	Kenneth French's Website
LIQ	Pastor-Stambaugh Liquidity Factor	Robert Stambaugh's Website
DP	Dividend-Price Ratio	Amit Goyal's Website
DY	Dividend Yield	Amit Goyal's Website
EP	Earnings-Price Ratio	Amit Goyal's Website
DE	Payout Ratio	Amit Goyal's Website
BM	Book-to-Market Ratio	Amit Goyal's Website
SVAR	Stock Variance	Amit Goyal's Website
NTIS	Net Equity Expansion	Amit Goyal's Website
BAML	ICE BofA US Corporate Total Return Index	St. Louis Fed's FRED Database
TBL	Treasury Bill Rate	Amit Goyal's Website
LTY	Long-term Yield	Amit Goyal's Website
LTR	Long-term Return	Amit Goyal's Website
TMS	Term Spread	Amit Goyal's Website
DFY	Default Yield Spread	Amit Goyal's Website
DFR	Default Return Spread	Amit Goyal's Website
CP	Cochrane-Piazzesi Bond Factor	Center for Research in Security Prices
INDPRO	Industrial Production	St. Louis Fed's FRED Database
M1	Money supply M1	St. Louis Fed's FRED Database
CPI	Consumer Price Index	St. Louis Fed's FRED Database
PPI	Producer Price Index	St. Louis Fed's FRED Database
CAP	Capacity Utilization	St. Louis Fed's FRED Database
UNRAT		
E	Unemployment Rate	St. Louis Fed's FRED Database
IS	Inventory-Sales Ratio	St. Louis Fed's FRED Database
HOUST	Housing Starts	St. Louis Fed's FRED Database
Credit	Total Consumer Credit	St. Louis Fed's FRED Database
GSCI	S&P Goldman Sachs Commodity Total Return Index	Thomson Reuters Eikon
USAU	Australian dollar - US dollar exchange rate	St. Louis Fed's FRED Database
USCA	Canadian dollar - US dollar exchange rate	St. Louis Fed's FRED Database
USIN	Indian rupee - US dollar exchange rate	St. Louis Fed's FRED Database
USNZ	New Zealand dollar - US dollar exchange rate	St. Louis Fed's FRED Database
VXO	CBOE Volatility Index	St. Louis Fed's FRED Database
UMSEN		
T	University of Michigan: Consumer Sentiment Index	St. Louis Fed's FRED Database
CFNAI	Chicago Fed National Activity Index	Chicago Fed's Website
REA	Real Economic Activity Index	Lutz Kilian's Website
GAC	Philadelphia Fed Business Outlook Survey Current Activity Index	Philadelphia Fed's Website
GAF	Philadelphia Fed Business Outlook Survey Future Activity Index	Philadelphia Fed's Website
UFin	Financial Uncertainty Index	Sydney C. Ludvigson's Website
UMacro	Macroeconomic Uncertainty Index	Sydney C. Ludvigson's Website

TABLE A5 Summary Statistics of Predictors

	Mean	St. Dev.	Skewness	Kurtosis	Minimum	Maximum
MKT	0.0103	0.0439	-0.7410	2.3274	-0.2264	0.1289
SMB	0.0009	0.0287	0.4104	4.5328	-0.1491	0.1832
HML	0.0023	0.0294	0.1548	1.9267	-0.1118	0.1287
RMW	0.0036	0.0234	-0.4153	12.3216	-0.1834	0.1333
CMA	0.0026	0.0200	0.3934	1.8686	-0.0686	0.0956
MOM	0.0056	0.0447	-1.3289	10.6047	-0.3439	0.1836
LIQ	0.0000	0.0003	-0.0859	0.8868	-0.0013	0.0012
DP	-0.0022	0.0435	0.7472	2.9340	-0.1214	0.2512
DY	-0.0022	0.0434	0.7505	2.9843	-0.1214	0.2512
EP	-2.9461	0.4374	-0.7005	3.0488	-4.8365	-1.8987
DE	-0.7895	0.3528	2.9668	13.8100	-1.2442	1.3795
BM	-0.0018	0.0238	0.2478	11.5176	-0.1446	0.1634
SVAR	0.0025	0.0051	9.0499	102.9008	0.0002	0.0709
NTIS	0.0034	0.0204	-0.2910	-0.2716	-0.0577	0.0457
BAML	0.0066	0.0178	0.1700	8.0024	-0.0994	0.1131
TBL	-0.0219	0.4571	-2.2107	30.8913	-4.6200	2.6100
LTY	6.4225	3.1197	0.6233	-0.3841	1.6300	14.8200
LTR	0.7625	3.2371	0.3923	2.0110	-11.2400	15.2300
TMS	0.0047	0.4694	0.7653	20.1537	-3.2800	4.2300
DFY	-0.0009	0.1246	0.8870	11.8964	-0.6300	0.9400
DFR	0.0169	1.5418	-0.4974	6.9311	-9.7500	7.3700
CP	0.0587	1.7022	0.2810	1.2937	-6.5855	6.1543
INDPRO	0.0015	0.0067	-1.1533	6.0317	-0.0443	0.0203
M1	0.0049	0.0081	1.1409	7.7167	-0.0345	0.0579
CPI	0.0000	0.0027	-0.2820	2.6973	-0.0122	0.0108
PPI	0.0000	0.0099	-1.0818	6.8567	-0.0575	0.0357
CAP	-0.0145	0.5198	-0.9985	4.9776	-3.4255	1.4692
UNRATE	-0.0052	0.1686	0.2929	1.2951	-0.7000	0.6000
IS	-0.0002	0.0173	0.6280	5.1817	-0.0620	0.1136
HOUST	0.1854	105.7384	-0.2583	2.3863	-597.0000	363.0000
Credit	0.0052	0.0053	0.6956	12.7641	-0.0323	0.0446
GSCI	0.0028	0.0561	-0.5964	2.9825	-0.3313	0.2065
USAU	-0.0010	0.0259	-0.9516	4.3938	-0.1732	0.0713
USCA	-0.0002	0.0154	-0.4985	6.2952	-0.1129	0.0601
USIN	-0.0045	0.0178	-3.0368	28.4617	-0.1956	0.0594
USNZ	-0.0008	0.0272	-0.4649	2.4203	-0.1433	0.0810
VXO	19.9162	7.7392	1.8688	6.2694	8.0164	67.1510
UMSENT	0.0798	3.8838	-0.0686	1.2159	-12.7000	17.3000
CFNAI	-0.0388	0.5673	-1.2617	4.1197	-2.9100	1.5700
REA	-0.0931	15.2902	-0.7146	6.6443	-100.1851	69.4538
GAC	7.8512	16.9927	-0.8063	1.2231	-57.1000	49.5000
GAF	0.1333	8.8601	-0.0634	0.7618	-31.3000	37.7000
UFin	0.0001	0.0342	0.3709	6.4373	-0.1943	0.2115
UMacro	0.6508	0.0985	1.9164	3.8795	0.5389	1.0764

Notes: All predictors are checked for stationarity using ADF tests and non-stationary predictors are adjusted by taking first differences.

TABLE A6 Portfolio Performance Based on Average Return for Alternative Rebalancing Frequencies

Model	A. 3-month rebalancing				B. 12-month rebalancing			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	5.73	5.73	5.73	5.73	5.73	5.73	5.73	5.73
HA	4.78	8.24	7.01	9.49	5.08	8.37	7.06	9.61
KS	7.44	11.61	7.83	11.28	10.04	14.34	6.59	8.13
Forecast Combinations of Bivariate Prediction Models								
MC	5.00	8.67	6.93	9.59	4.89	7.94	7.46	10.39
MDC	6.56	10.17	7.53	10.11	5.28	8.25	7.00	9.86
TMC	5.06	8.62	7.11	9.73	5.18	8.19	7.34	10.45
Rank	6.10	8.44	5.91	8.40	5.07	8.48	6.63	10.13
CL2	4.55	8.04	6.40	9.24	4.47	7.19	7.08	10.09
CL3	5.21	8.23	5.76	8.40	4.13	6.69	7.16	10.46
DMSFE1	4.94	8.62	6.93	9.60	4.87	7.90	7.44	10.35
DMSFE09	4.92	8.57	6.92	9.58	4.83	7.85	7.46	10.38
Shrinkage Methods								
Ridge	3.84	5.68	5.87	9.06	9.89	13.81	6.61	10.09
Lasso	4.37	5.73	5.06	8.05	11.94	16.64	7.88	10.96
EN	4.11	5.26	5.00	8.04	11.84	16.57	7.83	10.78
aLasso	3.92	5.31	3.41	5.88	11.82	16.68	7.56	10.76
Bridge	4.54	5.90	5.01	7.88	10.71	15.18	6.33	8.73
SCAD	3.09	3.96	4.54	7.54	9.89	14.29	6.18	8.78
MCP	2.98	4.07	6.06	9.34	11.07	15.43	6.71	9.37
SICA	2.07	3.12	4.02	6.61	10.40	15.26	7.13	10.25
Dimensionality Reduction Methods								
PCA	5.19	7.92	6.05	8.76	9.88	14.23	9.44	13.09
SPCA	3.01	6.41	5.45	7.79	<u>13.06</u>	<u>19.34</u>	10.64	14.57
PLS	5.61	9.10	6.31	9.78	4.22	7.24	4.39	6.24
SPLS	1.80	4.33	4.19	6.21	6.51	8.69	3.46	5.27
ICA	4.65	6.64	4.65	7.05	10.78	17.83	9.51	12.97
RICA	4.10	7.02	6.06	9.15	10.30	13.92	7.18	9.12
Non-Linear Machine Learning Methods								
RF	4.92	6.57	4.46	6.63	3.30	5.20	5.47	8.57
ERT	4.36	6.57	5.78	8.42	6.79	10.30	7.75	10.83
GBM	2.97	4.99	4.48	7.75	7.49	10.02	6.35	10.75
RGBM	-0.11	-1.94	1.36	3.98	6.94	10.99	6.33	9.60
e-SVM	3.96	7.47	6.36	8.63	6.70	11.42	7.54	9.76
nu-SVM	0.49	2.89	4.36	6.81	9.36	14.05	9.26	12.58
MLP1	<u>12.43</u>	<u>17.39</u>	<u>9.83</u>	<u>13.00</u>	7.54	11.19	6.90	9.74
MLP2	7.46	9.33	2.09	2.04	7.41	13.70	<u>12.14</u>	<u>18.14</u>
MLP3	4.59	4.94	3.38	5.54	5.92	9.57	5.96	9.30
Forecast Combinations of Machine Learning Models								
MC _{ML}	1.72	3.16	3.24	5.77	8.40	10.80	5.61	9.03
MDC _{ML}	2.24	4.00	3.91	6.96	5.80	8.25	4.75	8.73
TMC _{ML}	1.87	3.18	3.28	6.44	8.81	11.78	5.69	8.68
Rank _{ML}	5.25	7.36	3.54	5.14	9.56	12.27	6.87	11.81
CL2 _{ML}	4.07	5.24	3.68	6.00	7.58	9.86	5.61	8.93
CL3 _{ML}	1.19	2.25	1.10	2.37	7.90	11.27	7.91	10.86
DMSFE1 _{ML}	2.01	3.09	3.34	6.78	7.12	9.12	5.00	8.37
DMSFE09 _{ML}	0.53	1.78	2.89	5.49	6.11	7.98	4.88	8.83

Notes: This table reports the annualized average return of the mean-variance portfolios, with quarterly (Panel A) and annual (Panel B) rebalancing, for the out-of-sample period of January 2000 to December 2019. The portfolios are based on the graphical lasso estimator. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0, 1]$) and portfolios with leverage ($w_j \in [0, 1.5]$). The models with the highest average return are underlined.

TABLE A7 Portfolio Performance Based on Sharpe Ratio for Alternative Rebalancing Frequencies

Model	A. 3-month rebalancing				B. 12-month rebalancing			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>
HA	0.33	0.40	0.63	0.73	0.33	0.39	0.61	0.71
KS	0.29	0.31	0.39	0.42	0.41	0.40	0.34	0.30
Forecast Combinations of Bivariate Prediction Models								
MC	0.34	0.41	0.62	0.72	0.30	0.36	0.67	0.76
MDC	0.45	0.49	0.68	0.78	0.35	0.39	0.62	0.74
TMC	0.34	0.41	0.63	0.74	0.32	0.37	0.66	0.77
Rank	0.35	0.34	0.48	0.54	0.26	0.32	0.54	0.62
CL2	0.29	0.37	0.56	0.68	0.26	0.31	0.63	0.71
CL3	0.32	0.36	0.49	0.59	0.23	0.28	0.64	0.72
DMSFE1	0.33	0.41	0.62	0.72	0.30	0.36	0.67	0.76
DMSFE09	0.33	0.41	0.62	0.72	0.30	0.35	0.67	0.76
Shrinkage Methods								
Ridge	0.15	0.16	0.31	0.35	0.43	0.41	0.35	0.38
Lasso	0.18	0.17	0.27	0.32	0.52	0.50	0.43	0.43
EN	0.17	0.15	0.27	0.32	0.51	0.50	0.43	0.42
aLasso	0.16	0.15	0.19	0.23	0.51	0.51	0.43	0.43
Bridge	0.18	0.17	0.27	0.31	0.41	0.41	0.34	0.33
SCAD	0.12	0.11	0.24	0.29	0.39	0.40	0.33	0.33
MCP	0.12	0.11	0.32	0.36	0.46	0.45	0.35	0.35
SICA	0.09	0.09	0.22	0.26	0.51	0.51	0.41	0.41
Dimensionality Reduction Methods								
PCA	0.27	0.30	0.43	0.47	0.47	0.49	0.68	0.76
SPCA	0.11	0.17	0.35	0.38	0.50	0.54	0.75	0.81
PLS	0.24	0.28	0.37	0.42	0.17	0.20	0.23	0.24
SPLS	0.07	0.13	0.24	0.26	0.28	0.27	0.19	0.21
ICA	0.18	0.18	0.32	0.38	0.39	0.46	0.60	0.62
RICA	0.21	0.26	0.41	0.45	0.50	0.50	0.49	0.45
Non-Linear Machine Learning Methods								
RF	0.23	0.23	0.30	0.34	0.15	0.16	0.35	0.40
ERT	0.27	0.28	0.48	0.55	0.40	0.43	0.61	0.65
GBM	0.12	0.15	0.22	0.26	0.31	0.29	0.31	0.38
RGBM	0.00	-0.04	0.05	0.10	0.26	0.29	0.29	0.31
e-SVM	0.25	0.33	0.52	0.57	0.39	0.47	0.62	0.70
nu-SVM	0.03	0.13	0.34	0.40	0.53	0.56	0.66	0.70
MLP1	0.55	0.54	0.56	0.55	0.36	0.38	0.40	0.40
MLP2	0.38	0.34	0.14	0.10	0.31	0.40	0.62	0.65
MLP3	0.19	0.14	0.17	0.21	0.29	0.33	0.38	0.43
Forecast Combinations of Machine Learning Models								
MC _{ML}	0.07	0.09	0.19	0.25	0.38	0.35	0.33	0.38
MDC _{ML}	0.08	0.11	0.21	0.26	0.23	0.23	0.24	0.33
TMC _{ML}	0.08	0.09	0.19	0.27	0.40	0.37	0.33	0.37
Rank _{ML}	0.19	0.19	0.20	0.22	0.36	0.34	0.37	0.49
CL2 _{ML}	0.16	0.14	0.21	0.26	0.29	0.26	0.29	0.36
CL3 _{ML}	0.05	0.06	0.06	0.10	0.32	0.31	0.40	0.42
DMSFE1 _{ML}	0.08	0.09	0.20	0.29	0.31	0.28	0.29	0.36
DMSFE09 _{ML}	0.02	0.05	0.17	0.24	0.26	0.24	0.28	0.38

Notes: This table reports the annualized Sharpe ratio of the mean-variance portfolios, with quarterly (Panel A) and annual (Panel B) rebalancing, for the out-of-sample period of January 2000 to December 2019. The portfolios are based on the graphical lasso estimator. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0, 1]$) and portfolios with leverage ($w_j \in [0, 1.5]$). The models with the highest Sharpe ratio are underlined.

TABLE A8 Portfolio Performance Using Time-Varying Risk Aversion Parameter

Model	A. Average Return		B. Sharpe Ratio		C. Average Turnover	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	5.73	5.73	0.78	0.78	0.00	0.00
HA	4.60	7.83	0.36	0.43	9.45	12.79
KS	25.34	36.93	1.16	1.24	182.52	273.09
Forecast Combinations of Bivariate Prediction Models						
MC	5.41	8.81	0.41	0.48	34.55	43.35
MDC	5.74	9.31	0.45	0.51	18.28	23.07
TMC	5.42	8.72	0.42	0.48	32.09	39.51
Rank	8.01	11.35	0.51	0.55	113.31	153.18
CL2	4.96	7.94	0.37	0.44	52.95	67.68
CL3	5.07	8.32	0.37	0.46	72.29	94.66
DMSFE1	5.46	8.74	0.42	0.48	34.86	43.88
DMSFE09	5.38	8.76	0.41	0.48	34.77	43.53
Shrinkage Methods						
Ridge	24.11	34.42	1.10	1.15	182.93	273.84
Lasso	21.73	30.21	0.96	0.97	177.89	267.32
EN	21.47	29.66	0.94	0.95	178.65	268.46
aLasso	21.16	29.81	0.93	0.95	178.26	267.13
Bridge	20.50	29.35	0.90	0.96	182.43	272.94
SCAD	20.34	29.69	0.89	0.97	182.99	273.55
MCP	24.22	34.24	1.09	1.14	183.36	274.18
SICA	20.59	28.76	0.91	0.95	180.90	269.38
Dimensionality Reduction Methods						
PCA	9.38	14.72	0.54	0.64	146.10	206.07
SPCA	11.62	17.07	0.53	0.63	151.31	218.31
PLS	23.92	33.75	1.15	1.16	177.56	264.38
SPLS	19.36	27.45	0.90	0.92	177.70	264.22
ICA	10.09	13.76	0.46	0.50	151.54	219.21
RICA	8.70	12.91	0.47	0.53	150.01	216.95
Non-Linear Machine Learning Methods						
RF	12.03	16.82	0.64	0.66	148.90	215.16
ERT	5.98	9.92	0.42	0.52	76.41	104.17
GBM	11.43	18.17	0.49	0.56	177.86	265.31
RGBM	10.34	16.41	0.47	0.52	180.90	268.68
e-SVM	9.07	13.79	0.56	0.62	118.66	170.82
nu-SVM	7.60	11.21	0.50	0.54	116.91	169.96
MLP1	23.57	32.56	1.02	1.03	164.63	247.93
MLP2	12.87	18.22	0.62	0.66	177.39	266.56
MLP3	6.83	11.11	0.39	0.48	185.31	275.74
Forecast Combinations of Machine Learning Models						
MC _{ML}	23.22	32.25	1.16	1.18	175.89	260.59
MDC _{ML}	24.60	34.05	1.12	1.14	176.63	261.41
TMC _{ML}	23.68	32.86	1.18	1.20	176.37	260.50
Rank _{ML}	20.17	30.04	0.95	1.06	167.59	245.56
CL2 _{ML}	23.83	33.65	1.06	1.05	165.30	244.16
CL3 _{ML}	23.48	33.98	1.03	1.07	164.51	241.42
DMSFE1 _{ML}	24.52	33.83	1.21	1.22	175.19	259.66
DMSFE09 _{ML}	26.56	37.04	1.30	1.33	175.34	260.08

Notes: This table reports the annualized average return, Sharpe ratio and average turnover of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. The portfolios are based on the graphical lasso estimator and use the time-varying risk aversion (RA_{bex}) proposed by Bekaert, Engstrom and Xu (2019). The performance is reported for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$). The models with the highest average return, Sharpe ratio and turnover are underlined.

TABLE A9 Portfolio Performance Based on the Shrinkage Covariance Estimator

Model	A. Average Return				B. Sharpe Ratio			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
EW	5.73	5.73	5.73	5.73	0.78	0.78	0.78	0.78
HA	4.04	7.00	6.20	9.46	0.29	0.36	0.58	0.65
KS	25.41	37.50	<u>21.43</u>	28.63	1.02	1.11	<u>1.27</u>	1.25
Forecast Combinations of Bivariate Prediction Models								
MC	5.46	8.60	6.82	10.58	0.37	0.42	0.63	0.73
MDC	5.59	9.21	6.79	10.26	0.39	0.46	0.62	0.70
TMC	5.55	8.55	6.66	10.33	0.38	0.42	0.62	0.71
Rank	8.21	12.88	7.55	11.13	0.44	0.52	0.67	0.76
CL2	4.57	7.59	6.28	9.80	0.29	0.36	0.59	0.67
CL3	5.06	7.83	6.58	10.32	0.31	0.36	0.62	0.71
DMSFE1	5.47	8.56	6.82	10.59	0.37	0.42	0.63	0.73
DMSFE09	5.49	8.57	6.78	10.57	0.37	0.42	0.63	0.73
Shrinkage Methods								
Ridge	25.78	37.65	20.91	29.01	1.03	1.11	1.25	1.29
Lasso	24.20	34.05	18.93	25.85	0.93	0.96	1.09	1.13
EN	24.02	33.65	18.69	25.66	0.92	0.95	1.08	1.12
aLasso	22.97	33.14	18.88	26.14	0.89	0.93	1.08	1.13
Bridge	22.98	31.73	19.23	27.63	0.88	0.90	1.16	1.23
SCAD	21.98	30.92	19.66	27.45	0.85	0.88	1.18	1.22
MCP	25.78	36.88	21.17	<u>29.41</u>	1.02	1.08	1.26	<u>1.30</u>
SICA	22.31	31.68	17.96	24.85	0.84	0.90	1.08	1.11
Dimensionality Reduction Methods								
PCA	9.13	13.72	9.46	13.02	0.43	0.50	0.73	0.76
SPCA	8.77	16.45	9.92	12.97	0.34	0.46	0.73	0.77
PLS	24.00	34.94	18.25	22.85	1.05	1.09	1.10	1.01
SPLS	21.71	30.43	14.53	18.74	0.91	0.92	0.87	0.83
ICA	12.26	17.52	8.48	11.29	0.51	0.52	0.60	0.61
RICA	8.24	12.19	7.92	11.64	0.37	0.41	0.57	0.63
Non-Linear Machine Learning Methods								
RF	12.51	17.62	9.69	13.75	0.58	0.60	0.69	0.74
ERT	6.04	10.06	7.59	11.96	0.36	0.45	0.70	0.81
GBM	11.04	16.45	11.26	16.90	0.45	0.47	0.61	0.70
RGBM	9.92	15.22	10.11	14.19	0.42	0.45	0.54	0.57
e-SVM	9.73	13.89	8.86	12.90	0.55	0.57	0.75	0.85
nu-SVM	6.67	11.13	6.51	10.37	0.41	0.49	0.55	0.66
MLP1	25.43	35.90	18.70	25.72	1.01	1.01	1.08	1.13
MLP2	13.04	19.59	10.82	14.45	0.56	0.61	0.74	0.78
MLP3	7.22	10.02	6.71	9.95	0.37	0.38	0.54	0.62
Forecast Combinations of Machine Learning Models								
MC _{ML}	26.17	35.78	18.11	23.14	1.15	1.15	1.18	1.16
MDC _{ML}	24.83	36.68	18.72	23.47	1.02	1.08	1.13	1.07
TMC _{ML}	26.12	36.60	18.12	23.66	1.15	1.17	1.19	1.19
Rank _{ML}	19.10	29.86	17.17	22.68	0.82	0.91	1.13	1.17
CL2 _{ML}	22.86	34.86	19.71	24.98	0.96	1.01	1.16	1.16
CL3 _{ML}	24.11	35.43	18.94	24.58	0.97	1.00	1.09	1.10
DMSFE1 _{ML}	26.19	36.73	18.47	23.38	1.15	1.17	1.21	1.17
DMSFE09 _{ML}	<u>27.84</u>	<u>39.88</u>	19.25	24.35	<u>1.21</u>	<u>1.25</u>	1.25	1.22

Notes: This table reports the annualized average return and Sharpe ratio of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. The portfolios are based on the Ledoit and Wolf shrinkage estimator of the covariance matrix. The performance is reported for different levels of risk aversion ($\gamma = 2, 10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1.5]$). The models with the highest average return and Sharpe ratio are underlined.

TABLE A10 Portfolio Performance of Shrinkage Methods Using Power Series and Interaction Terms

Model	A. Average Return				B. Sharpe Ratio			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
KS	13.41	19.56	12.16	16.47	0.48	0.47	0.51	0.50
Ridge	14.95	21.62	13.58	20.48	0.57	0.56	0.62	0.67
Lasso	15.40	23.17	15.16	21.18	0.48	0.49	0.59	0.61
EN	16.43	23.77	15.52	23.35	0.54	0.54	0.69	0.76
aLasso	17.11	25.23	15.25	22.30	0.55	0.56	0.61	0.67
Bridge	6.30	9.95	7.02	11.81	0.26	0.28	0.34	0.41
SCAD	10.53	14.46	9.46	14.30	0.41	0.39	0.45	0.49
MCP	6.98	9.53	7.24	11.32	0.27	0.25	0.33	0.37
SICA	13.86	21.26	14.79	22.57	0.49	0.52	0.65	0.71

Notes: This table reports the annualized average return and Sharpe ratio of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. Panel A reports the performance based on the average return, Panel B reports the results based on Sharpe ratio. The performance is reported for different levels of risk aversion ($\gamma = 2,10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1]$). The set of predictors includes the original variables raised to the second and third power and two-way interactions for a total of 1034 predictors.

TABLE A11 Portfolio Performance of Various Machine Learning Methods Using Huber Loss

Model	A. Average Return				B. Sharpe Ratio			
	Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)		Aggressive ($\gamma=2$)		Conservative ($\gamma=10$)	
	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]	[0,1]	[0,1.5]
Ridge _H	16.86	25.77	16.64	22.88	0.67	0.72	0.90	0.90
Lasso _H	19.87	29.12	16.99	23.89	0.85	0.90	1.05	1.11
EN _H	10.35	15.74	11.41	16.10	0.39	0.41	0.60	0.65
GBM _H	9.62	15.96	11.99	17.76	0.41	0.48	0.65	0.72
RGBM _H	9.36	14.71	10.70	14.58	0.35	0.40	0.55	0.57

Notes: This table reports the annualized average return and Sharpe ratio of the mean-variance portfolios, with monthly rebalancing, for the out-of-sample period of January 2000 to December 2019. Panel A reports the performance based on the average return, Panel B reports the results based on Sharpe ratio. The performance is reported for different levels of risk aversion ($\gamma = 2,10$), for unleveraged portfolios ($w_j \in [0,1]$) and portfolios with leverage ($w_j \in [0,1]$).