

# A Framework for Measures of Risk under Uncertainty

Tolulope Fadina\*

Yang Liu<sup>†</sup>Ruodu Wang<sup>‡</sup>

## Abstract

A risk analyst assesses potential financial losses based on multiple sources of information. Often, the assessment does not only depend on the specification of the loss random variable, but also various economic scenarios. Motivated by this observation, we design a unified axiomatic framework for risk evaluation principles which quantifies jointly a loss random variable and a set of plausible probabilities. We call such an evaluation principle a generalized risk measure. We present a series of relevant theoretical results. The worst-case, coherent, and robust generalized risk measures are characterized via different sets of intuitive axioms. We establish the equivalence between a few natural forms of law invariance in our framework, and the technical subtlety therein reveals a sharp contrast between our framework and the traditional one. Moreover, coherence and strong law invariance are derived from a combination of other conditions, which provides additional support for coherent risk measures such as Expected Shortfall over Value-at-Risk, a relevant issue for risk management practice.

**Keywords:** Risk management; model uncertainty; regulatory capital; variational preferences; law invariance; decision theory.

---

\*Department of Mathematical Sciences, University of Essex, United Kingdom. ✉ [t.fadina@essex.ac.uk](mailto:t.fadina@essex.ac.uk).

<sup>†</sup>Department of Statistics and Actuarial Science, University of Waterloo, Canada. ✉ [yang.liu1@uwaterloo.ca](mailto:yang.liu1@uwaterloo.ca).

<sup>‡</sup>Department of Statistics and Actuarial Science, University of Waterloo, Canada. ✉ [wang@uwaterloo.ca](mailto:wang@uwaterloo.ca).

# 1 Introduction

Risk measures are widely used in both financial regulation and economic decisions. Since the seminal work of Artzner et al. (1999), risk measures are commonly defined as functionals on a space of random variables, or, with the assumption of law invariance, on the set of their distribution functions. The most popular risk measures are the Value-at-Risk (VaR) and the Expected Shortfall (ES); see Artzner et al. (1999) and Föllmer and Schied (2016) for the classic theory of risk measures, and documents from the Basel Committee on Banking Supervision (BCBS), e.g., BCBS (2019), for regulatory practice in banking.

In this paper, we propose a novel framework for measures of risk under uncertainty. Let us first explain our motivation. We take market risk as our primary example, although our discussions naturally apply to many other types of risks. A portfolio is associated with a future loss random variable  $X$  representing the portfolio risk. The loss  $X$  has two important practical aspects: the *specification* and the *modeling*.

1. The specification refers to how  $X$  is defined in terms of the underlying risk factors (e.g., asset prices, exchange rates, credit scores, volatilities, etc). More precisely,  $X$  is the financial loss (or gain) from holding assets, derivatives, or other investments in the portfolio. Mathematically, the specification of  $X$  is represented by the function  $X : \Omega \rightarrow \mathbb{R}$ , which maps each state of the future financial world (each element of the sample space  $\Omega$ ) into a realized loss.
2. The modeling refers to the statistical assessment of the likelihood and the severity of the loss  $X$ . The modeling of  $X$  is usually summarized by a distribution, or a collection of distributions in case of model uncertainty, under some estimated or hypothetical (e.g., in stress testing) probability measures  $\mathbb{P} \in \mathcal{P}$ , where  $\mathcal{P}$  is the set of probability measures on the sample space  $\Omega$ .

In the classic framework of Artzner et al. (1999), a risk measure  $\rho$  is defined on a set  $\mathcal{X}$  of random variables, and the risk value  $\rho(X)$  is thereby determined by the specification of  $X$ . The modeling of  $X$  is, however, implicit in this setting: through a given probability  $\mathbb{P}$ , assuming available to the end-user, the distribution of  $X$  under  $\mathbb{P}$  is determined by its specification.

There is a visible gap in the classic setting  $\rho : \mathcal{X} \rightarrow \mathbb{R}$ . In practice, neither  $X$  nor  $\mathbb{P}$  is generally fixed. A change in  $X$  means adjusting positions via trading financial securities. A change in  $\mathbb{P}$  means an update of the modeling, estimation, and calibration of the random world. In financial practice, both  $X$  and  $\mathbb{P}$  evolve on a daily basis for a trading desk; yet they are modified daily for very different reasons.

For another concrete example, suppose that a regulator specifies a risk measure (e.g., ES at level 0.975 as in [BCBS \(2019\)](#)), and two firms assess the risk of the same portfolio separately. Due to different modeling and data processing techniques used by the two firms, their reported ES values are typically not the same. However, the loss random variable  $X$  from the portfolio is the same for both firms. Therefore, the risk measure should not be only determined by the specification of  $X$ , but also the modeling information. In practice, modeling is always subject to uncertainty (called ambiguity in decision theory). Even in the simple estimation of a parametric model, the plausible models are not unique; see [Gilboa and Schmeidler \(1989\)](#) for a classic treatment of ambiguity.

Motivated by the above observations, we propose a new framework of risk measures taking into account both the specification and the modeling of random losses. We choose a set of probability measures instead of a single probability measure as the input for the modeling component. Formally, we introduce the *generalized risk measure*  $\Psi : \mathcal{X} \times 2^{\mathcal{P}} \rightarrow [-\infty, \infty]$ , which has two input arguments: a random variable  $X \in \mathcal{X}$  representing the specification of the loss, and a set of probability measures  $\mathcal{Q} \subseteq \mathcal{P}$  representing the modeling of the random world; each probability measure in  $\mathcal{P}$  is called a *scenario*. Our framework includes the traditional risk measure as a special case when  $\mathcal{Q}$  is a singleton, and when  $\mathcal{Q}$  is fixed but not a singleton, our generalized risk measure can be seen as a scenario-based risk measure of [Wang and Ziegel \(2021\)](#). The framework also incorporates other complicated decision criteria addressing model uncertainty in the literature, which will be discussed later.

We take the perspective of a regulator, who designs a regulatory capital assessment scheme that will be complied with by financial institutions. Financial institutions (or their trading desks) can choose their portfolio positions with losses  $X \in \mathcal{X}$ , and, subject to passing regulatory backtests for statistical prudence, they can also choose their internal models  $\mathcal{Q} \subseteq \mathcal{P}$ . The generalized risk measure is crucial to the design of the capital assessment procedure, because it acts on portfolios and models from the financial institutions, and computes regulatory capital requirement. Therefore, our theoretical framework closely resembles the regulatory practice in the Fundamental Review of the Trading Book (FRTB) of [BCBS \(2019\)](#); see [Wang and Ziegel \(2021\)](#) for discussions on the risk assessment practice of FRTB, and [Cambou and Filipovic \(2017\)](#) for model and scenario aggregation methods in solvency assessment. It is important to note that the input scenario set  $\mathcal{Q}$  does not necessarily contain the decision maker’s subjective probability governing the random world, because most models are simplifications or approximations, as argued by [Cerreia-Vioglio et al. \(2021\)](#).

Figure 1 illustrates a stylized risk assessment procedure, reflecting many of the above considerations. There are four roles: regulator (external), risk analyst (internal), portfolio manager




Figure 1: A stylized procedure for risk assessment practice

(internal), and model risk manager (internal). In Figure 1, except for the regulator’s actions, the other actions are changing dynamically on a daily (or similar) basis, making it clear that one should take both  $\mathcal{Q}$  and  $X$  as inputs and allow them to vary in a unified framework.

## 1.1 Contribution and structure of the paper

As explained above, in the literature on the axiomatic theory of risk measures, one often first designs axioms to identify desirable risk measures without model uncertainty, and then puts model uncertainty into the model as an exogenous object. This approach, although easy to apply, is unsatisfactory from a decision-theoretic point of view, as it does not identify desirable axioms for risk measures when model uncertainty is taken as an input. One of our main contributions is to provide an axiomatic framework of generalized risk measures which allows us to consider properties

on both the model uncertainty and the random losses, thus addressing this practical issue for the first time. The rigorous mathematical formulation of generalized risk measures is laid out in Section 2.

Since generalized risk measures are defined as a mapping from  $\mathcal{X} \times 2^{\mathcal{P}}$  to the (extended) real line, the mathematical structure is much more complicated than traditional risk measures. We establish several relevant theoretical results related to this new framework. In Section 3, we obtain an axiomatic characterization of worst-case generalized risk measures via some simple properties (Theorem 1). The worst-case generalized risk measures are the most practical and they appear extensively in the literature of risk measure and optimization.

Law invariance is a crucial property that connects loss random variables to statistical models. In the traditional framework, law-invariant risk measures can be equivalently expressed as functionals on a space of distributions; this is no longer true in our generalized framework. We provide three different forms of law invariance which reflect different considerations: strong law invariance, loss law invariance, and scenario law invariance; see Section 4 for details. In general, the three notions of law invariance are not equivalent and they reflect very different modeling considerations. Indeed, if strong law invariance is assumed, our framework can be converted to the traditional setting without many mathematical difficulties. However, in practice, strong law invariance may not be desirable, and technical complications arise when it has to be weakened. In Section 4, we show the equivalence between strong law invariance and a combination of the two weaker versions of law invariance under mild conditions (Theorem 2). Moreover, we express the worst-case generalized risk measures with various kinds of law invariance as functions defined on the distributions (Proposition 2). Therefore, from traditional risk measures defined on distributions, we can easily construct generalized risk measures satisfying certain desirable properties.

In Section 5, we focus on coherent generalized risk measures, an analogue to the coherent traditional risk measures of Artzner et al. (1999), and characterize the simplest form (expectation-type) in Theorem 3. Moreover, we propose the notion of ambiguity sensitivity, and establish an equivalence between strong law invariance and a combination of a weaker law invariance and ambiguity sensitivity (Theorem 4). In addition, the combination, with a few simple properties, implies coherence, which supports coherent risk measures in the traditional framework from a completely novel angle.

In Section 6, we discuss some connections of our framework to decision theory. Particularly, we characterize the multi-prior expected utility of Gilboa and Schmeidler (1989) with several properties (Proposition 3) and obtain an axiomatic characterization for the robust generalized risk measure

(Proposition 4), closely related to the variational preferences of [Maccheroni et al. \(2006\)](#). Section 7 contains further discussions and remarks. Proofs of all theorems and propositions are put in the appendix.

## 1.2 Connections to other frameworks in the literature

Our framework is in sharp contrast to the existing ones in the literature of risk management. We have already discussed the difference between our framework and the classic frameworks of risk measures (see [Artzner et al. \(1999\)](#) and [Föllmer and Schied \(2016\)](#)) or preferences (see [Wakker \(2010\)](#) for a comprehensive treatment), which are all defined on  $\mathcal{X}$ . The setting of scenario-based risk measures of [Wang and Ziegel \(2021\)](#) is conceptually close to ours and also motivated by the regulatory framework of [BCBS \(2019\)](#), but mathematically quite different. Scenario-based risk measures are mappings on  $\mathcal{X}$  determined by the distributions of random losses under a collection of pre-specified scenarios. Since the scenarios are fixed, the key question of how a risk measure reacts when scenarios change is left unaddressed.

Model uncertainty is an important topic in economic decision theory. In the classic setting of [Anscombe and Aumann \(1963\)](#), a risk (called a lottery) is represented by a collection of possible distributions, whereas in our framework, the input consists of a random variable and a collection of probability measures, which interact with each other. There are many recent developments in this stream of literature, which focus on characterization of preferences under uncertainty via some axioms. For a non-exclusive list, we mention the multi-prior expected utility of [Gilboa and Schmeidler \(1989\)](#), the multiplier preferences of [Hansen and Sargent \(2001\)](#), the smooth ambiguity preference of [Klibanoff et al. \(2005\)](#), the variational preference of [Maccheroni et al. \(2006\)](#), and the model misspecification preference of [Cerreia-Vioglio et al. \(2021\)](#). They can be formulated as examples of our framework, which will be illustrated in Example 5.

Some conceptual frameworks in decision theory reflect similar considerations towards risk and uncertainty as ours. In particular, [Cerreia-Vioglio et al. \(2021\)](#) studied preferences under model misspecification, and their set of structured models corresponds to our set of scenarios  $\mathcal{Q}$ . An earlier work closely related to our framework is [Gajdos et al. \(2008\)](#), where the authors studied preferences defined on the outcome mapping (an act) and the set of possible probabilities; thus conceptual similarity is clear. Nevertheless, since the main context of our work is financial risk assessment instead of decision making, the axioms and properties considered in this paper, as well as technical results and their implications, are completely different from [Gajdos et al. \(2008\)](#) and [Cerreia-Vioglio et al. \(2021\)](#).

In the operations research literature, [Delage et al. \(2019\)](#) recently investigated a model for decision making with and without uncertainty, and analyzed the conditions under which random decisions are strictly better than deterministic ones. Model uncertainty also widely appears in robust optimization; see [El Ghaoui et al. \(2003\)](#), [Zhu and Fukushima \(2009\)](#) and [Zymler et al. \(2013\)](#) for optimizing risk measures under uncertainty. In the above literature, model uncertainty is generally pre-specified and regarded as an objective fact, whereas we study the properties of risk measures taking model uncertainty as an input argument that can vary over all possible choices.

## 2 A framework for measures of risk and uncertainty

### 2.1 Notation

We begin by stating some notation which will be used throughout. Let  $(\Omega, \mathcal{F})$  be a measurable space and let  $\mathcal{P}$  be the class of atomless probability measures defined on the measurable space.<sup>1</sup> The set of all subsets of  $\mathcal{P}$  is denoted by  $2^{\mathcal{P}}$ . Let  $\mathcal{X}$  be the space of bounded random variables, and  $\mathcal{M}$  be the set of compactly supported distributions on  $\mathbb{R}$ . For  $X, Y \in \mathcal{X}$  and  $P, Q \in \mathcal{P}$ , we write  $X|_P \stackrel{d}{=} Y|_Q$  if the distribution of  $X$  under  $P$  is identical to that of  $Y$  under  $Q$ . We denote by  $F_{X|P} \in \mathcal{M}$  the distribution of  $X$  under  $P$ . For an increasing set function  $\nu : \mathcal{F} \rightarrow \mathbb{R}$  with  $\nu(\emptyset) = 0$ , a Choquet integral (e.g., [Föllmer and Schied \(2016\)](#)) with respect to  $\nu$  is defined as

$$\int X \, d\nu = \int_{-\infty}^0 (\nu(X \geq x) - \nu(\Omega)) \, dx + \int_0^{\infty} \nu(X \geq x) \, dx, \quad X \in \mathcal{X}.$$

### 2.2 A new and generalized framework for risk measures

Traditionally, risk measures in the sense of [Artzner et al. \(1999\)](#) and [Föllmer and Schied \(2016\)](#) are mappings from  $\mathcal{X}$  to  $\mathbb{R}$ . We will call them *traditional* risk measures.

In the new framework that we work with in this paper, the input of a risk measure is a combination of the loss  $X$  and a set of possible probability measures  $\mathcal{Q}$  that represents the best knowledge of the underlying random nature. To distinguish from the traditional setting, we shall refer to these functionals as *generalized* risk measures.

**Definition 1.** A *generalized risk measure*<sup>2</sup> is a mapping  $\Psi : \mathcal{X} \times 2^{\mathcal{P}} \rightarrow [-\infty, \infty]$ . The generalized risk measure  $\Psi$  is called *standard* if  $\Psi(s|\mathcal{Q}) = s$  for all  $s \in \mathbb{R}$  and  $\mathcal{Q} \subseteq \mathcal{P}$ . A *single-scenario risk measure* is a mapping  $\Psi : \mathcal{X} \times \mathcal{P} \rightarrow [-\infty, \infty]$ .

<sup>1</sup>A probability measure  $P$  on  $(\Omega, \mathcal{F})$  is atomless if there exists a uniform random variable on  $(\Omega, \mathcal{F}, P)$ .

<sup>2</sup>We use the notation  $\Psi(X|\mathcal{Q})$  instead of  $\Psi(X, \mathcal{Q})$  to emphasize the different roles of  $X \in \mathcal{X}$  and  $\mathcal{Q} \in 2^{\mathcal{P}}$ .

Clearly, a single-scenario risk measure is precisely a generalized risk measure with its second argument confined to singletons of scenarios. For a singleton  $\{P\} \subseteq \mathcal{P}$ , we use the simpler notation  $\Psi(X|P) = \Psi(X|\{P\})$ . For a fixed  $\mathcal{Q}$ ,  $\Psi(\cdot|\mathcal{Q})$  is a risk measure in the traditional sense.

The requirement of standardization reflects the consideration that, for any fixed constant  $s$ , there is no uncertainty associated with it, and  $\Psi(s|\mathcal{Q})$  should not depend on the input scenarios  $\mathcal{Q}$ . The range of  $\Psi$  is chosen as  $[-\infty, \infty]$  in our general framework to allow for the greatest generality. In practical applications, one may restrict the range to be  $\mathbb{R}$  or  $(-\infty, \infty]$ .

Generalized risk measures are much more complicated as a mathematical object than traditional risk measures, since their input includes both a random loss  $X$  and a set of probability measures  $\mathcal{Q}$ . Below we collect some basic properties to consider for a generalized risk measure  $\Psi$ .

- (A1) Uncertainty aversion:  $\Psi(X|\mathcal{Q}) \leq \Psi(X|\mathcal{R})$  for all  $X \in \mathcal{X}$  and  $\mathcal{Q} \subseteq \mathcal{R} \subseteq \mathcal{P}$ .
- (A2) Scenario monotonicity:  $\Psi(X|\mathcal{Q}) \leq \Psi(Y|\mathcal{Q})$  if  $\Psi(X|P) \leq \Psi(Y|P)$  for all  $P \in \mathcal{Q}$ .
- (A3) Scenario upper bound:  $\Psi(X|\mathcal{Q}) \leq \sup_{P \in \mathcal{Q}} \Psi(X|P)$  for all  $X \in \mathcal{X}$  and  $\mathcal{Q} \subseteq \mathcal{P}$ .

Property (A1) means that the evaluation of the risk weakly increases if model uncertainty increases, and this reflects an aversion to model uncertainty. Property (A2) means that, if under each possible scenario,  $X$  is evaluated to be less risky than  $Y$ , then the overall evaluation of the risk of  $X$  should not be more than  $Y$ . Property (A3) means that the overall evaluation of  $X$  is not more extreme than  $X$  evaluated under the worst-case scenario.

(A2) and (A3) are quite natural and they are satisfied by most examples of generalized risk measures in their various disguises in the risk management and decision theory literature; we will discuss some of them later.

Property (A1) is more specialized, as it will lead to worst-case risk evaluation or decision making (Theorem 1 below) axiomatized in decision theory by Gilboa and Schmeidler (1989). This property is not satisfied in models where uncertainty is aggregated in some form of averaging, such as taking a weighted average of risk evaluates such as the average ES of Wang and Ziegel (2021) or the smooth ambiguity model of Klibanoff et al. (2005). Indeed, if a new scenario  $P$  is added to an existing collection of scenarios  $\mathcal{Q}$ , and a random loss  $X$  is considered safe under  $P$ , then it may be desirable in risk management practice to reduce the assessment of riskiness of  $X$  by including  $P$ , that is,  $\Psi(X|\{\mathcal{Q}, P\}) < \Psi(X|\mathcal{Q})$ , violating (A1).

In decision theory, after a proper translation between two frameworks, the preferential version of (A2) appears in Cerreia-Vioglio et al. (2021) as *Q-separability*, and (A1) is genuinely weaker than *monotonicity in model ambiguity* of Cerreia-Vioglio et al. (2021) on preferences. For a fixed

set of scenarios  $\mathcal{Q}$ , (A1) and (A2) are, respectively, similar to *ambiguity aversion* and *ambiguity monotonicity* in [Delage et al. \(2019\)](#), which are formulated for distributions rather than random variables.

### 2.3 Examples: VaR and ES

We first give a few examples in this section, and more will be discussed later. The two popular traditional risk measures in banking and insurance are the Value-at-Risk (VaR) and the Expected Shortfall (ES); see [Embrechts et al. \(2014\)](#) for a review. Both risk measures in the classic formulation are defined with a fixed scenario  $P \in \mathcal{P}$ , and allowing  $P$  to vary we can treat them as single-scenario risk measures in [Definition 1](#). For a level  $\alpha \in (0, 1]$ , the VaR under  $P$  is defined as

$$\text{VaR}_\alpha(X|P) = \inf\{x \in \mathbb{R} : P(X \leq x) \geq \alpha\}, \quad X \in \mathcal{X},$$

and the ES under  $P$  is defined as

$$\text{ES}_\alpha(X|P) = \frac{1}{1-\alpha} \int_\alpha^1 \text{VaR}_\beta(X|P) \, d\beta, \quad X \in \mathcal{X}.$$

We first show some properties of VaR and ES in our setting. These properties follow from existing properties of VaR and ES with a fixed  $P$ , but the concavity or convexity with respect to scenarios is not formally studied in the literature since our framework is new.

**Proposition 1.** *Fix  $\alpha \in (0, 1)$ . The single-scenario risk measure  $(X, P) \mapsto \text{ES}_\alpha(X|P)$  is convex in  $X$  and concave in  $P$ , whereas  $(X, P) \mapsto \text{VaR}_\alpha(X|P)$  is neither convex nor concave in  $X$  or  $P$ .*

Building on the single-scenario VaR and ES, we can define generalized risk measures such as worst-case VaR and worst-case ES, via

$$\overline{\text{VaR}}_\alpha(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \text{VaR}_\alpha(X|P) \quad \text{and} \quad \overline{\text{ES}}_\alpha(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \text{ES}_\alpha(X|P), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

We refer to [El Ghaoui et al. \(2003\)](#) for optimization of the worst-case VaR, [Zhu and Fukushima \(2009\)](#) for optimization of the worst-case ES, and [Wang and Ziegel \(2021\)](#) for their theoretical properties. The worst-case VaR and the worst-case ES are both standard, and they satisfy (A1), (A2) and (A3).

For a given fixed  $\mathcal{Q} \subseteq \mathcal{P}$ , several other examples of ES and VaR with aggregated scenarios, such as averages (with respect to a pre-specified measure over  $\mathcal{Q}$ ) and inf-convolutions (for a finite  $\mathcal{Q}$ ), are also considered by [Wang and Ziegel \(2021\)](#) and [Castagnoli et al. \(2021\)](#). For instance, we can

define an average ES by

$$(X, \mathcal{Q}) \mapsto \int_{\mathcal{Q}} \text{ES}_{\alpha}(X|Q) \, d\mu_{\mathcal{Q}}(Q), \quad (1)$$

where  $\mu_{\mathcal{Q}}$  is a measure over  $\mathcal{Q}$  for each  $\mathcal{Q} \subseteq \mathcal{P}$ . The average ES in (1) is standard and it satisfies (A2) and (A3); it does not satisfy (A1) in general. We remark that, although sharing many common forms and examples, our framework is fundamentally different from the existing ones in the literature, as it is crucial for a generalized risk measure to use  $\mathcal{Q}$  as an input variable instead of a pre-specified collection.

### 3 Worst-case generalized risk measures

In this section, we present our first theoretical result, a characterization of generalized risk measures satisfying (A1) as the supremum of risk measures in the traditional sense. This result allows us to apply many results on traditional risk measures to generalized risk measures.

**Theorem 1.** *For a generalized risk measure  $\Psi : \mathcal{X} \times 2^{\mathcal{P}} \rightarrow \mathbb{R}$ :*

(i) *Suppose that  $\Psi$  is standard.  $\Psi$  satisfies (A1)-(A2) if and only if it admits a representation*

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \Psi(X|P), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}. \quad (2)$$

(ii)  *$\Psi$  satisfies (A1) and (A3) if and only if it admits a representation (2).*

Using Theorem 1, we can pin down the forms of possible generalized risk measures with properties on the simpler object  $\Psi(X|P)$  for  $X \in \mathcal{X}$  and  $P \in \mathcal{P}$ . Theorem 1 is a general functional form of the specific preferential characterization treated in Theorem 2 of [Cerrei-Vioglio et al. \(2021\)](#).

**Definition 2.** For a given generalized risk measure  $\Psi$ , the single-scenario risk measure  $(X, P) \mapsto \Psi(X|P)$  is called the *core* of  $\Psi$ .

By Theorem 1, the cores one-to-one correspond to standard generalized risk measures that satisfy (A1)-(A2) via (2). Note that in general, the core of  $\Psi$  does not determine  $\Psi$  on  $\mathcal{X} \times 2^{\mathcal{P}}$ , if the conditions in Theorem 1 are not satisfied.

In case (2) holds, we say that the core  $\Psi$  on  $\mathcal{X} \times \mathcal{P}$  induces the generalized risk measure  $\Psi$  on  $\mathcal{X} \times 2^{\mathcal{P}}$ . Many results in this paper are stated for cores instead of the generalized risk measure. Nevertheless, when we speak of cores, we do not need to assume the worst-case form (2) or any of (A1)-(A3).

Some simple examples for the worst-case generalized risk measures are collected below, and they appear in forms similar to those in the classic theory of risk measures.

**Example 1.** (i) The expectation core

$$\Psi(X|P) = \mathbb{E}^P[X], \quad (X, P) \in \mathcal{X} \times \mathcal{P}$$

induces the generalized risk measure

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \mathbb{E}^P[X], \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

For a fixed  $\mathcal{Q}$ ,  $\Psi(\cdot|\mathcal{Q})$  is the robust representation of a traditional coherent risk measure of [Artzner et al. \(1999\)](#). This class of risk measures is the most well studied in the literature, and we will pay special attention to it in [Section 5](#).

(ii) Let  $\gamma : \mathcal{P} \rightarrow \mathbb{R}$  be a non-constant function on  $\mathcal{P}$ . The penalized-mean core

$$\Psi(X|P) = \mathbb{E}^P[X] - \gamma(P), \quad (X, P) \in \mathcal{X} \times \mathcal{P}$$

induces the generalized risk measure

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \{\mathbb{E}^P[X] - \gamma(P)\}, \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

For a fixed  $\mathcal{Q}$ ,  $\Psi(\cdot|\mathcal{Q})$  is the robust representation of a traditional convex risk measure of [Föllmer and Schied \(2016\)](#).

(iii) For  $\alpha \in (0, 1)$ , the VaR core  $(X, P) \mapsto \text{VaR}_\alpha(X|P)$  induces the worst-case VaR in [Section 2.3](#).

(iv) For  $\alpha \in (0, 1)$ , the ES core  $(X, P) \mapsto \text{ES}_\alpha(X|P)$  induces the worst-case ES in [Section 2.3](#).

## 4 Three formulations of law invariance

For a given  $P$ , the functional  $X \mapsto \Psi(X|P)$  is a traditional risk measure, and properties can be imposed for this traditional risk measure. The more interesting and non-trivial question is the interplay between  $X$  and  $P$  for the core  $\Psi$ , which we will address below. Since  $P \in \mathcal{P}$  is interpreted as a scenario for us to generate a statistical model for the loss  $X$ , the evaluation of the risk should depend on the distribution of  $X$ . Motivated by this consideration, we can consider three forms of

law invariance on the generalized risk measure  $\Psi$  or its core:<sup>3</sup>

(B1) Strong law invariance:  $\Psi(X|P) = \Psi(Y|Q)$  for  $X, Y \in \mathcal{X}$  and  $P, Q \in \mathcal{P}$  with  $X|_P \stackrel{d}{=} Y|_Q$ .

(B2) Loss law invariance:  $\Psi(X|P) = \Psi(Y|P)$  for  $X, Y \in \mathcal{X}$  and  $P \in \mathcal{P}$  with  $X|_P \stackrel{d}{=} Y|_P$ .

(B3) Scenario law invariance:  $\Psi(X|P) = \Psi(X|Q)$  for  $X \in \mathcal{X}$  and  $P, Q \in \mathcal{P}$  with  $X|_P \stackrel{d}{=} X|_Q$ .

Clearly, (B1) is stronger than both (B2) and (B3). Each of (B1), (B2) and (B3) reflects the consideration that the probability measures  $P$  in  $\Psi(X|P)$  is used to model the distribution of the loss  $X$ . More precisely, (B1) is an agreement of risk assessment for the same distribution across different scenarios and different losses, whereas (B2) only yields the agreement for each particular scenario, and (B3) only yields the agreement for each particular loss. The following example shows that (B1), (B2) and (B3) are genuinely different concepts.

**Example 2.** (i) The cores in Example 1 (i), (iii) and (iv) are strongly law-invariant.

(ii) The core in Example 1 (ii) is loss law-invariant, but generally not scenario law-invariant.

(iii) Let  $\beta : \mathcal{X} \rightarrow \mathbb{R}$  be a non-constant function on  $\mathcal{X}$ . The core

$$\Psi(X|P) = \mathbb{E}^P[X] - \beta(X), \quad (X, P) \in \mathcal{X} \times \mathcal{P}$$

is scenario law-invariant, but generally not loss law-invariant.

Since (B1) implies both (B2) and (B3), one may wonder whether (B2) and (B3) jointly imply (B1), which turns out to be a tricky question. In other words, we aim to show from (B2) and (B3) that  $\Psi(X|P) = \Psi(Y|Q)$  holds for  $P, Q \in \mathcal{P}$  and  $X, Y \in \mathcal{X}$  satisfying  $X|_P \stackrel{d}{=} Y|_Q$ . Denote by  $F$  the distribution of  $X$  under  $P$ , which is the same as that of  $Y$  under  $Q$ . If there exists  $Z \in \mathcal{X}$  which has distribution  $F$  under both  $P$  and  $Q$ , then we have the desired chain of equalities  $\Psi(X|P) = \Psi(Z|P) = \Psi(Z|Q) = \Psi(Y|Q)$ . Unfortunately, the existence of such  $Z$  depends on the specification of  $P, Q$  and it cannot be expected in general; this problem is non-trivial and has been studied in detail by [Shen et al. \(2019\)](#). In the result below, we show that, under the extra assumption that the measurable space  $(\Omega, \mathcal{F})$  is standard Borel (i.e., isomorphic to the Borel space on  $[0, 1]$ ), it is possible to find an intermediate measure  $R$  and two random variables  $Z, W \in \mathcal{X}$  such

---

<sup>3</sup>In this paper, all properties (Ax) reflect how  $\Psi$  reacts to  $\mathcal{Q}$ , all properties (Bx) reflect how  $\Psi$  reacts to the distributions of the risk, all properties (Cx) reflect consideration for  $\Psi$  in terms of a traditional risk measure, all properties (Dx) are relevant to a mapping defined on the set of measures  $\mathcal{P}$ , and all properties (Ex) reflect consideration on decision-theoretic preference.

that the chain of equalities

$$X|_P \stackrel{d}{=} Z|_P \stackrel{d}{=} Z|_R \stackrel{d}{=} W|_R \stackrel{d}{=} W|_Q \stackrel{d}{=} Y|_Q$$

holds, and it gives the desired statement  $\Psi(X|P) = \Psi(Y|Q)$  needed for (B1).

**Theorem 2.** *For a core  $\Psi$ , (B1) implies both (B2) and (B3). If  $(\Omega, \mathcal{F})$  is standard Borel, then (B2) and (B3) together are equivalent to (B1).*

*Remark 1.* It remains unclear whether the equivalence  $(B2+B3) \Leftrightarrow (B1)$  holds if  $(\Omega, \mathcal{F})$  is not standard Borel. For applications in finance and risk management, it is typically sufficient to use a standard Borel space, because one can construct countably many independent Brownian motions on the corresponding probability space. The assumption of a standard Borel space is used in some classic literature on risk measures, e.g., [Delbaen \(2002\)](#) and [Jouini et al. \(2006\)](#).

Loss law invariance (B2) seems to be always desirable to assume in practice, because if two random losses  $X$  and  $Y$  share the same distribution under a chosen scenario  $P$  of interest, then it is natural to assign the same risk value to these two losses. For a fixed collection  $\mathcal{Q} \in 2^{\mathcal{P}}$ , this property defines a  $\mathcal{Q}$ -based risk measure of [Wang and Ziegel \(2021\)](#). On the other hand, it may not always be desirable to assume (B3); although two scenarios may give the same distribution of a loss  $X$ , the riskiness may not be understood as the same, as illustrated by the following example.

**Example 3** ((B3) is not always desirable). Let  $P$  represent a good economic scenario, and  $Q$  represent an adverse economic scenario (e.g., COVID-19). Assume that the distribution of  $X$  is the same under  $P$  and  $Q$ , which means that  $X$  is independent of the particular economic factor which generates  $P$  and  $Q$ . The value  $\Psi(X|P)$  quantifies the riskiness of  $X$  when  $P$  is the chosen scenario, and  $\Psi(X|Q)$  quantifies the riskiness of  $X$  when  $Q$  is the chosen scenario. Since  $P$  is a better economy, the risk manager may think that  $X$  is more acceptable in this situation, leading to  $\Psi(X|P) < \Psi(X|Q)$ . For instance, the core in [Example 1 \(ii\)](#), the robust representation of convex risk measures, reflects this consideration, and it is not scenario law-invariant.

Next, we collect some representation results based on (A1), (A3) and (B1)-(B3). Throughout, we define

$$\Sigma = \{\psi : \mathcal{M} \rightarrow [-\infty, \infty]\}.$$

Each mapping  $\psi \in \Sigma$  represents a traditional law-invariant risk measure treated as a functional on  $\mathcal{M}$  instead of on  $\mathcal{X}$ .

**Proposition 2.** *Let  $\Psi$  be a generalized risk measure.*

(i)  $\Psi$  satisfies (A1), (A3) and (B1) if and only if there exists  $\psi \in \Sigma$  such that

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \psi(F_{X|P}), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

(ii)  $\Psi$  satisfies (A1), (A3) and (B2) if and only if there exists  $\{\psi_P : P \in \mathcal{P}\} \subseteq \Sigma$  such that

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \psi_P(F_{X|P}), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

(iii)  $\Psi$  satisfies (A1), (A3) and (B3) if and only if there exists  $\{\psi_X : X \in \mathcal{X}\} \subseteq \Sigma$  such that

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \psi_X(F_{X|P}), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}.$$

## 5 Coherent generalized risk measures

In this section, we pay special attention to the most important class of traditional risk measures: coherent risk measures of Artzner et al. (1999). We first provide a characterization for a generalized risk measure to have the form of coherent risk measures in Example 1, and then we discuss a few additional properties that are specific to our setting.

### 5.1 A characterization for coherent risk measures

We give a simple characterization of the coherent risk measures in Example 1 (i). Coherent risk measures including ES (for a fixed scenario) are the most well-studied class of risk measures in the finance and engineering literature. We first list some properties of traditional risk measures of Artzner et al. (1999) and Föllmer and Schied (2016). These properties are formulated for the traditional risk measure  $X \mapsto \Psi(X|\mathcal{Q})$  on  $\mathcal{X}$  for each fixed  $\mathcal{Q} \subseteq \mathcal{P}$ , and we denote this by  $\Psi_{\mathcal{Q}}$ .

(C1) Monotonicity:  $\Psi_{\mathcal{Q}}(X) \leq \Psi_{\mathcal{Q}}(Y)$  for all  $X, Y \in \mathcal{X}$  with  $X \leq Y$ ;

(C2) Cash-additivity:  $\Psi_{\mathcal{Q}}(X + m) = \Psi_{\mathcal{Q}}(X) + m$  for all  $X \in \mathcal{X}$  and  $m \in \mathbb{R}$ ;

(C3) Positive homogeneity:  $\Psi_{\mathcal{Q}}(\lambda X) = \lambda \Psi_{\mathcal{Q}}(X)$  for all  $\lambda > 0$  and  $X \in \mathcal{X}$ ;

(C4) Subadditivity:  $\Psi_{\mathcal{Q}}(X + Y) \leq \Psi_{\mathcal{Q}}(X) + \Psi_{\mathcal{Q}}(Y)$  for all  $X, Y \in \mathcal{X}$ .

Following the terminology for traditional risk measures, a generalized risk measure  $\Psi$  is *monetary* if it satisfies (C1)-(C2) and *coherent* if it satisfies (C1)-(C4). We further state a strong property imposed on the cores.

(C0) Additivity of the core:  $\Psi(X + Y|P) = \Psi(X|P) + \Psi(Y|P)$  for all  $X, Y \in \mathcal{X}$  and  $P \in \mathcal{P}$ .

The property (C0) will be a key property to pin down the form of coherent traditional risk measures.

**Theorem 3.** *A standard generalized risk measure  $\Psi$  satisfies (A1), (A2), (B2), (C1) and (C0) if and only if it is uniquely given by*

$$\Psi(X|Q) = \sup_{P \in Q} \mathbb{E}^P[X], \quad (X, Q) \in \mathcal{X} \times 2^{\mathcal{P}}. \quad (3)$$

Moreover,  $\Psi$  in (3) satisfies (C2)-(C4).

The most important property used in Theorem 3 is the additivity of the core (C0), which may be seen as quite strong. As a primary example of coherent risk measures of the form (3) in financial practice, the Chicago Mercantile Exchange (CME) uses (3) to determine margin requirements for portfolio of instruments; see McNeil et al. (2015, p.64). In the CME approach, under each fixed scenario, the risk factors move in a particular deterministic way, and hence the portfolio loss assessment is additive; thus (C0) is natural in this context.

## 5.2 Ambiguity sensitivity and comonotonic additive risk measures

As we have seen from Example 2, strong law invariance (B1) is genuinely stronger than the weaker notions of (B2) and (B3). In the following result, we connect weak and strong law invariance via an additional property on the generalized risk measure  $\Psi$  via its core.

(B4) Ambiguity sensitivity: For  $X \in \mathcal{X}$ ,  $P, Q \in \mathcal{P}$  and  $\lambda \in [0, 1]$ ,  $\Psi(X|\lambda P + (1 - \lambda)Q) \geq \lambda \Psi(X|P) + (1 - \lambda)\Psi(X|Q)$ . Moreover,  $\Psi(\mathbb{1}_A|\lambda P + (1 - \lambda)Q) = \lambda \Psi(\mathbb{1}_A|P) + (1 - \lambda)\Psi(\mathbb{1}_A|Q)$  for all  $A \in \mathcal{F}$  such that  $P(A) = Q(A)$ .

The first statement of (B4) intuitively means that, due to ambiguity on the distribution of  $X$ , the risk of  $X$  under a mixture is larger than the mixture of its risks under  $P$  and  $Q$ ; this is the concavity in  $P$  in Proposition 1. For instance, for a random variable  $X$  which is constant under both  $P$  and  $Q$ , it may be random (Bernoulli) under  $\lambda P + (1 - \lambda)Q$ , and hence its risk should be larger under the mixture than under the individual scenarios. Regarding the second statement of (B4), if the probability measures  $P$  and  $Q$  agree on how likely event  $A$  is, then there is no ambiguity on  $A$ , and its risk under a mixture should be simply a mixture of its risks under  $P$  and  $Q$ . Another explanation may be illustrated by the following example.

**Example 4** (Ambiguity sensitivity). Assume that  $P$  is used by a risk analyst and  $Q$  is used by another risk analyst. The manager would like to use  $\lambda P + (1 - \lambda)Q$ , a mixture of  $P$  and  $Q$ , to reflect

the knowledge of both analysts. For simplicity, the random loss  $X$  is assumed to be an indicator of a loss event  $A$ . If  $P$  and  $Q$  give different assessments of the probability of  $A$ , then the manager would be worried about the discrepancy in the models, and her final risk assessment  $\Psi(X|\lambda P + (1 - \lambda)Q)$  is more than  $\lambda\Psi(X|P) + (1 - \lambda)\Psi(X|Q)$ , the weighted average of the two analysts' assessments. On the other hand, if  $P$  and  $Q$  give the same probability of  $A$ , then there is no disagreement in predicting  $A$ . In this case, her risk assessment of  $\mathbb{1}_A$  is the same as the weighted average of the two analysts' assessments.

Another property that is essential to our next characterization result is comonotonic additivity, which is intimately linked to Choquet integrals; see e.g., [Wang et al. \(2020\)](#).

(C5) Comonotonic additivity:  $\Psi_Q(X + Y) = \Psi_Q(X) + \Psi_Q(Y)$ , for all  $X, Y \in \mathcal{X}$  which are comonotonic.<sup>4</sup>

The following result characterizes loss law-invariant risk measures with ambiguity sensitivity, which turns out to be equivalent to strongly law-invariant risk measures without this assumption. The proof of this result is quite technical and it relies on Lyapunov's convexity theorem, as well as a few characterization results on Choquet integrals in [Wang et al. \(2020\)](#).

**Theorem 4.** *For a core  $\Psi$ , the following are equivalent:*

- (i)  $\Psi$  is loss law-invariant, ambiguity sensitive, monetary, and comonotonic additive, i.e.,  $\Psi$  satisfies (B2), (B4), (C1), (C2) and (C5).
- (ii)  $\Psi$  is strongly law-invariant, coherent, and comonotonic additive, i.e.,  $\Psi$  satisfies (B1) and (C1)-(C5).
- (iii) There exists an increasing concave function  $h : [0, 1] \rightarrow [0, 1]$  with  $h(0) = 0 = 1 - h(1)$  such that

$$\Psi(X|P) = \int X \, d(h \circ P), \quad (X, P) \in \mathcal{X} \times \mathcal{P}. \quad (4)$$

There has been an extensive debate in both academia and industry on whether subadditivity (C4) proposed by [Artzner et al. \(1999\)](#) is a good criterion for risk measures used in regulatory practice, as (C4) is key property which distinguishes VaR and ES; see [Embrechts et al. \(2018, 2021\)](#) and the reference therein. By Theorem 4, from the perspective of multiple models, we can obtain (C4) by using ambiguity sensitivity (B4). Hence, our framework and results offer a novel decision-theoretic reason to support coherent risk measures (in particular, ES over VaR) without directly assuming subadditivity (C4).

---

<sup>4</sup>Two random variables  $X$  and  $Y$  are *comonotonic*  $(X(\omega) - X(\omega'))(Y(\omega) - Y(\omega')) \geq 0$  for all  $(\omega, \omega') \in \Omega \times \Omega$ .

## 6 Connection to decision theory

In this section, we discuss the connection of our generalized risk measures to classic notions in decision theory, as model uncertainty has been dealt with extensively in the decision theoretic literature, and traditional risk measures are intimately linked to decision preferences in various forms; see e.g., [Drapeau and Kupper \(2013\)](#). We first present a list of decision-theoretic criteria as examples to our framework, followed by characterization results of two classic notions: the multi-prior expected utility of [Gilboa and Schmeidler \(1989\)](#) and the variational preferences of [Maccheroni et al. \(2006\)](#).

### 6.1 Examples of generalized risk measures in decision theory

Our framework includes many criteria in decision theory as typical examples. Although the considerations of these criteria are different from our paper, the following examples show the generality of our framework.

**Example 5.** (i) The multi-prior expected utility of [Gilboa and Schmeidler \(1989\)](#) has a numerical representation

$$\Psi(X|\mathcal{Q}) = u^{-1} \left( \min_{P \in \mathcal{Q}} \mathbb{E}^P[u(X)] \right),$$

where  $u$  is a strictly increasing utility function.

(ii) The variational preference of [Maccheroni et al. \(2006\)](#) has a numerical representation

$$\Psi(X|\mathcal{Q}) = \min_{P \in \mathcal{Q}} (\mathbb{E}^P[u(X)] - \gamma(P)),$$

where  $u$  is a strictly increasing utility function and  $\gamma : \mathcal{P} \rightarrow [-\infty, \infty)$  is a penalty function. The multiplier preferences of [Hansen and Sargent \(2001\)](#) correspond to a special choice of  $\gamma$  which is the Kullback–Leibler divergence from a reference scenario.

(iii) Let  $\mathcal{Q} \subseteq \mathcal{P}$  be pre-specified and  $\mu$  be a probability measure on  $\mathcal{Q}$ . The smooth ambiguity preference of [Klibanoff et al. \(2005\)](#) has a numerical representation

$$\Psi(X|\mathcal{Q}) = \phi^{-1} \left( \int_{\mathcal{Q}} \phi(u^{-1}(\mathbb{E}^P[u(X)])) \, d\mu(P) \right),$$

where  $u$  is a strictly increasing utility function and  $\phi$  is a strictly increasing function. Note that in this formulation,  $\mu$  needs to be specified with  $\mathcal{Q}$ , and hence it should be considered as an input of  $\Psi$  in our framework; see [Section 7](#) for more discussion on this.

(iv) The imprecise information preference of [Gajdos et al. \(2008\)](#) has a numerical representation

$$\Psi(X|\mathcal{Q}) = u^{-1} \left( \min_{P \in \phi(\mathcal{Q})} \mathbb{E}^P[u(X)] \right),$$

where  $u$  is a strictly increasing utility function and  $\phi$  is a selecting function (assumed to exist) reflecting the decision maker's attitude to imprecision.

(v) The model misspecification preference of [Cerrea-Vioglio et al. \(2021\)](#) has a numerical representation

$$\Psi(X|\mathcal{Q}) = \min_{P \in \mathcal{P}} \left\{ \mathbb{E}^P[u(X)] + \min_{Q \in \mathcal{Q}} c(P, Q) \right\},$$

where  $u$  is a strictly increasing utility function and  $c$  is a distance on the set of measures which penalizes the model misspecification.

## 6.2 Multi-prior expected utilities

[Gilboa and Schmeidler \(1989\)](#) proposed the notion of *multi-prior expected utility* in decision theory. Motivated by the multi-prior expected utility, we consider a preference on  $\mathcal{X} \times \mathcal{S}$  which is represented by a total pre-order  $\preceq$ , where  $\mathcal{S}$  is the collection of all finite subsets of  $\mathcal{P}$ . For tractability, we consider  $\mathcal{S}$ , all the finite subsets of  $\mathcal{P}$ , instead of  $2^{\mathcal{P}}$  in this subsection. The decision is to compare a risk and a set of scenarios with another risk and another set of scenarios. This setting was studied by [Gajdos et al. \(2008\)](#). We denote by  $\simeq$  the equivalence under this preference. As above, we use  $(X, P)$  if the set of scenarios has only one element  $P$ . For decisions among  $(X_1, \mathcal{Q}_1), (X_2, \mathcal{Q}_2) \in \mathcal{X} \times \mathcal{S}$ , we propose the following axioms similar to what we have seen so far in this paper, but defined for preferences instead of generalized risk measures.

(E1) Strong law invariance:  $(X, P) \simeq (Y, Q)$  for any  $P, Q \in \mathcal{P}$  and  $X, Y \in \mathcal{X}$  satisfying  $X|_P \stackrel{d}{=} Y|_Q$ .

(E2) Uncertainty aversion:  $(X, \mathcal{Q}) \preceq (X, \mathcal{R})$  for any  $X \in \mathcal{X}$  and  $\mathcal{R}, \mathcal{Q} \in \mathcal{S}$  with  $\mathcal{R} \subseteq \mathcal{Q}$ .

(E3) Uncertainty bound: for any  $X \in \mathcal{X}$  and  $\mathcal{Q} \in \mathcal{S}$ , there exists some  $P \in \mathcal{Q}$  such that  $(X, P) \preceq (X, \mathcal{Q})$ .

(E4) Independence: for any  $P, Q \in \mathcal{P}$ , any  $X, Y \in \mathcal{X}$  satisfying  $X|_Q \stackrel{d}{=} Y|_Q$ , and any  $\alpha \in (0, 1)$  we have  $(X, P) \preceq (Y, P) \iff (X, \alpha P + (1 - \alpha)Q) \preceq (Y, \alpha P + (1 - \alpha)Q)$ .

(E5) Continuity: for any  $P, Q, R \in \mathcal{P}$  and any  $X \in \mathcal{X}$ , if  $(X, P) \preceq (X, Q) \preceq (X, R)$ , then there exists  $\alpha \in [0, 1]$  such that  $(X, \alpha P + (1 - \alpha)R) \simeq (X, Q)$ .

Proposition 3 illustrates a decision-theoretic characterization for the multi-prior expected utility. The proof is based on Theorem 1 and the classic result of Von Neumann and Morgenstern (1944).

**Proposition 3.** *A preference  $\preceq$  on  $\mathcal{X} \times \mathcal{S}$  satisfies (E1)-(E5) if and only if it is a multi-prior expected utility, i.e., there exists a function  $u : \mathbb{R} \rightarrow \mathbb{R}$  such that*

$$(X_1, \mathcal{Q}_1) \preceq (X_2, \mathcal{Q}_2) \iff \min_{P \in \mathcal{Q}_1} \mathbb{E}^P[u(X_1)] \leq \min_{P \in \mathcal{Q}_2} \mathbb{E}^P[u(X_2)]. \quad (5)$$

The strong law invariance (E1), which allows us to translate  $\preceq$  to a preference on the set of distributions on the real line, is crucial to the representation result. (E2) and (E3) are reasonable for uncertainty-averse decision makers, and they correspond to (A1) and (A3), respectively, in the framework of generalized risk measures. (E4) and (E5) correspond to the independence axiom and the continuity axiom of Von Neumann and Morgenstern (1944), respectively.

### 6.3 Robust generalized risk measures

In addition to the worst-case generalized risk measure characterized in Theorem 1, another popular form of risk measures involving multiple probability measures arises from the robust representation of convex risk measures as in Example 1 (ii). More precisely, a traditional convex risk measure  $\rho$  of Föllmer and Schied (2016) takes the form, for some  $\mathcal{Q} \subseteq \mathcal{P}$ ,

$$\rho(X) = \sup_{P \in \mathcal{Q}} \{\mathbb{E}^P[X] - \gamma(P)\}, \quad X \in \mathcal{X}, \quad (6)$$

where  $\gamma : \mathcal{P} \rightarrow (-\infty, \infty]$  is a penalty function. Moreover, the variational preference of Maccheroni et al. (2006) takes a similar form to (6) with the mean  $\mathbb{E}$  replaced by an expected utility;<sup>5</sup> see Example 5 (ii). Inspired by (6) and the variational preference of Maccheroni et al. (2006), we consider generalized risk measures with the form, for some  $\psi \in \Sigma$ ,

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \{\psi(F_{X|P}) - \gamma(P)\}, \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}. \quad (7)$$

Clearly, if  $\psi$  is the mean functional, then (7) yields the traditional (convex) risk measure (6) for a given  $\mathcal{Q}$ . The generalized risk measure in (7) is loss law-invariant (B2) but neither scenario law-invariant (B3) nor strongly law-invariant (B1). In order to characterize (7), we further impose the following technical property, which says that the difference between the values of the core evaluated on  $P$  and  $Q$  for identically distributed losses only depends on  $P$  and  $Q$  but not the random losses.

---

<sup>5</sup>In the setting of numerical representation of preferences, a negative sign needs to be applied to the generalized risk measures to transform it to a preference functional.

(B5) If  $X|_P \stackrel{d}{=} Y|_Q$  and  $Z|_P \stackrel{d}{=} W|_Q$ , then  $\Psi(X|P) - \Psi(Y|Q) = \Psi(Z|P) - \Psi(W|Q)$ .

**Proposition 4.** *Let  $\Psi$  be a generalized risk measure.  $\Psi$  satisfies (A1), (A3), (B2) and (B5) if and only if there exist a penalty function  $\gamma : \mathcal{P} \rightarrow \mathbb{R}$  and some  $\psi \in \Sigma$  such that the representation (7) holds.*

Property (B5) can be roughly interpreted as that the magnitude of penalization for a given scenario  $P$  is independent of the risky position  $X$  being evaluated. This property may be seen as a bit artificial. Our characterization in Proposition 4 is mainly motivated by the great popularity of the robust representation of convex risk measures and variational preferences, and we omit a detailed discussion of the economic desirability or undesirability of (B5).

## 7 Concluding remarks

The new framework of generalized risk measures introduced in this paper allows for a unified systemic formulation of measures of risk and uncertainty. Our results are only first attempts to understand the new setting, and many further questions arise, especially regarding the interplay between the risk variable  $X$  and the uncertainty collection  $\mathcal{Q}$  for a generalized risk measures. Both new economic and mathematical questions arise as the new functionals are more sophisticated than traditionally studied objects by definition.

Worst-case generalized risk measures are characterized with a few axioms in Theorem 1. Another popular way of handling model uncertainty is to use a weighted average of the risk evaluations. In case of a finite collection  $\mathcal{Q}$ , we can always use an arithmetic average as the risk evaluation, that is, to generate  $\Psi$  via its core by

$$\Psi(X|\mathcal{Q}) = \frac{1}{|\mathcal{Q}|} \sum_{Q \in \mathcal{Q}} \Psi(X|Q).$$

Certainly, such a formulation does not satisfy (A1) but it satisfies (A2) and (A3). In general, to allow for different weights and infinite collections, one needs to associate each collection  $\mathcal{Q}$  with a measure as in (1) or the smooth ambiguity preference of Klibanoff et al. (2005) in Example 5 (iii). Such a measure can either be pre-specified or treated as an input of  $\Psi$ , thus slightly extending our framework.

We studied several most popular properties such as law-invariance, coherence and comonotonic additivity, but many more properties on the new framework remain to be explored, as the literature on traditional risk measures is very rich. In particular, the desirability of theoretic properties in

risk management practice requires thorough study, as they may have different interpretation from its traditional counterpart. For instance, additivity of the core may be sensible in our framework (Theorem 3) and it nicely connects to the scenario-based margin calculation used by CME. However, such a property is not desirable for traditional risk measures, as it essentially forces the risk measure to collapse to the mean; see e.g., Liebrich and Munari (2021) and Chen et al. (2021) .

Finally, we mention that in some formulations of generalized risk measures, not all choices of the input scenario  $\mathcal{Q}$  are economically meaningful. In particular, for a given penalty function  $\gamma$  on  $\mathcal{P}$ , the core  $\Psi(X|P) = \mathbb{E}^P[X] - \gamma(P)$  in Example 1 (ii) or  $\Psi(X|P) = \mathbb{E}^P[u(X)] - \gamma(P)$  in Example 5 (ii) is not meant to be used directly with a single  $P$ ; the use of  $\gamma$  already implicitly implies that there are some level of model uncertainty, and it is supposed to be coupled with the worst-case operation. The value  $\Psi(X|P)$  for a standalone  $P$  is thus difficult to interpret, and should not be used for decision making without properly specifying the uncertainty collection  $\mathcal{Q}$ . On the other hand, such a situation does not happen in, for instance, the worst-case or average-type generalized risk measures based on traditional risk measures, such as the worst-case ES.

## Acknowledgments

YL and RW acknowledge financial support from the Natural Sciences and Engineering Research Council of Canada (RGPIN-2018-03823, RGPAS-2018-522590).

## References

- Anscombe, F. J. and Aumann, R. J. (1963). A definition of subjective probability. *Annals of Mathematical Statistics*, **34**(1), 199–205.
- Artzner, P., Delbaen, F., Eber, J.-M. and Heath, D. (1999). Coherent measures of risk. *Mathematical Finance*, **9**(3), 203–228.
- BCBS (2019). *Minimum Capital Requirements for Market Risk. February 2019*. Basel Committee on Banking Supervision. Document d457, Basel: Bank for International Settlements.
- Castagnoli, E., Cattelan, G., Maccheroni, F., Tebaldi, C. and Wang, R. (2021). Star-shaped risk measures. *arXiv*: 2103.15790.
- Correia-Vioglio, S., Hansen, L. P., Maccheroni, F. and Marinacci, M. (2021). Making decisions under model misspecification. *SSRN*: 3666424.
- Chen, S., Gao, N., Leung, D. H. and Li, L. (2021). Do law-invariant linear functionals collapse to the mean? *arXiv*: 2107.11239.

- Cambou, M. and Filipovic, D. (2017). Model uncertainty and scenario aggregation. *Mathematical Finance*, **27**(2), 534–567.
- Delage, E., Kuhn, D. and Wiesemann, W. (2019). “Dice”-sion-making under uncertainty: When can a random decision reduce risk? *Management Science*, **65**(7), 3282–3301.
- Delbaen, F. (2002). Coherent risk measures on general probability spaces. In *Advances in Finance and Stochastics* (pp. 1–37). Springer, Berlin, Heidelberg.
- Delbaen, F. (2012). *Monetary Utility Functions*. Osaka University Press, Osaka.
- Drapeau, S. and Kupper, M. (2013). Risk preferences and their robust representation. *Mathematics of Operations Research*, **38**(1), 28–62.
- Durrett, R. (2010). *Probability: Theory and Examples*. Cambridge University Press. Fourth Edition.
- El Ghaoui, L., Oks, M. and Oustry, F. (2003). Worst-case value-at-risk and robust portfolio optimization: A conic programming approach. *Operations Research*, **51**(4), 543–556.
- Embrechts, P., Puccetti, G., Rüschendorf, L., Wang, R. and Beleraj, A. (2014). An academic response to Basel 3.5. *Risks*, **2**(1), 25–48.
- Embrechts, P., Liu, H. and Wang, R. (2018). Quantile-based risk sharing. *Operations Research*, **66**(4), 936–949.
- Embrechts, P., Schied, A. and Wang, R. (2021). Robustness in the optimization of risk measures. *Operations Research*, forthcoming.
- Föllmer, H. and Schied, A. (2016). *Stochastic Finance. An Introduction in Discrete Time*. Walter de Gruyter, Berlin, Fourth Edition.
- Gajdos, A.T., Hayashi, T., Tallon, J.-M. and Vergnaud, J.-C. (2008). Attitude toward imprecise information. *Journal of Economic Theory*, **140**(1), 27–65.
- Gilboa, I. and Schmeidler, D. (1989). Maxmin expected utility with a non-unique prior. *Journal of Mathematical Economics*, **18**, 141–153.
- Hansen, L. P. and Sargent, T. J. (2001). Robust control and model uncertainty. *American Economic Review*. **91**(2), 60–66.
- Jouini, E., Schachermayer, W. and Touzi, N. (2006). Law invariant risk measures have the Fatou property. *Advances in Mathematical Economics*, **9**, 49–71.
- Klibanoff, P., Marinacci, M. and Mukerji, S. (2005). A smooth model of decision making under uncertainty. *Econometrica*, **73**(6), 1849–1892.
- Liebrich, F. B. and Munari, C. (2021). Law-invariant functionals that collapse to the mean: Beyond convexity. *arXiv*: 2106.01281.
- Maccheroni, F., Marinacci, M. and Rustichini, A. (2006). Ambiguity aversion, robustness, and the

- variational representation of preferences. *Econometrica*, **74**(6), 1447–1498.
- McNeil, A. J., Frey, R. and Embrechts, P. (2015). *Quantitative Risk Management: Concepts, Techniques and Tools*. Revised Edition. Princeton, NJ: Princeton University Press.
- Von Neumann, J. and Morgenstern, O. (1944). *Theory of Games and Economic Behavior*. Princeton, NJ: Princeton University Press.
- Rudin, W. (1991). *Functional Analysis*. Second Ed., McGraw-Hill, New York.
- Shen, J., Shen, Y., Wang, B. and Wang, R. (2019). Distributional compatibility for change of measures. *Finance and Stochastics*, **23**(3), 761–794.
- Wakker, P. P. (2010). *Prospect Theory: For Risk and Ambiguity*. Cambridge University Press.
- Wang, R., Wei, Y. and Willmot, G. E. (2020). Characterization, robustness and aggregation of signed Choquet integrals. *Mathematics of Operations Research*, **45**(3), 993–1015.
- Wang, R. and Ziegel, J. F. (2021). Scenario-based risk evaluation. *Finance and Stochastics*, **25**, 725–756.
- Wang, R. and Zitikis, R. (2021). An axiomatic foundation for the expected shortfall. *Management Science*, **67**(3), 1413–1429.
- Zhu, S. and Fukushima, M. (2009). Worst-case conditional value-at-risk with application to robust portfolio management. *Operations Research*, **57**(5), 1155–1168.
- Zymler, S., Kuhn, D. and Rustem, B. (2013). Worst-case value-at-risk of nonlinear portfolios. *Management Science*, **59**(1), 172–188.

## A Proofs of all technical results

*Proof of Proposition 1.* For a fixed  $P$ , convexity of  $\text{ES}_\alpha(\cdot|P)$  is well-known since  $\text{ES}_\alpha(\cdot|P)$  is a coherent risk measure; see e.g., Föllmer and Schied (2016). The non-convexity and non-concavity of  $\text{VaR}_\alpha(\cdot|P)$  are due to the fact that  $\text{VaR}_\alpha(\cdot|P)$  has a non-convex and non-concave distortion function; see e.g., Theorem 3 of Wang et al. (2020). For a fixed  $X$ , note that a mixture on scenarios leads to a mixture of the distribution of  $X$ , that is, the distribution of  $X$  under  $\lambda P + (1 - \lambda)Q$  is  $\lambda F_{X|P} + (1 - \lambda)F_{X|Q}$ . Thus, concavity with respect to scenarios corresponds to mixture concavity studied in Wang et al. (2020). Again, using Theorem 3 of Wang et al. (2020),  $\text{ES}_\alpha(X|\cdot)$  is concave and  $\text{VaR}_\alpha(X|\cdot)$  is neither convex nor concave.  $\square$

*Proof of Theorem 1.* (i) The “if” statement can be directly checked since (2) satisfies (A1)-(A2) for a standard generalized risk measure. We show the “only if” statement below. Using (A1), we have  $\Psi(X|Q) \geq \Psi(X|P)$  for all  $P \in \mathcal{Q}$ , which implies “ $\geq$ ” in (2). Define  $s_X =$

$\sup_{P \in \mathcal{Q}} \Psi(X|P)$ . As  $s_X$  is a constant random variable under every  $P \in \mathcal{Q}$ , we have  $\Psi(X|P) \leq s_X = \Psi(s_X|P)$  for all  $P \in \mathcal{Q}$ . Using (A2), we have  $\Psi(X|\mathcal{Q}) \leq \Psi(s_X|\mathcal{Q}) = s_X$ . Thus, “ $\leq$ ” in (2) follows.

(ii) For the “only if” statement, (A1) gives the “ $\geq$ ” direction of (2) and the (A3) gives the “ $\leq$ ” direction of (2). The “if” statement is straightforward to check.  $\square$

*Proof of Theorem 2.* The first statement can be directly checked. For the second statement, it suffices to show (B2+B3) $\Rightarrow$ (B1). Take  $P, Q \in \mathcal{P}$  and  $X, Y \in \mathcal{X}$  such that  $X|_P \stackrel{d}{=} Y|_Q$ , and we denote this common distribution by  $F$ . As explained above, we aim to show that  $\Psi(X|P) = \Psi(Y|Q)$ .

Let  $P' = (P + Q)/2$  which is a probability measure dominating both  $P$  and  $Q$ . Since both  $(\Omega, \mathcal{F}, P)$  and  $(\Omega, \mathcal{F}, Q)$  are atomless, so is  $(\Omega, \mathcal{F}, P')$ . Hence, there exist iid uniform  $[0, 1]$  random variables  $U$  and  $V$  under  $P'$ . Take an arbitrary  $x \in (0, 1)$  and define a probability measure  $R$  as the regular conditional probability

$$R(A) = P'(A|U = x), \quad A \in \mathcal{F}.$$

Note that  $R$  is a well-defined probability measure since the measurable space  $(\Omega, \mathcal{F})$  is standard Borel (see e.g., Theorem 5.1.9 of [Durrett \(2010\)](#)). It is clear that  $R$  is atomless since  $V$  is uniformly distributed under  $R$ . Moreover,  $R$  and  $P'$  are mutually singular. Since  $P, Q \ll P'$ , we know that  $R$  and  $P$  are mutually singular, and so are  $R$  and  $Q$ . By Remark 3.13 and Theorem 3.17 of [Shen et al. \(2019\)](#), there exists a random variable  $Z$  such that the distribution of  $Z$  is  $F$  under both  $P$  and  $R$ . Similarly, there exists a random variable  $W$  such that the distribution of  $W$  is  $F$  under both  $Q$  and  $R$ . Therefore, we obtain the chain of equalities

$$X|_P \stackrel{d}{=} Z|_P \stackrel{d}{=} Z|_R \stackrel{d}{=} W|_R \stackrel{d}{=} W|_Q \stackrel{d}{=} Y|_Q,$$

which implies

$$\Psi(X|P) = \Psi(Z|P) = \Psi(Z|R) = \Psi(W|R) = \Psi(W|Q) = \Psi(Y|Q).$$

Hence,  $\Psi$  satisfies (B1).  $\square$

*Proof of Proposition 2.* For the proof of (i), an application of Theorem 1 shows that  $\Psi$  satisfies

(A1) and (A3) if and only if

$$\Psi(X|Q) = \sup_{P \in \mathcal{Q}} \Psi(X|P), \quad (X, Q) \in \mathcal{X} \times 2^{\mathcal{P}}. \quad (8)$$

It remains to show that  $\Psi$  satisfies (B1) if and only if there exists a function  $\psi \in \Sigma$  such that  $\Psi(X|P) = \psi(F_{X|P})$  for any  $(X, P) \in \mathcal{X} \times \mathcal{P}$ . The “if” statement can be checked directly.

Now we prove the “only if” statement. For any  $F \in \mathcal{M}$ , there exists a random variable  $X \in \mathcal{X}$  and a probability measure  $P \in \mathcal{P}$  such that  $F$  is the distribution of  $X$  under  $P$ . According to (B1),  $\Psi(X|P)$  is irrelevant to the selection of  $X$  and  $P$ . Hence, we define a functional  $\psi \in \Sigma$  via

$$\psi(F) = \Psi(X|P), \quad F \in \mathcal{M}. \quad (9)$$

Combining (8) and (9), we complete the proof of (i). One can prove (ii) and (iii) similarly.  $\square$

*Proof of Theorem 3.* It is straightforward to check that the risk measure  $\Psi$  defined by (3) satisfies (A1), (A2), (B1), (C1)-(C4) and (C0). Next, we show that these properties pin down (3). Using Theorem 1, it suffices to show that, with (B2), (C1) and (C0), the mapping  $X \mapsto \Psi(X|P)$  has to be the expectation under  $P$ . It is a well-known result that a monotone, standard, law-invariant, and additive functional has to be the mean; see e.g., the proof of Lemma A.1 of Wang and Zitikis (2021). Hence, these properties are enough to pin down (3).  $\square$

*Proof of Theorem 4.* First, we show the equivalence (ii) $\Leftrightarrow$ (iii). To show the direction (ii) $\Rightarrow$ (iii), based on (B2) and (C1)-(C5), for a fixed  $P \in \mathcal{P}$ , we get, by Theorem 1, Lemma 2 and Theorem 3 of Wang et al. (2020), there exists an increasing concave function  $h_P : [0, 1] \rightarrow [0, 1]$  with  $h_P(0) = 0 = 1 - h_P(1)$  such that

$$\Phi(X|P) = \int X d(h_P \circ P), \quad (X, P) \in \mathcal{X} \times \mathcal{P}. \quad (10)$$

For every  $x \in [0, 1]$  and  $P, Q \in \mathcal{P}$ , there exist  $A, B \in \mathcal{F}$  with  $P(A) = Q(B) = x$  by Theorem 1 of Delbaen (2012). Strong law invariance of  $\Phi$  implies

$$\Phi(\mathbb{1}_A|P) = \Phi(\mathbb{1}_B|Q) = h_P(P(A)) = h_Q(Q(B)).$$

Thus,  $h_P(x) = h_Q(x)$  for all  $x \in [0, 1]$ . This shows that  $h_P$  does not depend on  $P$ , and writing  $h = h_P$ , (10) leads to (4).

To show the direction (iii) $\Rightarrow$ (ii), for fixed  $P \in \mathcal{P}$ , using Theorem 3 of Wang et al. (2020),  $\Phi(\cdot|P)$

is subadditive. Monotonicity, cash-additivity, positive homogeneity and comonotonic additivity follow from Theorems 4.88 and 4.94 of [Föllmer and Schied \(2016\)](#). Therefore,  $\Phi$  is coherent and comonotonic-additive. Strong law invariance follows from the fact that  $X|P \stackrel{d}{=} Y|Q$  implies

$$\begin{aligned} \int X d(h \circ P) &= \int_{-\infty}^0 (h \circ P(X \geq x) - 1) dx + \int_0^{\infty} h \circ P(X \geq x) dx \\ &= \int_{-\infty}^0 (h \circ Q(Y \geq x) - 1) dx + \int_0^{\infty} h \circ Q(Y \geq x) dx = \int Y d(h \circ Q). \end{aligned}$$

Next, we show (i) $\Leftrightarrow$ (iii). To show (iii) $\Rightarrow$ (i), as the other properties are straightforward to check, we only show ambiguity sensitivity. Because of cash-additivity of (4), it suffices to check it for  $X \geq 0$ . For  $X \geq 0$  and  $P, Q \in \mathcal{P}$ ,

$$\begin{aligned} \Psi(X|\lambda P + (1 - \lambda)Q) &= \int X d(h \circ (\lambda P + (1 - \lambda)Q)) \\ &= \int_0^{\infty} h(\lambda P(X \geq x) + (1 - \lambda)Q(X \geq x)) dx \\ &\geq \int_0^{\infty} (\lambda(h \circ P)(X \geq x) + (1 - \lambda)(h \circ Q)(X \geq x)) dx \\ &= \lambda \int X d(h \circ P) + (1 - \lambda) \int X d(h \circ Q) \\ &= \lambda \Psi(X|P) + (1 - \lambda) \Psi(X|Q). \end{aligned}$$

Moreover, for all  $A \in \mathcal{F}$  and  $P \in \mathcal{P}$ , we compute

$$\Psi(\mathbb{1}_A|P) = \int_0^{\infty} h(P(\mathbb{1}_A \geq x)) dx = \int_0^1 h(P(\mathbb{1}_A \geq x)) dx = \int_0^1 h(P(A)) dx = h(P(A)).$$

For all  $A \in \mathcal{F}$  and  $P, Q \in \mathcal{P}$  such that  $P(A) = Q(A)$ , we have

$$\begin{aligned} \Psi(\mathbb{1}_A|\lambda P + (1 - \lambda)Q) &= h((\lambda P + (1 - \lambda)Q)(A)) = h(P(A)) \\ &= \lambda h(P(A)) + (1 - \lambda)h(Q(A)) = \lambda \Psi(\mathbb{1}_A|P) + (1 - \lambda) \Psi(\mathbb{1}_A|Q). \end{aligned}$$

To show (i) $\Rightarrow$ (iii), by Theorem 1 and Lemma 2 of [Wang et al. \(2020\)](#), based on (B2), (C1), (C2) and (C5), for a fixed  $P \in \mathcal{P}$ , there exists an increasing function  $h_P : [0, 1] \rightarrow [0, 1]$  with  $h_P(0) = 0 = 1 - h_P(1)$  such that

$$\Psi(X|P) = \int X d(h_P \circ P), \quad (X, P) \in \mathcal{X} \times \mathcal{P}. \quad (11)$$

Also note that  $h_P(x) = \Psi(\mathbb{1}_A|P)$  for all  $P \in \mathcal{P}$  and  $A \in \mathcal{F}$  with  $P(A) = x$ . Further, note that for

any  $P \in \mathcal{P}$  and any  $t \in [0, 1]$ ,

$$\Psi(\mathbf{1}_{\{U \leq t\}} | P) = \int \mathbf{1}_{\{U \leq t\}} d(h_P \circ P) = h_P(t).$$

Lyapunov's convexity theorem (Theorem 5.5 of [Rudin \(1991\)](#)) states that the set  $R(P, Q) := \{(P(A), Q(A)) : A \in \mathcal{F}\}$  is closed and convex. Hence, as  $(0, 0), (1, 1) \in R(P, Q)$ , for every  $x \in [0, 1]$  and  $P, Q \in \mathcal{P}$ , there exists  $A \in \mathcal{F}$  with  $P(A) = Q(A) = x$ . Hence, the second condition in (B4) implies

$$h_{\lambda P + (1-\lambda)Q}(x) = \lambda h_P(x) + (1-\lambda)h_Q(x). \quad (12)$$

Assume  $P \neq Q$ . There exists  $B \in \mathcal{F}$  such that  $P(B) > Q(B)$ , which also implies  $P(B^c) < Q(B^c)$ . Therefore,  $R(P, Q)$  contains at least one point above the diagonal line and at least one point below the diagonal line. Using Lyapunov's convexity theorem again, since

$$(0, 0), (1, 1), (P(B), Q(B)), (P(B^c), Q(B^c)) \in R(P, Q),$$

for any  $\varepsilon \in (0, \frac{1}{2})$ , there exists a "rectangle" set in  $R(P, Q)$ , i.e., there exists some  $\delta' > 0$  such that

$$\{(x + \delta, x - \delta) : x \in (\varepsilon, 1 - \varepsilon), \delta \in (-\delta', \delta')\} \subseteq R(P, Q). \quad (13)$$

Hence, for any  $\delta \in (0, \delta')$ , for any  $x \in (\varepsilon, 1 - \varepsilon)$ , there exists some  $B_\delta \in \mathcal{F}$  satisfying  $P(B_\delta) = x + \delta$  and  $Q(B_\delta) = x - \delta$ . The first condition in (B4) further gives

$$\begin{aligned} h_{P/2+Q/2}(x) &= h_{P/2+Q/2}\left(\frac{P(B_\delta)}{2} + \frac{Q(B_\delta)}{2}\right) \\ &\geq \frac{1}{2}h_P(P(B_\delta)) + \frac{1}{2}h_Q(Q(B_\delta)) = \frac{1}{2}h_P(x + \delta) + \frac{1}{2}h_Q(x - \delta). \end{aligned}$$

It then follows from (12) that

$$h_Q(x) - h_Q(x - \delta) \geq h_P(x + \delta) - h_P(x),$$

which implies

$$\frac{1}{\delta}(h_Q(x) - h_Q(x - \delta)) \geq \frac{1}{\delta}(h_P(x + \delta) - h_P(x)). \quad (14)$$

By the symmetry of  $P$  and  $Q$ , for any  $x \in (\varepsilon, 1 - \varepsilon)$  and  $\delta \in (0, \delta')$  sufficiently small, we similarly have

$$\frac{1}{\delta}(h_P(x) - h_P(x - \delta)) \geq \frac{1}{\delta}(h_Q(x + \delta) - h_Q(x)). \quad (15)$$

For any  $x \in (\varepsilon, 1 - \varepsilon)$  and  $\delta \in (0, \delta')$  sufficiently small, substituting  $x + \delta$  in (15), we have

$$\frac{1}{\delta}(h_P(x + \delta) - h_P(x)) \geq \frac{1}{\delta}(h_Q(x + 2\delta) - h_Q(x + \delta)), \quad (16)$$

Combining (14) and (16), for any  $x \in (\varepsilon, 1 - \varepsilon)$  and  $\delta \in (0, \delta')$  sufficiently small, we have

$$\frac{1}{\delta}(h_Q(x) - h_Q(x - \delta)) \geq \frac{1}{\delta}(h_Q(x + 2\delta) - h_Q(x + \delta)), \quad (17)$$

which implies that  $h_Q$  is concave on  $(\varepsilon, 1 - \varepsilon)$ . As  $\varepsilon \in (0, \frac{1}{2})$  is arbitrary, we have  $h_Q$  is concave on  $(0, 1)$ . Concavity implies that  $h_Q$  is absolutely continuous on  $(0, 1)$ . Similarly,  $h_P$  is also concave and hence absolutely continuous on  $(0, 1)$ .

Note that  $h_P$  and  $h_Q$ , as increasing functions, have derivatives almost everywhere. Letting  $\delta \downarrow 0$  in (14) gives

$$\frac{d}{dx}h_Q(x) \geq \frac{d}{dx}h_P(x) \quad \text{for a.e. } x \in (0, 1).$$

By symmetry in the positions of  $P$  and  $Q$ , we have

$$\frac{d}{dx}h_P(x) \geq \frac{d}{dx}h_Q(x) \quad \text{for a.e. } x \in (0, 1),$$

which further implies that

$$\frac{d}{dx}h_P(x) = \frac{d}{dx}h_Q(x) \quad \text{for a.e. } x \in (0, 1). \quad (18)$$

Using the Newton-Leibnitz formula, for any  $x \in (0, 1]$ , we have

$$h_P(1-) - h_P(x) = \int_x^1 \frac{d}{dt}h_P(t) dt = \int_x^1 \frac{d}{dt}h_Q(t) dt = h_Q(1-) - h_Q(x). \quad (19)$$

We proceed to prove that  $h_P(1) = h_P(1-)$  for any  $P \in \mathcal{P}$ . It is clear that  $h_P(1) \geq h_P(1-)$  because  $h_P$  is increasing. Write

$$\tilde{h}_P(x) = \begin{cases} h_P(x), & x \in [0, 1); \\ h_P(1-), & x = 1. \end{cases}$$

Hence, for any  $X \in \mathcal{X}$  satisfying  $0 \leq X \leq 1$ , we have

$$\begin{aligned}
\Psi(X|P) &= \int X dh_P \circ P \\
&= \int X d(h_P(1) - \tilde{h}_P(1)) \circ P + \int X d\tilde{h}_P \circ P \\
&= (h_P(1) - \tilde{h}_P(1)) \cdot \text{ess-inf}(X|P) + \int X d\tilde{h}_P \circ P \\
&= (h_P(1) - \tilde{h}_P(1)) \cdot \text{ess-inf}(X|P) + \int_0^1 \tilde{h}_P \circ P(X \geq x) dx \\
&\leq (h_P(1) - \tilde{h}_P(1)) \cdot \text{ess-inf}(X|P) + \tilde{h}_P(1).
\end{aligned} \tag{20}$$

For any  $\lambda \in [0, 1]$ , let  $X$  be a random variable satisfying  $X|P \sim \text{Bernoulli}(\lambda)$ . On one hand, according to (20), we have

$$\Psi(X|P) \leq \tilde{h}_P(1). \tag{21}$$

On the other hand, we define two probability measures:  $Q(\cdot) = P(\cdot|X = 0)$  and  $R(\cdot) = P(\cdot|X = 1)$ . In fact, we have  $P = (1 - \lambda)Q + \lambda R$ . Hence,  $Q$  and  $R$  are mutual singular with  $Q(X = 0) = 1$  and  $R(X = 1) = 1$ . Hence,  $\Psi(X|Q) = 0$  and  $\Psi(X|R) = 1$ . According to (B4), we have

$$\Psi(X|P) \geq (1 - \lambda)\Psi(X|Q) + \lambda\Psi(X|R) = \lambda. \tag{22}$$

Combining (21) and (22), we have  $\tilde{h}_P(1) \geq \lambda$  for any  $\lambda \in [0, 1]$ , which implies that  $h_P(1-) = \tilde{h}_P(1) = 1 = h_P(1)$ .

Finally, combining with (19), we have  $h_P = h_Q$  on  $(0, 1]$ . Hence, together with  $h_P(0) = h_Q(0) = 0$ , we have  $h_P = h_Q$ . This shows that  $h_P$  does not depend on  $P$ , and we write  $h = h_P$ . As  $h$  is concave on  $(0, 1)$ ,  $h(1-) = h(1)$  and  $h(0+) \geq h(0)$ , we know that  $h$  is concave on  $[0, 1]$ . Hence, we complete the proof of (4).  $\square$

*Proof of Proposition 3.* It is straightforward to check that the multi-prior expected utility in (5) satisfies (E1)-(E5). Below we will show the representation (5) from (E1)-(E5).

By (E1), to compare  $(X, P)$  with  $(Y, Q)$ , it suffices to compare the distributions  $F_{X|P}$  and  $F_{Y|Q}$  as elements of  $\mathcal{M}$ , the set of distributions on  $\mathbb{R}$ . Hence, the restriction of  $\preceq$  to  $\{(X, P) : X \in \mathcal{X}, P \in \mathcal{P}\}$  is described equivalently by a binary relation  $\preceq^*$  on  $\mathcal{M}$ , via

$$(X, P) \preceq (Y, Q) \iff F_{X|P} \preceq^* F_{Y|Q}.$$

By letting  $F_{X|P} = F$ ,  $F_{Y|P} = G$  and  $F_{X|Q} = F_{Y|Q} = H$ , we can translate (E4) into the following

property: For any  $F, G, H \in \mathcal{M}$  and  $\alpha \in (0, 1)$ , it holds

$$F \preceq^* G \iff \alpha F + (1 - \alpha)H \preceq^* \alpha G + (1 - \alpha)H;$$

this is the independence axiom of [Von Neumann and Morgenstern \(1944\)](#) on  $\preceq^*$ . Similarly, (E5) can be translated into the continuity axiom of [Von Neumann and Morgenstern \(1944\)](#) on  $\preceq^*$ . Using the Von Neumann-Morgenstern utility theorem, there exists a function  $u : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$(X, P) \preceq (Y, Q) \iff F_{X|P} \preceq^* F_{Y|Q} \iff \mathbb{E}^P[u(X)] \leq \mathbb{E}^Q[u(Y)]. \quad (23)$$

Next, we consider two general objects  $(X_1, \mathcal{Q}_1) \in \mathcal{X} \times \mathcal{S}$  and  $(X_2, \mathcal{Q}_2) \in \mathcal{X} \times \mathcal{S}$ . First, by (E2) and (E3), there exist  $Q_1^* \in \mathcal{Q}_1$  and  $Q_2^* \in \mathcal{Q}_2$  such that

$$(X_1, Q_1^*) \simeq (X_1, \mathcal{Q}) \quad \text{and} \quad (X_2, Q_2^*) \simeq (X_2, \mathcal{Q}_2)$$

Moreover, by (E2), we have  $(X_1, Q_1^*) \simeq (X_1, \mathcal{Q}_1) \preceq (X_1, Q)$  for all  $Q \in \mathcal{Q}_1$ . By using (23), we have

$$\mathbb{E}^{Q_1^*}[u(X_1)] \leq \mathbb{E}^Q[u(X_1)] \quad \text{for all } Q \in \mathcal{Q}_1.$$

Thus,  $\mathbb{E}^{Q_1^*}[u(X_1)] = \min_{Q \in \mathcal{Q}_1} \mathbb{E}^Q[u(X_1)]$ . Similarly,  $\mathbb{E}^{Q_2^*}[u(X_2)] = \min_{Q \in \mathcal{Q}_2} \mathbb{E}^Q[u(X_2)]$ . Suppose that  $\mathbb{E}^{Q_1^*}[u(X_1)] \leq \mathbb{E}^{Q_2^*}[u(X_2)]$ . By using (23) again, we have

$$(X_1, Q_1^*) \preceq (X_2, Q) \quad \text{for all } Q \in \mathcal{Q}_2.$$

Using (E3), this implies  $(X_1, \mathcal{Q}_1) \simeq (X_1, Q_1^*) \preceq (X_2, \mathcal{Q}_2)$ . Reverting the positions of  $(X_1, \mathcal{Q}_1)$  and  $(X_2, \mathcal{Q}_2)$ , we obtain that (5) holds true.  $\square$

*Proof of Proposition 4.* Proposition 2 shows that  $\Psi$  satisfies (A1), (A3) and (B2) if and only if there exists  $\{\psi_P : P \in \mathcal{P}\} \subseteq \Sigma$  such that

$$\Psi(X|\mathcal{Q}) = \sup_{P \in \mathcal{Q}} \Psi(X|P), \quad (X, \mathcal{Q}) \in \mathcal{X} \times 2^{\mathcal{P}}$$

and

$$\Psi(X|P) = \psi_P(F_{X|P}), \quad (X, P) \in \mathcal{X} \times \mathcal{P}. \quad (24)$$

It remains to show that  $\Psi$  satisfies (B5) if and only if there exists  $\gamma : \mathcal{P} \rightarrow (-\infty, \infty]$  and  $\psi \in \Sigma$

such that

$$\Psi(X|P) = \psi_P(F_{X|P}) = \psi(F_{X|P}) - \gamma(P), \quad (X, P) \in \mathcal{X} \times \mathcal{P}. \quad (25)$$

The “if” statement can be checked directly. Now we proceed to prove the “only if” statement.

Assume that (B5) holds. For any  $X, Y, Z, W \in \mathcal{X}$  and  $P, Q \in \mathcal{P}$  satisfying  $X|P \stackrel{d}{=} Y|Q$  and  $Z|P \stackrel{d}{=} W|Q$ , according to (24) we have  $\psi_P(F_{X|P}) - \psi_Q(F_{Y|Q}) = \psi_P(F_{Z|P}) - \psi_Q(F_{W|Q})$ . Write  $F_1 = F_{X|P} = F_{Y|Q}$  and  $F_2 = F_{Z|P} = F_{W|Q}$ . Hence, for any  $F_1, F_2 \in \mathcal{M}$  and  $P, Q \in \mathcal{P}$ , we have

$$\psi_P(F_1) - \psi_Q(F_1) = \psi_P(F_2) - \psi_Q(F_2),$$

which means  $\psi_P(F) - \psi_Q(F)$  is a constant for any  $F \in \mathcal{M}$ . That is, there exists a functional  $g : \mathcal{P} \times \mathcal{P} \rightarrow \mathbb{R}$  such that

$$\psi_P(F) - \psi_Q(F) = g(P, Q), \quad P, Q \in \mathcal{P}, \quad F \in \mathcal{M}.$$

Fix  $P \in \mathcal{P}$  and set  $Q = \delta_0$ , the Dirac measure at zero. Then

$$\psi_P(F) = \psi_{\delta_0}(F) + g(P, \delta_0), \quad F \in \mathcal{M}.$$

Define  $\psi(\cdot) = \psi_{\delta_0}(\cdot)$  and  $\gamma(\cdot) = -g(\cdot, \delta_0)$ . We have

$$\psi_P(F) = \psi(F) - \gamma(P), \quad F \in \mathcal{M}. \quad (26)$$

Since (26) holds for any  $P \in \mathcal{P}$ ,

$$\psi_P(F) = \psi(F) - \gamma(P), \quad (P, F) \in \mathcal{P} \times \mathcal{M}.$$

For any  $X \in \mathcal{X}$  and  $P \in \mathcal{P}$ , we consider  $F = F_{X|P}$  and hence have

$$\Psi(X|P) = \psi_P(F_{X|P}) = \psi(F_{X|P}) - \gamma(P), \quad (X, P) \in \mathcal{X} \times \mathcal{P},$$

which is exactly (25). □