

1 Title:

2 Field-based and overspeed potentiated warm-ups increase clubhead speed and drive carry  
3 distance in skilled collegiate golfers

4 Authors:

5 Bliss, A<sup>1</sup>., Livingstone, H<sup>1</sup>., Tallent, J<sup>1</sup>.,

6 Corresponding Author:

7 Alex Bliss. Programme Director, MSc Applied Strength and Conditioning Science, Senior  
8 Lecturer, St Mary's University, Twickenham, UK [alex.bliss@stmarys.ac.uk](mailto:alex.bliss@stmarys.ac.uk)

9 Affiliations:

10 1: Faculty of Sport, Allied Health and Performance Sciences. St Mary's University.  
11 Twickenham, London, UK, TW1 4SX.

12

13

14

15

16

17

18

19

20

21

22

23 **Abstract:**

24 Warm-ups utilising post-activation performance enhancement (PAPE) strategies have been  
25 shown to increase clubhead speed (CHS) in golfers. However, the effectiveness of overspeed  
26 training using weighted clubs to elicit PAPE in CHS is unknown. The purpose of this  
27 investigation was to compare traditional, field-based warm-up activities with no potentiation  
28 activity (CON), against a field-based potentiated warm-up using high rate of force development  
29 bodyweight movements (BWP), and an overspeed warm-up using speed sticks (SSP) as the  
30 potentiation method. Thirteen skilled adult male golfers (handicap  $1.0 \pm 2.1$ ) completed three  
31 testing sessions, separated by seven days. The CON, BWP and SSP warm-ups were  
32 identical, except for the potentiation method. After each warm-up condition, ten shots,  
33 separated by one minute, were recorded using a doppler rader launch monitor (Trackman 4)  
34 with CHS, ball speed (BS), carry distance (CD) and total distance (TD) recorded. A repeated  
35 measures one-way ANOVA with Bonferroni *post hoc* pairwise comparisons revealed  
36 increases in CHS in the BWP ( $p= 0.004$ ) and SSP ( $p= 0.003$ ) groups against CON, with no  
37 difference between BWP and SSP. Increased CD was observed for BWP ( $p= 0.034$ ) and SWP  
38 ( $p= 0.030$ ) against CON with no differences between BWP and SSP. No differences for BS or  
39 TD were observed. Warm-ups with BWP or SSP activities should be considered if players are  
40 attempting to increase CHS or CD of drives, although utilising overspeed potentiation methods  
41 appear to confer no additional benefit to bodyweight PAPE exercises in skilled collegiate  
42 golfers.

43 **Keywords:** golf, warm-ups, overspeed, speed sticks, post activation potentiation effect

44 **Introduction:**

45 Effective warm-ups for athletic performance typically follow the sequential “Raise, Activate,  
46 Mobilise, Potentiate” (RAMP) model originally proposed by Jeffreys<sup>1</sup> where body  
47 temperature and heart rate are raised, muscles are activated and joints mobilised, before the  
48 musculature is primed or potentiated for the task about to be performed in a sequential

49 manner. Golf warm-ups that contain these elements have been shown to improve  
50 determinants of drive performance in golf including clubhead speed (CHS), driving distance  
51 and strike quality<sup>2</sup>. Conversely, warm-ups that focus on static stretching and do not adhere to  
52 the RAMP model have been demonstrated to contribute to decrements in these performance  
53 measures.<sup>3</sup> A recent review of warm-ups in golf has provided a thorough overview of the area,  
54 suggesting that to be practically viable, warm-ups should include some form of resistance  
55 exercise but with minimal equipment.<sup>4</sup> However, none of the studies systematically  
56 investigated contained golf-specific overspeed potentiation methods, or directly compared  
57 bodyweight resistance exercises and golf swing specific potentiation methods. The work of  
58 Tilley and McFarlane<sup>5</sup> did use a weighted club, but this was used at the start of the warm-up.  
59 Overspeed potentiation methods have been shown to confer increases in swing speed in  
60 sports with a similar rotational striking movements such as baseball.<sup>6,7</sup> However, there is  
61 currently no evidence on overspeed potentiation methods in golf as an acute strategy to  
62 enhance CHS. Therefore, understanding whether warm-ups containing an overspeed  
63 potentiation strategy deliver maximal performance improvements is necessary.

64 Research in this area is useful as it may offer simple methods by which to increase CHS, and  
65 subsequently drive distance. Although drive distance is underpinned by a myriad of factors,  
66 the principal component for increased drive distance is increased CHS.<sup>8</sup> For example, CHS  
67 is strongly correlated with handicap index in amateur golfers, with better players demonstrating  
68 a strong correlation with CHS.<sup>9</sup> At the elite level, long-hitting golfers are more likely to score  
69 better on par four and five holes on the PGA tour.<sup>10</sup>

70 Post activation potentiation (PAP) is a commonly used technique by strength & conditioning  
71 practitioners to acutely improve physical qualities of athletes that are required to perform  
72 forceful muscular contractions.<sup>11</sup> Traditionally, PAP is observed by evoking a muscle twitch  
73 using electrical stimulation after an intense voluntary contraction, although it has also recently  
74 been defined as a voluntary force or power enhancement after a high-intensity warm-up.<sup>12</sup>  
75 This linked, but separate phenomenon is termed the post-activation performance

76 enhancement (PAPE) effect and is thought to result from increases in muscle temperature,  
77 muscle and muscle fibre water content, and other central and peripheral mechanisms to  
78 improve muscle activation.<sup>12</sup> Previous studies in golf have shown that PAPE activities can elicit  
79 positive and transferable effects to golf driving performance and CHS. Research conducted  
80 by Read, Miller & Turner has shown that skilled golfers increased CHS by 2.25 miles per hour  
81 (mph) after completing a series of bodyweight countermovement jumps (CMJ).<sup>13</sup> However,  
82 golfers may be reluctant to perform this type of warm-up because it is not common amongst  
83 their peers or (because it is a generic athletic movement rather than a golf movement) they  
84 may not know how to.<sup>4</sup> Conversely, a study of skilled golfers undergoing professional training  
85 demonstrated that warm-ups are perceived to be beneficial for golf performance, and that over  
86 50% of players undertake air swings with a golf club as part of their preparations.<sup>14</sup>  
87 Furthermore, studies investigating changes to CHS in golf following weighted club warm-ups  
88 are lacking. Based on the research of Ehlert & Wilson<sup>4</sup>, this type of warm-up may be more  
89 attractive as it mimics the golf swing, but it does involve specialist equipment.

90 Enhancements in muscular force production from PAPE activities have been observed  
91 following dynamic, high-speed activities.<sup>12</sup> Studies from sports with similar rotational  
92 hitting/striking profiles to golf such as baseball have found that performing maximal effort  
93 swings as part of a warm-up with lighter than normal, or normally weighted bats can increase  
94 subsequent normal bat swing velocity by approximately 4%, but heavier bats confer no  
95 benefit.<sup>6</sup> Therefore, it was the purpose of this study to compare the effects of both high-rate of  
96 force development bodyweight PAPE exercise (BWP) or an overspeed warm-up using speed  
97 sticks (SSP) on golf drive performance.

## 98 **Methods**

### 99 ***Participants***

100 Thirteen skilled adult male golfers (age= 20 ± 1 yrs; height= 1.82 ± 0.08 m; body mass= 77.55  
101 ± 7.11 kg; handicap= 1.0 ± 2.1) were recruited to the study. To be included in the study,

102 participants must have been a category one handicap (5.4 or lower) or professional. Twelve  
103 participants were amateur and one was professional, who was given a handicap of zero for  
104 the purposes of the study. Participants were recruited from a research advert which was  
105 placed at a golf college in the United Kingdom (UK) and golf clubs local to the university. All  
106 participants were free from injury. Power analysis was carried out using G\*Power (v3.1.9.7) *a*  
107 *priori*, determining that with an estimated effect size of 0.6 (based on the similar work of  
108 Coughlan et al.<sup>15</sup>) and an alpha level of 0.05. 12 participants were required to achieve a power  
109 >80%. The study was conducted in accordance with the principles of the Declaration of  
110 Helsinki (2013) and ethical approval was granted by the institution's ethics committee.

### 111 ***Procedure and measurements:***

112 Participants attended all testing sessions at the same time of day, separated by one-week.  
113 Participants were instructed to avoid strenuous activity 24-h prior to assessment and to arrive  
114 in a rested condition. Participants were asked to avoid eating or drinking anything other than  
115 water at least 2-h prior to assessment, and to avoid consumption of any nutritional  
116 supplements on the day of assessment. For the golf assessment, all testing sessions were  
117 carried out in an outdoor, covered driving range in the UK in similar weather conditions. A  
118 computerised launch monitor (Trackman 4, Trackman Golf, Denmark) was used to collect shot  
119 data. Participants used their own drivers, although the same balls (Srixon Range Balls, Srixon  
120 Sports Europe, UK) were used for each participant. The launch monitor was calibrated and  
121 set to a "normalised" setting for all testing sessions to account for variables such as wind  
122 direction, ground conditions, ball quality etc. Data fields recorded were: CHS, ball speed (BS),  
123 carry distance (CD) and total distance (TD). Previous research has demonstrated that the  
124 Trackman 3e (the previous model to the 4) has a median accuracy of 0.18m/s and 0.09m/s  
125 for CHS and BS respectively.<sup>16</sup> The Trackman 4 is a newer model than the 3e and is expected  
126 to be as accurate, if not more accurate than its predecessor.<sup>17</sup> If an error occurred and the  
127 launch monitor did not record all of these fields the participant was asked to re-hit.

128 Participants undertook three separate protocols. Each was categorised by the type of warm-  
129 up. Each warm-up was identical in nature, except for the final activities which aimed to elicit a  
130 PAPE effect. Protocol one (CON) consisted of players completing the standardised warm-up  
131 (Table 1) with no potentiating activity and acted as a control. Protocol two added high rate of  
132 force development bodyweight plyometric exercises as a potentiating activity to the standard  
133 warm-up (BWP). Protocol three added overspeed training using Speed Sticks (SuperSpeed  
134 Golf, Tulsa, OK, USA) to the CON protocol to act as the potentiating activity (SSP)). The Speed  
135 Sticks were light (20% lighter than a standard men's driver), medium (10% lighter) and heavy  
136 (around standard driver weight or up to 5% heavier). After completion of the warm-up,  
137 participants would rest for one minute before hitting 10 maximum effort drives with a 60 second  
138 rest between shots in accordance with previous research.<sup>18</sup> Participants were asked to "swing  
139 as hard as possible, but with a technique that you would use when playing a real course".

#### 140 *Statistical Analysis:*

141 A statistical package (IBM SPSS Statistics, v24.0, IBM Corporation, USA) was utilised for data  
142 analysis. Descriptive statistics are presented as mean  $\pm$  standard deviation. The score for  
143 each dependent variable was taken as the mean value of all shots performed per condition  
144 after any outliers were removed in accordance with previous research.<sup>18</sup> The outlier analysis  
145 employed box-and-whisker plots to remove any mishit shots. Values outside of 1.5\* the lower  
146 bound for each dependent variable were removed. A one-way repeated measures analysis of  
147 variance (ANOVA) with partial eta squared ( $\eta_p^2$ ) effect size calculations was conducted to  
148 compare means of the three groups for each dependent variable. Data were checked for  
149 sphericity using Mauchly's test, with any violations adjusted using the Greenhouse-Geiser  
150 correction. Effect sizes were classified as  $\geq 0.1$ = small;  $\geq 0.3$ = medium;  $\geq 0.5$ = large.<sup>18</sup> Where  
151 significant effects were observed, Bonferroni *post hoc* comparisons were used. An alpha level  
152 of  $\leq 0.05$  was used for significance.

#### 153 **Results:**

154 From 390 shots performed, the outlier removal process disregarded 24 shots. All participants  
155 had at least seven data points for each dependent variable for each protocol following outlier  
156 removal. Descriptive data are displayed in Table 2. ANOVA revealed significant, large effects  
157 of warm-up on CHS ( $F_{(2,24)} = 14.822$ .  $p = <0.001$ .  $\eta_p^2 = 0.553$ ) and significant medium effects  
158 on CD ( $F_{(2,24)} = 5.569$ .  $p = 0.01$ .  $\eta_p^2 = 0.317$ ). Bonferroni *post hoc* comparisons revealed, when  
159 compared to the CON condition, increased CHS in the BWP ( $110.1 \pm 5.5$  vs  $111.6 \pm 5.1$  mph.  
160  $p = 0.004$ .  $ES = 0.28$ ) and SSP conditions ( $110.1 \pm 5.5$  mph vs  $111.6 \pm 5.2$ .  $p = 0.003$ .  $ES =$   
161  $0.28$ ), but no difference between BWP and SSP ( $p = 1.000$ ). Compared to the CON protocol,  
162 increased CD was observed for the BWP ( $261.5 \pm 16.4$  yards vs  $267.1 \pm 14.2$ .  $p = 0.034$ .  $ES =$   
163  $0.37$ ) and SSP conditions ( $261.5 \pm 16.4$  vs  $268.2 \pm 16.0$  yards.  $p = 0.030$ .  $ES = 0.41$ ), but no  
164 difference between BWP and SSP ( $p = 1.000$ ). No other significant effects were found for BS  
165 or TD (all  $p > 0.05$ ). The dependent variables with significant effects are displayed in Figure 1.

## 166 **Discussion:**

167 The aim of this study was to investigate three identical warm-up protocols that varied in  
168 potentiation method only and their effects on golf driving performance in skilled adult golfers.  
169 The novel element of this study is the use of overspeed training utilising weighted clubs as a  
170 potentiation method in a warm-up. The study found that utilising BWP or SSP methods can  
171 acutely increase CHS and CD in skilled golfers, but do not influence BS or TD.

172 Undertaking a warm-up prior to golf performance, despite recent evidence, appears to be a  
173 behaviour that is perceived as important by skilled professional golfers<sup>14</sup> but is not well  
174 established in amateur golfers.<sup>4</sup> This is surprising given that much recent research has  
175 demonstrated the positive benefits of doing so.<sup>2,5,15</sup> A key finding from this study is that  
176 undertaking maximal effort activity using BWP or SSP to finish the warm-up appears to cause  
177 a PAPE effect and creates increases in CHS and CD when compared to a warm-up with no  
178 potentiation activity. However, it also appears that there are no differences between the  
179 increase if the potentiating activity is generic (BWP) or sport-specific (SSP). This finding is  
180 similar to that of Langdown et al.<sup>2</sup> who reported that even though both conditions were greater

181 than the control group, there were no differences in any of the five drive metrics (BS, launch  
182 angle, total spin, dispersion, CD) monitored between their dynamic warm-up and resistance  
183 band-warm-up, with the exception of launch angle which showed a larger reduction in the  
184 dynamic group. Interestingly, while Langdown et al.<sup>2</sup> did not measure CHS (they report an  
185 increase in BS), they showed no difference in CD, but increases were found in this study. This  
186 may be explained by impact conditions (spin rates, launch angles etc.) or by the high-intensity,  
187 maximal effort potentiation activities utilised in this study in comparison to the multiple  
188 repetition or duration-based dynamic and banded activities undertaken in the work of  
189 Langdown et al.<sup>2</sup> To substantiate this contention, Read et al.<sup>13</sup> reported an increase in CHS  
190 when using CMJs to potentiate, with their increase (2.2% equating to 2.25 mph) greater than  
191 that reported here (1.4% equating to 1.50 mph) in the BWP group. While both increases were  
192 significant, the participants in this study had higher CHS ( $110.1 \pm 5.5$  mph in the CON no  
193 potentiation condition) than those in the Read study ( $106.9 \pm 6.6$  mph).<sup>13</sup> It may be that as the  
194 participant's "normal" CHS increases, that the effect size of a BWP warm-up becomes smaller.  
195 Future research could address this by comparing warm-ups designed to elicit a PAPE effect  
196 in high and low CHS participants.

197 Overspeed training is a practice that has garnered attention in other rotational striking sports  
198 such as baseball,<sup>6,7</sup> but has seen a recent revival in golf, through the use of weighted golf  
199 clubs. However, despite these implements being widely used across all levels of golf including  
200 the elite level, there is currently no peer-reviewed evidence to support their use. In baseball,  
201 warm-ups utilising maximal effort swings with lightweight or normally weighted bats elicited  
202 improvements (8.3% and 4.8% increases, respectively) in bat swing speed against using  
203 heavily weighted bats.<sup>6</sup> In a separate warm-up study utilising a range of weighted baseball  
204 bats from very light to very heavy as potentiation methods, bats within 10% of the weight of a  
205 normal bat produced the greatest swing speeds.<sup>7</sup>

206 A limitation of this study is that, even though the participants were accustomed to regular  
207 physical activity and we would not expect an order effect, the warm-up conditions were not



208 randomised. Additionally, assessment of muscular recruitment pattern or activity (via  
209 electromyography) or force production (via force platform) was not conducted. Therefore, the  
210 mechanism by which the improvements in CHS and CD can only be speculatively attributed  
211 to a PAPE effect. Future research should investigate how kinetic and kinematic factors that  
212 underpin CHS or CD are enhanced as a result of a RAMP warm-up.

213 Although CHS and CD were enhanced in both BWP and SSP conditions, no other dependent  
214 variables demonstrated an improvement. This finding likely demonstrates that increases in  
215 CHS, while a major determinant of drive distance, is not the only factor that underpins drive  
216 performance. Launch angles (vertical and horizontal), spin rates, and centredness of strike on  
217 the clubface are also key factors that underpin early ball flight characteristics and ultimately  
218 TD.<sup>20</sup> Furthermore, Parker, Hellstrom, and Ollson<sup>21</sup> demonstrated that individual swing  
219 techniques are a crucial aspect of CHS in males and females of comparable handicap and  
220 age to those in this study, although CD was less influenced by individual variance in technique.  
221 It was also suggested by that the factors that underpin CHS and CD are not transferable in  
222 males and females.<sup>21</sup> In this regard, kinetic and kinematic variables relating to individual swing  
223 technique were not collected during the testing protocols and are limitations of this study.  
224 Further, it was conducted in a male only cohort and as such the findings should not be  
225 considered generalisable to female golfers. Future research should investigate whether there  
226 are kinetic and kinematic alterations to swing technique as a result of BWP or SSP activities  
227 in addition to monitoring drive performance.

228 Lastly, it is acknowledged that there were large interindividual differences in response to the  
229 BWP and SSP warm-up conditions. As an extreme example, one participant experienced a  
230 20 yard increase in CD in the SSP condition vs CON, as where another saw a decrease of 9  
231 yards when using a SSP warm-up versus no potentiating activity. This variation in response  
232 to warm-ups aiming to elicit a PAPE effect has been previously reported. These findings are  
233 similar to those of Langdown et al<sup>2</sup> who stated, that even though all participants in their study  
234 (and this study) were category 1, skilled players, there was considerable variability in response

235 to warm-up conditions. Additionally, a study by Till & Cooke<sup>22</sup> showed a variance of 15.3%  
236 between individual responses to PAP activities on sprint and jump performance in academy  
237 footballers. The authors stated that athletes with greater muscular strength and high training  
238 exposure had greater individual responses to PAP interventions.<sup>22</sup> Furthermore, athletes with  
239 greater training experience have greater responses to PAP due to physiological make up of  
240 muscle fibres and motor units.<sup>23</sup> Athletes with limited or no training experience have reduced  
241 responses to potentiating activity<sup>23</sup> and lack of training experience or fitness levels is also  
242 shown to inhibit potentiating effects.<sup>24</sup> Therefore, it is likely that the participant's strength  
243 characteristics will influence how they respond to RAMP based warm-ups and golfers with  
244 greater physical training experience may experience the most benefit. Limitations of this study  
245 were that strength characteristics of the participants were not measured and internal load was  
246 not monitored and therefore whether the individual responses to the BWP and SSP warm-up  
247 conditions could be attributed to strength levels is unknown. Future research in this area  
248 should collect field or laboratory measures of the participants' force generating capabilities or  
249 internal load (through heart rate or rating of perceived exertion as examples) to provide useful  
250 information that may support or help to explain the variations in drive performance between  
251 participants.

## 252 **Conclusions:**

253 A warm-up that follows the RAMP protocol and contains either BWP or SSP activities elicit  
254 improvements in CHS and CD in skilled amateur male golfers. However, there were no  
255 differences between using BWP or SSP and therefore the type of potentiation activity at the  
256 end of a warm-up appears to be comparable. It is important that potentiation activities are  
257 performed at maximum effort. However, BWP and SSP warm-ups did not improve BS or TD  
258 and therefore the other kinetic and kinematic determinants of drive performance such as  
259 centredness of strike, launch angle, and spin rate need to be maintained when attempting to  
260 increase CHS and CD. Golfers can acutely increase CHS or CD through a physical warm-up  
261 if they perform BWP or SSP activities. This increase could support training or competition play

262 and may help golfers improve their drive performance on the opening hole, which will acutely  
263 improve players' scoring potential. However, it is unknown how long these performance  
264 benefits will last and future research which studies the effects of a BWP or SSP warm- up over  
265 a longer playing duration than the opening drive is warranted.

#### 266 **Conflict of Interest:**

267 The authors declare no conflict of interest

#### 268 **Acknowledgements:**

269 The authors would like to thank The Golf College for their support of the project and the  
270 participants for providing their time and efforts undertaking the study.

#### 271 **References:**

- 272 1. Jeffreys, I. Warm-up revisited: The ramp method of optimizing warm-ups. *Professional*  
273 *Strength and Conditioning* 2007. 6, 15-19.
- 274 2. Langdown B, Wells J, Graham S, et al. Acute effects of different warm-up protocols on  
275 highly skilled golfers' drive performance. *J Sports Sci* 2019. 37(6): 656-664
- 276 3. Gergley J. Acute effects of passive static stretching during warm-up on driver clubhead  
277 speed, distance, accuracy, and consistent ball contact in young male competitive  
278 golfers. *J Strength Cond Res* 2009., 23(3), 863-867.
- 279 4. Ehlert A., & Wilson P. A systematic review of golf warm-ups: behaviors, injury, and  
280 performance. *J Strength Cond Res* 2019. 33(12): 3444-3462
- 281 5. Tilley N, & McFarlane A. Effects of different warm-up programs on golf performance in  
282 elite male golfers. *Int J Sports Phys Ther.* 2012. 7(4): 388-395
- 283 6. Montoya B, Brown L, Coburn J, et al. Effect of warm-up with different weighted bats  
284 on normal baseball bat velocity. *J Strength Cond Res* 2009. 23(5): 1566-1569
- 285 7. DeRenne C, Ho K, Hetzler R et al. Effects of warm up with various weighted  
286 implements on baseball bat swing velocity. *J App Sport Sci Res* 1992. 6(4): 214-218

- 287 8. Hume P, Keogh J, Reid D. The role of biomechanics in maximising distance and  
288 accuracy of golf shots. *Sports Med* 2005. 35(5): 429-449
- 289 9. Fradkin A, Sherman C, and Finch C. How well does golf club headspeed correlate with  
290 golf handicaps? *J Sci Med Sport* 2004 7: 465–472
- 291 10. Hellström J, Nilsson J, & Isberg L. Drive for dough. PGA Tour golfers' tee shot  
292 functional accuracy, distance and hole score. *J Sports Sci* 2014. 32(5): 462–9.
- 293 11. Evetovich T, Conley D, & McCawley P. Postactivation potentiation enhances upper-  
294 and lower-body athletic performance in collegiate male and female athletes. *J Strength*  
295 *Cond Res* 2015. 29(2): 336-342.
- 296 12. Blazeovich A, & Babault N. Post-activation potentiation versus post-activation  
297 performance enhancement in humans: historical perspective, underlying mechanisms,  
298 and current issues. *Front. Phys.* 2019. doi: 10.3389/fphys.2019.01359
- 299 13. Read P, Miller S, & Turner A. The effects of postactivation potentiation on golf club  
300 head speed. *J Strength Cond Res* 2013. 27(6): 1579-1582.
- 301 14. Wells J, & Langdown, B. Sports science for golf: a survey of high-skilled pgolfers'  
302 "perceptions" and "practices". *Journal of Sports Sciences* 2020. 38(8): 918-927
- 303 15. Coughlan D, Taylor M, & Jackson J. The impact of warm-up on youth golfer clubhead  
304 speed and self-reported shot quality. *Int J Sports Phys Ther* 2018. 13(5): 828-834
- 305 16. Leach R, Forrester S, Mears A, Roberts J. How valid and accurate are measurements  
306 of golf impact parameters obtained using commercially available radar and  
307 stereoscopic optical launch monitors? *Measurement* 2017 112:125–136.
- 308 17. Turner J, Forrester S, Mears A, Roberts J. Reliability of repeat golf club testing  
309 sessions with modified club moment of inertia. *Sports Engineering*. 23(8). 2020.
- 310 18. Bliss A, McCulloch H, & Maxwell N. The effects of an eight-week plyometric training  
311 program on golf swing performance characteristics in skilled adolescent golfers. *Int J*  
312 *Golf Sci* 2015. 4(2), 120-135
- 313 19. Cohen J. Statistical power analysis for the behavioral sciences (2<sup>nd</sup> ed.). 1988 New  
314 Jersey: Lawrence Erlbaum.

315 20. Sweeney M, Mills P, Alderson J, et al. The influence of club-head kinematics on early  
316 ball flight characteristics in the golf drive. *Sports Biomech* 2013. 12(3): 247-258

317 21. Parker J, Hellstrom J, Olsson C. Differences in kinematics and driver performance in  
318 elite female and male golfers. *Sports Biomech* 2019.  
319 DOI:10.1080/14763141.2019.1683221

320 22. Till K, & Cooke C. The effects of postactivation potentiation on sprint and jump  
321 performance of male academy soccer players. *J Strength Cond Res* 2009. 23(7): 1960-  
322 1967

323 23. Rixon K, Lamont H, & Bemben M. Influence of type of muscle contraction, gender, and  
324 lifting experience on postactivation potentiation performance. *J Strength Cond Res*  
325 2007. 21(2): 500-505

326 24. Chiu L, Fry A, Schilling B, et al. Neuromuscular fatigue and potentiation following two  
327 successive high intensity resistance exercise sessions. *Eur J App Phys* 2004. 92: 385-  
328 392

329

330

331

332

333

334

335

336

337

338

339 Table 1: Standardised sequential RAMP-based warm-up protocol

---

**Raise**

Skipping (2 minutes)

**Activation and mobilisation**

Leg swings x 10 ES

Resistance band shoulder external rotations 10 ES x 2

Single leg kneeling kickbacks x 10 ES

Lunges with rotations x 10 ES

Overhead squats with golf club x 12

**Golf Swing Specific**

Sand wedge pitch shots x 3

Sand wedge full shots x 3

7 iron full shots x 2

Driver full shots x 2

**Potentiation**

---

<b>Condition</b>		
<b>CON</b>	<b>BWP</b>	<b>SSP</b>
None	CMJ 10 reps x 3	SSS Light Dominant Side x 10
	Plyometric Press Ups 10 reps x 2	SSS Light Non-Dominant Side x 10
		SSS Medium Dominant Side x 10
		SSS Heavy Dominant Side x 10

---

340 ES = Each side. CMJ= Countermovement Jump. SSS= Super Speed Stick. Reps =  
 341 repetitions

342

343

344

345

346

347

348

349 *Table 2: Mean ( $\pm$  SD) values for drive variables across warm-up conditions*

	<b>CON</b>	<b>BWP</b>	<b>SSP</b>
<b>CHS (mph)</b>	110.1 $\pm$ 5.5	111.6 $\pm$ 5.1*	111.6 $\pm$ 5.2*
<b>BS (mph)</b>	160.5 $\pm$ 8.0	161.8 $\pm$ 7.2	161.9 $\pm$ 7.9
<b>CD (yards)</b>	261.5 $\pm$ 16.4	267.1 $\pm$ 14.3*	268.2 $\pm$ 16.0*
<b>TD (yards)</b>	285.1 $\pm$ 17.8	287.7 $\pm$ 15.7	289.2 $\pm$ 18.0

350 \*= statistically significant ( $p \leq 0.05$ ) increase vs CON condition

351

352

353

354

355

356

357

358

359

360

361


362

363

364

365

366



367

368 **Figure 1:** Mean CHS (top) and CD (bottom) for all warm-up conditions. Error bars represent  
 369 SD. Grey lines represent individual responses. \*= significant difference ( $p < 0.05$ )

370