



On the Minkowski problem for p -harmonic measures

Murat Akman¹ · Shirsho Mukherjee¹

Received: 14 December 2023 / Accepted: 22 October 2024 / Published online: 16 December 2024
© Crown 2024

Abstract

We study the Minkowski problem corresponding to the p -harmonic measures and obtain results previously known for harmonic measures due to Jerison (Invent Math 105(2):375–400, 1991). We show that a class of Borel measures on spheres can be prescribed by p -harmonic measures on convex domains.

Mathematics Subject Classification 31B05 · 35J08 · 35J25 · 42B37 · 42B25 · 42B99

Contents

1	Introduction	1
2	Notations and preliminaries	3
2.1	Convex domains	4
2.2	Convex geometry	7
2.3	Hausdorff distance and approximations	11
2.4	p -harmonic functions and measures	12
2.5	Minkowski problem	14
3	Existence for p -harmonic Minkowski problem	15
3.1	First variation of density of p -harmonic measures	16
3.2	Convergence of p -harmonic measures	27
3.3	Proof of existence	32
	References	37

1 Introduction

One of the most fundamental problems in the geometric analysis of convex bodies is the Minkowski problem. The original form of the problem seeks to find a convex polyhedron that has a specified set of vectors and real numbers as its normals and surface areas of its faces under stipulated conditions. It can be reformulated more generally as seeking to find

Communicated by O. Savin.

Both authors have been supported by the EPSRC New Investigator award [grant number EP/W001586/1].

✉ Shirsho Mukherjee
shirsho.mukherjee@essex.ac.uk ; m.shirsho@gmail.com

Murat Akman
murat.akman@essex.ac.uk

¹ Department of Mathematical Sciences, University of Essex, Colchester CO4 3SQ, UK

a convex domain that induces a specified measure on a sphere as the image of its surface measure mapped by its normals. In this form, the original problem is identical to the case of having a discrete measure on a sphere. Similar related problems are to show the uniqueness of such domains up to translations and to find convex domains with smooth boundaries when the regularity of the Gaussian curvature of its boundary as a function of its normal is specified.

The literature goes back to Minkowski [41, 42] who proved the existence and uniqueness of a convex polyhedron with specified discrete spherical measure if and only if the centroid is at the origin and support of the measure is not concentrated on an equator (or great sub-spheres). The problem for measures having continuous density was also illustrated by Minkowski and later solved for general measures by Alexandrov [5, 7] and Fenchel-Jessen [19]. The problem corresponding to convex domains with smooth boundaries involves establishing regularity of solutions for Monge-Ampère type equations on the sphere, which is rather difficult and has involved many significant works including Lewy [38], Pogorelov [44], Nirenberg [43], Cheng-Yau [13], Caffarelli [10–12], etc. Furthermore, the harmonic measures on the boundary of convex domains being equivalent to the surface measures (see Dahlberg [16]), an analogous Minkowski problem with surface measures being replaced by harmonic measures, has been studied by Jerison [26]. The objective of the present paper is to study the Minkowski problem corresponding to p -harmonic measures for $1 < p < \infty$, which, to our knowledge, remained open.

For $n \geq 3$, we consider convex domains $\Omega \subset \mathbb{R}^n$ and the Gauss map $\mathbf{g}_\Omega : \partial\Omega \rightarrow \mathbb{S}^{n-1}$ where $\mathbf{g}_\Omega(x)$ is the outer unit normal at $x \in \partial\Omega$, which is defined almost everywhere on the boundary as convexity implies that $\partial\Omega$ is locally Lipschitz. We consider a positive finite Borel measure μ on \mathbb{S}^{n-1} that satisfies the following necessary conditions:

$$\begin{aligned} (i) \quad & \int_{\mathbb{S}^{n-1}} |\langle \zeta, \xi \rangle| d\mu(\xi) > 0, \quad \forall \zeta \in \mathbb{S}^{n-1}, \\ (ii) \quad & \int_{\mathbb{S}^{n-1}} \xi d\mu(\xi) = 0. \end{aligned} \tag{1.1}$$

The original Minkowski problem seeks the existence up to a translation of a unique compact convex domain Ω with non-empty interior such that $(\mathbf{g}_\Omega)_* \mathcal{H}^{n-1} \llcorner_{\partial\Omega} = \mu$, where $(\mathbf{g}_\Omega)_* \mathcal{H}^{n-1}$ is the pushforward measure on \mathbb{S}^{n-1} and μ satisfies (i) and (ii) of the above. We show the existence of the domain corresponding to \mathcal{H}^{n-1} being replaced by p -harmonic measures.

Given any convex set $K \subset \mathbb{R}^n$, the p -harmonic measure ω_p supported on ∂K associated to a function $u = u_K$ is given by

$$\omega_p(E) = \int_{\partial K \cap E} |\nabla u|^{p-1} d\mathcal{H}^{n-1},$$

for any measurable $E \subseteq \mathbb{R}^n$, where u vanish on ∂K and is non-negative and p -harmonic in a neighborhood of it. This includes K being open or the closure of an open convex domain as well as the case of K being a convex set of empty interior. It also includes the case of u having a possible blow-up or being arbitrarily large somewhere in the interior of the domain away from the boundary, that resembles the behavior of Green’s function for the linear case. The p -harmonic measure has been studied by Lewis [36], Lewis-Nyström-Vogel [33], see also [3, 25, 29, 30], etc. The Gauss map \mathbf{g}_K , which is defined almost everywhere from ∂K to \mathbb{S}^{n-1} when K is convex and of non-empty interior with $\mathbf{g}_K(x) \in \mathbb{S}^{n-1}$ being the outer unit normal for \mathcal{H}^{n-1} -a.e. $x \in \partial K$, can be generalized to be defined as a set-valued map where $\mathbf{g}_K(x)$ constitutes the set of all normals at any $x \in \partial K$ as in [26]; the notion of push forward $(\mathbf{g}_K)_*$ for measures can be accordingly generalized, which coincides with the case

when \mathbf{g}_K is single-valued. This is necessary to include convex sets of empty interior and lower dimensional sets. Further details are in Sect. 2.

The main result of this paper is the following.

Theorem 1.2 *Given a finite regular Borel measure μ on \mathbb{S}^{n-1} satisfying conditions (1.1), there exists a bounded convex domain Ω with non-empty interior such that $(\mathbf{g}_\Omega)_*\omega_p \llcorner_{\partial\Omega} = \mu$, where \mathbf{g}_Ω is the \mathcal{H}^{n-1} -a.e. defined Gauss map and ω_p is any p -harmonic measure on $\partial\Omega$ for $1 < p < \infty$.*

Many further problems of similar character as the Minkowski problem have been studied with different measures as replacement of the surface measures, e.g. capacity measures, curvature measures, etc. We refer to [8, 14, 22–24, 40], and references therein. The proof of the existence of domains involves variational methods with respect to the Minkowski sum of convex domains $\Omega + t\Omega'$ as in Jerison [26, 27] and Colesanti et al. [15] (see also [1, 2, 4]); adaptations of such methods form the key elements of our proof of existence. The problem of the existence of domains is converted to a constrained minimization problem of functions by virtue of the dualistic relation between convex domains to their support functions and positive continuous functions to their Wulff shapes, see Sect. 2 for details. However, contrary to surface measures, harmonic measures, capacity measures, and others, the p -harmonic measures are not uniquely defined on the boundary of convex domains as their definition in general involves the choice of a p -harmonic function. Nevertheless, we show that the arbitrariness of the choice can be maintained as the characteristics of being p -harmonic at the boundary are similar for all p -harmonic measures and they admit natural approximants that can be uniquely defined for the purpose of establishing existence via limiting argument.

The paper is organized as follows. In Sect. 2, we develop our notations and provide a self-contained account of preliminaries on convex domains, geometric features of their boundaries, and approximations. We also enlist the definitions and known results on p -harmonic functions, p -harmonic measures, and Minkowski problems. In Sect. 3, we compute the first variation of densities of p -harmonic measures defined on smooth convex domains of the form $\Omega + t\Omega'$. Then we provide an account of weak convergence of p -harmonic measures which are used to construct approximants of the measures. The properties of the first variation are used in the variational methods for the existence of the domains for discrete measures, which are further used as approximations for the existence of the domains for general measures in the proof of Theorem 1.2 at the end.

The uniqueness up to translation and regularity of the domains are nontrivial and have not been addressed in this paper. These objectives are to be pursued in the future.

2 Notations and preliminaries

In this section, we introduce some notations and review some well-known properties of convex domains, p -harmonic measures and Minkowski problems.

For $n \geq 1$, we denote points in \mathbb{R}^n as $x = (x_1, \dots, x_n)$ so that if $\{e_1, \dots, e_n\}$ is the standard basis then $(x_1, \dots, x_n) = \sum_{i=1}^n x_i e_i$. The standard inner product on \mathbb{R}^n shall be denoted by $\langle \cdot, \cdot \rangle$ and the Euclidean norm by $|\cdot|$. For functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $F = (f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$, the gradient defined by $\nabla f = \sum_{i=1}^n (\partial_i f) e_i$, the Jacobian defined by $DF = \sum_{i=1}^m \sum_{j=1}^n (\partial_i f_j) e_i \otimes e_j$ and the Hessian defined by $D^2 f = D(\nabla f) = \sum_{i,j=1}^n (\partial_{i,j}^2 f) e_i \otimes e_j$, are as usual.

In general, hyperplanes in \mathbb{R}^n shall be denoted as

$$H_{y,\alpha} := \{x \in \mathbb{R}^n : \langle x, y \rangle = \alpha\}, \tag{2.1}$$

for some $y \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$; the half-spaces corresponding to $\langle x, y \rangle < \alpha$ and $\langle x, y \rangle > \alpha$ shall be respectively denoted as $H_{y,\alpha}^-$ and $H_{y,\alpha}^+$. Thus, we have $H_{y,\alpha}^- = H_{-y,-\alpha}^+$, $H_{y,\alpha}^+ = H_{-y,-\alpha}^-$ and $H_{y,\alpha} = H_{-y,-\alpha}$, for any $y \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$. In particular, $H_{e_n,0}^+$ is denoted by \mathbb{R}_+^n .

Given a measure space (X, μ) and a map $f : X \rightarrow Y$, the pushforward measure $f_*\mu$ on Y is defined on any measurable subset $E \subseteq Y$ as

$$(f_*\mu)(E) = \mu(f^{-1}(E)),$$

which is absolutely continuous with respect to μ and in the infinitesimal form, for any function $g : Y \rightarrow \mathbb{R}$ it is written as $g d(f_*\mu) = (g \circ f)d\mu$. For any Borel set $E \subseteq \mathbb{R}^n$, the k -dimensional Hausdorff measure on \mathbb{R}^n is defined by

$$\mathcal{H}^k(E) = \lim_{\delta \rightarrow 0^+} \inf \left\{ \sum_j r_j^k : E \subset \bigcup_j B_{r_j}(x_j), r_j \leq \delta \right\},$$

where $B_r(x) = \{y \in \mathbb{R}^n : |y - x| < r\}$ is the standard metric ball of radius r centered at $x \in \mathbb{R}^n$. The distance of E from a point $y \in \mathbb{R}^n$ is defined by $\text{dist}(y, E) = \inf\{|x - y| : x \in E\}$. The boundary of the unit sphere $B_1(0)$ of \mathbb{R}^n shall be denoted as \mathbb{S}^{n-1} . We have the transformation to polar coordinates $\mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty) \times \mathbb{S}^{n-1}$ as $x \mapsto (|x|, x/|x|)$ and its inverse $(r, \theta) \mapsto r\theta$. For any $F \in L^1(\mathbb{R}^n)$ we have

$$\int_{\mathbb{R}^n} F(x)dx = \omega_n \int_0^\infty \left[\int_{\mathbb{S}^{n-1}} F(r\theta)r^{n-1}d\theta \right] dr, \tag{2.2}$$

where the Lebesgue measure \mathcal{L}^n of \mathbb{R}^n and the uniform measure \mathcal{H}^{n-1} on \mathbb{S}^{n-1} are abbreviated as dx and $d\theta$ in infinitesimal form and $\omega_n = 2\pi^{n/2}/\Gamma(n/2)$ is the surface area of \mathbb{S}^{n-1} .

2.1 Convex domains

Here we shall denote $\Omega \subset \mathbb{R}^n$ as a convex domain, i.e. open and connected non-empty convex subset and $K \subset \mathbb{R}^n$ as a convex body, i.e. closure of a bounded convex domain, hence a compact convex subset with non-empty interior. We highlight some essential notions and properties related to convex sets and functions. For further details on the theory of convex bodies, we refer to Schneider [45] and Gardner [20].

For $A \subset \mathbb{R}^n$, a function $f : A \rightarrow \mathbb{R}$ is convex iff the epigraph $K = \{(x, c) \in A \times \mathbb{R} : f(x) \leq c\}$ is convex, which implies f is continuous and locally Lipschitz; we denote $\mathcal{K}(A)$ as the set of all convex functions $f : A \rightarrow \mathbb{R}$ and hence, $\mathcal{K}(A) \subset C_{\text{loc}}^{0,1}(A)$. We recall that $f \in C^1(\Omega) \cap \mathcal{K}(\Omega)$ iff $\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq 0$ for all $x, y \in \Omega$ and $f \in C^2(\Omega) \cap \mathcal{K}(\Omega)$ iff $D^2 f \geq 0$. In general, for $f \in \mathcal{K}(\mathbb{R}^n)$, the *sub-differential* of f at $x \in \mathbb{R}^n$, given by

$$\partial f(x) = \{v \in \mathbb{R}^n : f(y) \geq f(x) + \langle v, y - x \rangle\}, \tag{2.3}$$

is a convex subset and f is differentiable at x iff $\partial f(x)$ is a singleton. An important class of convex functions, is the class of *sub-linear functions* that are *sub-additive* and *homogeneous* of degree 1; in other words, they are defined by functions $h : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying

$$h(x + y) \leq h(x) + h(y) \quad \text{and} \quad h(\lambda x) = \lambda h(x) \quad \forall x, y \in \mathbb{R}^n, \lambda \geq 0. \tag{2.4}$$

A closed subset $K \subset \mathbb{R}^n$ is convex iff for any $z \in \mathbb{R}^n$ there exists a unique projection $p_K(z) \in K$ satisfying $|z - p_K(z)| = \text{dist}(z, K)$; in this case, we have $\text{dist}(\cdot, K) \in \mathcal{C}(\mathbb{R}^n) \cap C^1(\mathbb{R}^n \setminus K)$ and

$$\xi_K(z) := \nabla \text{dist}(z, K) = \frac{z - p_K(z)}{\text{dist}(z, K)}, \quad \forall z \in \mathbb{R}^n \setminus K. \tag{2.5}$$

Thus, for any $z \in \mathbb{R}^n \setminus K$, we have $p_K(z) \in \partial K$ and $|\nabla \text{dist}(z, K)| = 1$. For any $x \in K$ and $t \in [0, 1]$, since $t \mapsto |z - (tx + (1 - t)p_K(z))|^2$ attains minimum at $t = 0$, its slope at $t = 0$ is non-negative which leads to

$$\langle x - p_K(z), z - p_K(z) \rangle \leq 0, \quad \forall x \in K, z \in \mathbb{R}^n \setminus K. \tag{2.6}$$

Furthermore, the ray emerging from $p_K(z)$ connecting z being $R_{z,K} = \{p_K(z) + \lambda \xi_K(z) : \lambda \geq 0\}$, we note that $p_K(z') = p_K(z)$ and $\xi_K(z') = \xi_K(z)$ for all $z' \in R_{z,K}$. The continuity of p_K also ensures that for any $x \in \partial K$ there exists $z \in \mathbb{R}^n \setminus K$ such that $p_K(z) = x$ and hence $R_{z,K} \subseteq p_K^{-1}(x)$. If ∂K is smooth enough at $x \in \partial K$ then $R_{z,K} = p_K^{-1}(x)$ and $\xi_K(z)$ coincides with the outer unit normal at $x \in \partial K$. This leads to the following.

Definition 2.7 If K is closed and convex, for any $x \in \partial K$, the vector $\xi_K(z) \in \mathbb{S}^{n-1}$ as in (2.5) is called a (outer) *normal vector* of K at x if $z \in p_K^{-1}(x)$.

For any $A \subset \mathbb{R}^n$, the *support function* $h_A : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as

$$h_A(y) = \sup \{ \langle x, y \rangle : x \in A \}. \tag{2.8}$$

Thus, we have $h_\Omega = h_{\bar{\Omega}}$ and $h_K(y) = \max_{x \in K} \langle x, y \rangle$ for a compact convex subset $K \subset \mathbb{R}^n$. Note that h_K is linear (i.e. $h_K(y) = \langle x, y \rangle$ for every $y \in \mathbb{R}^n$) iff $K = \{x\}$. Furthermore, the convexity implies $x \in K$ iff $\langle x, y \rangle \leq h_K(y)$ for every y , i.e. in terms of the subdifferential (2.3), $\partial h_K(0) = K$. The *supporting hyperplane* of K with outer normal $y \in \mathbb{R}^n \setminus \{0\}$, is given by

$$H_K(y) := H_{y, h_K(y)} = \{x \in \mathbb{R}^n : \langle x, y \rangle = h_K(y)\}, \tag{2.9}$$

so that $K \subset H_{y, h_K(y)}^- = \{x \in \mathbb{R}^n : \langle x, y \rangle \leq h_K(y)\}$ (called *supporting half-space*) and it can be shown in terms of the subdifferential (2.3) that for any $y \in \mathbb{R}^n \setminus \{0\}$ we have

$$\partial h_K(y) = K \cap H_K(y) \subset \partial K,$$

see [45]. Thus, h_K is differentiable at $y \in \mathbb{R}^n \setminus \{0\}$ with $\nabla h_K(y) = x$ iff $K \cap H_K(y) = \{x\}$; in this case $h_{K \cap H_K(y)}(\cdot) = \langle x, \cdot \rangle = \langle \nabla h_K(y), \cdot \rangle$ and K is called *strictly convex* at x and if it holds for every point then the set is called strictly convex. For such domains, since h_K is homogeneous of degree 1, hence $y \mapsto \nabla h_K(y)$ is homogeneous of degree 0, i.e.

$$\nabla h_K(\lambda y) = \nabla h_K(y), \quad \forall \lambda \geq 0. \tag{2.10}$$

In general, it is evident that support functions are sub-linear and moreover, all sub-linear functions are support functions by virtue of the following theorem, see [45, Theorem 1.7.1].

Theorem 2.11 For any sublinear function $h : \mathbb{R}^n \rightarrow \mathbb{R}$, there exists a unique convex body $K = \{x \in \mathbb{R}^n : \langle x, y \rangle \leq h(y) \forall y \in \mathbb{R}^n\}$ such that $h_K = h$.

Due to homogeneity, it suffices to define the set $K = \{x \in \mathbb{R}^n : \langle x, \xi \rangle \leq h(\xi) \forall \xi \in \mathbb{S}^{n-1}\}$ in Theorem 2.11 having h as its support function and henceforth, it suffices to consider support

functions restricted to \mathbb{S}^{n-1} for all things and purposes. Also, this implies that for any convex set K , we have

$$K = \bigcap_{\xi \in \mathbb{S}^{n-1}} \{x \in \mathbb{R}^n : \langle x, \xi \rangle \leq h_K(\xi)\}. \tag{2.12}$$

Thus, it is evident that we have $K \subseteq K'$ iff $h_K \leq h_{K'}$. Now, alongside having supporting hyperplanes with a fixed normal, one can have a supporting hyperplane at a fixed point in the boundary. These two notions are related to each other via projection. Indeed, from (2.6), we note that $h_K(z - p_K(z)) = \langle p_K(z), z - p_K(z) \rangle$ for any $z \in \mathbb{R}^n \setminus K$ and hence, we have

$$h_K(\xi_K(z)) = \langle p_K(z), \xi_K(z) \rangle, \tag{2.13}$$

with $\xi_K(z)$ is as in (2.5). Therefore, for any $x \in \partial K$, there exists a supporting hyperplane of K at x given by

$$H_K[x] = \{x' \in \mathbb{R}^n : \langle x' - p_K(z), z - p_K(z) \rangle = 0\}, \quad \text{with } z \in p_K^{-1}(x); \tag{2.14}$$

which is related to (2.9) by $H_K[x] = H_K(z - p_K(z))$ from (2.13). Thus, from homogeneity of support functions, $H_K[x] = H_K(\xi_K(z))$ for any $z \in p_K^{-1}(x)$ where $\xi_K(z)$ is as in (2.5) and moreover, ∂K is smooth at $x \in \partial K$ iff $H_K[x]$ is unique with a unique outer unit normal.

We recall that, for any $E, F \subseteq \mathbb{R}^n$, the *Minkowski sum* is defined by

$$E + F = \{x + y : x \in E, y \in F\}; \tag{2.15}$$

also, we shall denote by $cE = \{cy : y \in E\}$ for any $c \in \mathbb{R}$. It is not hard to check that if E, F are convex then so is $E + F$ and αE for all $\alpha \geq 0$. Furthermore, for compact convex sets, the decomposition to Minkowski sums is unique, i.e. if $E + K_1 = E + K_2$ then $K_1 = K_2$. In particular, note that $B_r(x) = \{x\} + r B_1(0)$. One of the most important features of convex sets is that their support functions are Minkowski additive, i.e. for compact convex sets $E, F \subseteq \mathbb{R}^n$, we have

$$h_{\alpha E + \beta F} = \alpha h_E + \beta h_F \quad \forall \alpha, \beta \geq 0, \tag{2.16}$$

and hence, for supporting hyperplanes as in (2.9), $H_{\alpha E + \beta F}(\cdot) = \alpha H_E(\cdot) + \beta H_F(\cdot)$ holds as well. For any general $E \subseteq \mathbb{R}^n$, the *convex hull* of E , denoted by $\text{conv}(E)$, is defined as the smallest convex set (intersection of all convex sets) containing E and we have

$$\text{conv}(E + F) = \text{conv}(E) + \text{conv}(F).$$

If E is convex, then $\text{conv}(E) = E$, if E open (or closed) so is $\text{conv}(E)$. In fact, points in convex hull of a set comprise convex combinations of affinely independent points of the set; i.e. for any $x \in \text{conv}(E)$, there exists affinely independent $x_1, \dots, x_k \in E$ for $k \leq n + 1$ and $\lambda_1, \dots, \lambda_k \geq 0$ such that $x = \sum_j \lambda_j x_j$ where $\sum_j \lambda_j = 1$ (see [45, Theorem 1.1.4]).

Definition 2.17 The convex hulls of finitely many points are called *polytopes*. For any $m \leq n$, if the points $x_1, \dots, x_{m+1} \in \mathbb{R}^n$ are affinely independent then the polytope $P = \text{conv}(\{x_1, \dots, x_{m+1}\})$ is called a *m-simplex* with vertices x_1, \dots, x_{m+1} .

Thus, $\text{conv}(E)$ is the union of all m -simplices for $m \leq n$ with vertices in E . Conversely, for a convex set K , the set of points in K that cannot be written as $\lambda x + (1 - \lambda)y$ for any $x, y \in K$ is denoted as $\text{ext}(K)$ with elements called *extreme points*. Any compact convex set is the closed convex hull of its extreme points, i.e. $K = \text{conv}(\text{ext}(K))$ (Krein-Milman theorem) and hence, polytopes are the only convex bodies with finitely many extreme points. Therefore, for any polytopes P_1 and P_2 and $\alpha, \beta \geq 0$, evidently $\alpha P_1 + \beta P_2$ is also a polytope.

If H is a supporting hyperplane of a polytope $P = \text{conv}(\{x_1, \dots, x_k\})$, then we have

$$H \cap P = \text{conv}(H \cap \{x_1, \dots, x_k\}),$$

which is also a polytope of a lower dimension; such polytopes are called *faces* of P . In fact, all polytopes are finite intersections of their supporting half-spaces corresponding to their faces codimension 1, see [45, Theroem 2.4.3]. Furthermore, we have the following characterization theorem of polytopes, we refer to [45, Corollary 2.4.4 and Theorem 2.4.6] for the proof.

Theorem 2.18 *A convex set is a bounded intersection of finitely many closed half-spaces iff it is a polytope. Moreover, there exists $\xi_1, \dots, \xi_k \in \mathbb{S}^{n-1}$ with respect to which a polytope P is uniquely represented as $P = \{x \in \mathbb{R}^n : \langle x, \xi_i \rangle \leq h_P(\xi_i) \forall i \in \{1, \dots, k\}\}$.*

The vectors $\xi_1, \dots, \xi_k \in \mathbb{S}^{n-1}$ of Theorem 2.18 are normal vectors of P with $h_P(\xi_i)$'s uniquely determining the polytope and are chosen so that $P \cap H_P(\xi_i)$ are the 1-codimensional faces.

2.2 Convex geometry

We enlist some essential features and geometric notions of convex domains with smooth boundaries which have some generalizations to more general convex domains.

Recalling the notion of smoothness of boundaries, $\Omega \subset \mathbb{R}^n$ is said to be of class C^k (resp. $C^{k,\alpha}$) for $k \in \mathbb{N}$ and $0 < \alpha \leq 1$ if for any $x \in \partial\Omega$, there exists $r > 0$ and a bijection $\psi : B_r(x) \rightarrow \mathbb{R}^n$ with $\psi, \psi^{-1} \in C^k$ (resp. $C^{k,\alpha}$) such that $\psi(B_r(x) \cap \Omega) \subset \mathbb{R}_+^n$ and $\psi(B_r(x) \cap \partial\Omega) \subset \partial\mathbb{R}_+^n$; equivalently, for any $x \in \partial\Omega$ there exists a neighborhood U and $\phi \in C^k(\mathbb{R}^{n-1})$ (resp. $C^{k,\alpha}(\mathbb{R}^{n-1})$) such that after a possible rotation, we have

$$\begin{aligned} \Omega \cap U &= U \cap \{(x_0, x') \in \mathbb{R} \times \mathbb{R}^{n-1} : \phi(x') < x_0\}; \\ \partial\Omega \cap U &= U \cap \{(x_0, x') \in \mathbb{R} \times \mathbb{R}^{n-1} : \phi(x') = x_0\}. \end{aligned} \tag{2.19}$$

In addition, if Ω is convex then ϕ of (2.19) is a convex function. If Ω is of class C^k (resp. $C^{k,\alpha}$), then any $f \in C^k(\partial\Omega)$ (resp. $C^{k,\alpha}(\partial\Omega)$) can be extended to a function in $C^k(\bar{\Omega})$ (resp. $C^{k,\alpha}(\bar{\Omega})$).

If ∂K is of class C^1 for a convex domain K , then for every $x \in \partial K$ there exists a normal vector at x as in Definition 2.7 that is unique up to scaling. This gives rise to $\mathbf{g}_K : \partial K \rightarrow \mathbb{S}^{n-1}$, called *Gauss map*, where $\mathbf{g}_K(x)$ is the outer unit normal at $x \in \partial K$. Thus, notice that for any $z \in p_K^{-1}(x)$, we have $\mathbf{g}_K(x) = (z - x)/|z - x| = \xi_K(z)$ with $\xi_K(z)$ is as in (2.5) and in terms of support function, from (2.13), we have

$$h_K(\mathbf{g}_K(x)) = \langle x, \mathbf{g}_K(x) \rangle, \quad \forall x \in \partial K. \tag{2.20}$$

In particular, for balls we have $\mathbf{g}_{B_r(x_0)} : \partial B_r(x_0) \rightarrow \mathbb{S}^{n-1}$ given by $\mathbf{g}_{B_r(x_0)}(x) = (x - x_0)/r$ for any $x_0 \in \mathbb{R}^n$ and $r > 0$. Recall that the tangent space at $x \in \partial K$ is given by

$$T_x(\partial K) = \{y \in \mathbb{R}^n : \langle y, \mathbf{g}_K(x) \rangle = 0\}$$

and from (2.20), we observe that $H_K(\mathbf{g}_K(x)) = \{x\} + T_x(\partial K)$. The boundary ∂K being a C^1 -manifold, we have $\mathbf{g}_K \in C^1(\partial K, \mathbb{S}^{n-1})$ and we recall some familiar notions from differential geometry. Since $T_x(\partial K) = T_{\mathbf{g}_K(x)}(\mathbb{S}^{n-1})$, the following *Weingarten map*, is defined by

$$\mathcal{W}_K(x) := d\mathbf{g}_K(x) : T_x(\partial K) \rightarrow T_x(\partial K).$$

The *principal curvatures* are the eigenvalues $\kappa_1(x), \dots, \kappa_{n-1}(x)$ of $\mathcal{W}_K(x)$; the *mean curvature* and the *Gaussian curvature* are respectively given by

$$H(x) = \frac{\text{Tr}(\mathcal{W}_K(x))}{n-1} = \frac{1}{n-1}(\kappa_1 + \dots + \kappa_{n-1}) \quad \text{and} \quad \kappa(x) = \det(\mathcal{W}_K(x)) = \kappa_1 \dots \kappa_{n-1}.$$

If K is strictly convex then $K \cap H_K(\mathbf{g}_K(x)) = \{x\}$ for any $x \in \partial K$ and $\mathbf{g}_K^{-1} : \mathbb{S}^{n-1} \rightarrow \partial K$ is well defined; moreover, h_K being differentiable everywhere in $\mathbb{R}^n \setminus \{0\}$, from (2.20) we have

$$h_K(\xi) = \langle \xi, \mathbf{g}_K^{-1}(\xi) \rangle \quad \text{and} \quad \nabla h_K(\xi) = \mathbf{g}_K^{-1}(\xi), \quad \forall \xi \in \mathbb{S}^{n-1}. \tag{2.21}$$

Definition 2.22 A convex set K is said to be of class C_+^k (resp. $C_+^{k,\alpha}$) for any $k \in \mathbb{N}$, $\alpha \in (0, 1]$ if ∂K is of class C^k (resp. $C^{k,\alpha}$) and the Gauss map $\mathbf{g}_K : \partial K \rightarrow \mathbb{S}^{n-1}$ is a diffeomorphism.

The Gauss map being a diffeomorphism is equivalent to the Weingarten map being of maximal rank everywhere. This is further equivalent to all principal curvatures being non-zero or equivalently the Gaussian curvature being non-zero; the convexity implies that the curvatures are positive everywhere. The corresponding domains are called *strongly convex*.

If K is of class C_+^2 , note that $\mathbf{g}_K^{-1} \in C^1(\mathbb{S}^{n-1}, \partial K)$ and $h_K \in C^2(\mathbb{R}^n \setminus \{0\})$. Hence, from homogeneity of h_K , (2.21) and (2.10), we have

$$h_K(y) = \langle y, \nabla h_K(y) \rangle \quad \text{and} \quad D^2 h_K(y)y = 0, \quad \forall y \in \mathbb{R}^n \setminus \{0\}, \tag{2.23}$$

where the latter equality of the above can be obtained by differentiating the former. We have the inverse Weingarten map

$$\mathcal{W}_K^{-1}(\xi) := (\mathcal{W}_K(\mathbf{g}_K^{-1}(\xi)))^{-1} = d\mathbf{g}_K^{-1}(\xi) : T_\xi(\mathbb{S}^{n-1}) \rightarrow T_\xi(\mathbb{S}^{n-1})$$

defined for all $\xi \in \mathbb{S}^{n-1}$, which is the non-singular part of $D^2 h_K(\xi)$. Indeed, with $\xi \in \mathbb{S}^{n-1}$ being fixed, there exists an orthonormal basis $\{e^1, \dots, e^{n-1}, \xi\}$ of \mathbb{R}^n and hence, $\{e^i\}$ span the tangent space $T_\xi(\mathbb{S}^{n-1})$. Any $x \in \mathbb{R}^n$, in this basis, is given by

$$x = \sum_{k=1}^{n-1} x^k e^k + \langle x, \xi \rangle \xi, \quad \text{with} \quad x^k = \langle x, e^k \rangle. \tag{2.24}$$

Since $D^2 h_K(\xi)\xi = 0$ from (2.23), the only non-zero entries of $D^2 h_K(\xi)$ are $\langle D^2 h_K(\xi)e^j, e^i \rangle$ for $i, j \in \{1, \dots, n-1\}$ which, from (2.21), are the entries of $d\nabla h_K(\xi) = d\mathbf{g}_K^{-1}(\xi) = \mathcal{W}_K^{-1}(\xi)$, with respect to the above basis. In other words, we have

$$D^2 h_K(\xi) = \sum_{i,j=1}^{n-1} \langle \mathcal{W}_K^{-1}(\xi)e^j, e^i \rangle e^i \otimes e^j. \tag{2.25}$$

Thus, $D^2 h_K(\xi)$ has eigenvalues $\{1/\kappa_1(\mathbf{g}_K^{-1}(\xi)), \dots, 1/\kappa_{n-1}(\mathbf{g}_K^{-1}(\xi)), 0\}$, where κ_i 's are the principal curvatures of ∂K . With $\xi \in U \subset \mathbb{S}^{n-1}$ and a coordinate chart $\varphi : U \rightarrow V \subset \mathbb{R}^{n-1}$, the covariant derivatives of $f : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ of first and second orders are locally defined by

$$\begin{aligned} (i) \quad \nabla f &:= \sum_{i=1}^{n-1} (\nabla_i f) e^i, \quad \text{where} \quad \nabla_i f(x) := \partial_i(f \circ \varphi^{-1})(\varphi(x)), \\ (ii) \quad \nabla^2 f &:= \sum_{i,j=1}^{n-1} (\nabla_{i,j} f) e^i \otimes e^j, \quad \text{where} \quad \nabla_{i,j} f(x) := \partial_{i,j}(f \circ \varphi^{-1})(\varphi(x)), \end{aligned} \tag{2.26}$$

Letting $\chi = \varphi^{-1} : V \rightarrow U \subset \mathbb{S}^{n-1}$, note that since $|\chi|^2 = 1$ hence by differentiating successively we have $\langle \chi, \partial_j \chi \rangle = 0$ and $\langle \chi, \partial_{i,j} \chi \rangle = -\langle \partial_i \chi, \partial_j \chi \rangle$. We can choose U, φ such that $\chi(0) = \xi$ and that $\partial_j \chi(0) = e^j$ for all $j \in \{1, \dots, n-1\}$ taking $U = \mathbb{S}_+^{n-1} = \{x \in \mathbb{R}^n : \langle x, \xi \rangle = \sqrt{1 - \sum_{i=1}^{n-1} |x^i|^2}\}$ and $\varphi = (x^1, \dots, x^{n-1})$, leading to $\chi(z_1, \dots, z_{n-1}) = \sum_{i=1}^{n-1} z_i e^i + \sqrt{1 - \sum_{i=1}^{n-1} |z_i|^2} \xi$. Thus, we have $\partial_j \chi = e^j - (z_j / \sqrt{1 - \sum_{i=1}^{n-1} |z_i|^2}) \xi$ and $\partial_{i,j} \chi(0) = -\delta_{i,j} \chi(0)$ (one can define a similar chart on the other hemisphere for $U = \mathbb{S}_-^{n-1}$ corresponding to the negative square root). Taking h_K restricted to \mathbb{S}^{n-1} , (2.23) can be written in terms of local coordinates as

$$h_K \circ \chi = \langle \chi, \nabla h_K \circ \chi \rangle \quad \text{and} \quad (D^2 h_K \circ \chi) \chi = 0. \tag{2.27}$$

By differentiating and using the above, we get

$$\partial_j (h_K \circ \chi) = \langle \partial_j \chi, \nabla h_K \circ \chi \rangle + \langle \chi, (D^2 h_K \circ \chi) \partial_j \chi \rangle = \langle \partial_j \chi, \nabla h_K \circ \chi \rangle.$$

Evaluating at 0, we get $\nabla_j h_K(\xi) = \langle \nabla h_K(\xi), e^j \rangle$. Hence, from (2.24) and (2.27), notice that

$$\nabla h_K(\xi) = h_K(\xi) \xi + \nabla h_K(\xi). \tag{2.28}$$

By differentiating twice at 0 and using the choice of chart, we get

$$\begin{aligned} \partial_{i,j} (h_K \circ \chi)(0) &= \langle \partial_{i,j} \chi(0), \nabla h_K \circ \chi(0) \rangle + \langle \partial_j \chi(0), (D^2 h_K \circ \chi)(0) \partial_i \chi(0) \rangle \\ &= -\delta_{i,j} \langle \chi(0), \nabla h_K(\chi(0)) \rangle + \langle e^j, (D^2 h_K(\chi(0))) e^i \rangle \end{aligned}$$

which, from (2.23) and (2.25), further implies

$$\nabla^2 h_K(\xi) = \mathcal{W}_K^{-1}(\xi) - h_K(\xi) \mathbb{I}, \tag{2.29}$$

where \mathbb{I} is the identity matrix. Therefore, we have the following for C_+^2 domains;

$$\det(\nabla^2 h_K(\xi) + h_K(\xi) \mathbb{I}) = \det(\mathcal{W}_K^{-1}(\xi)) = 1/\kappa(\mathbf{g}_K^{-1}(\xi)), \tag{2.30}$$

where κ is the Gaussian curvature. One can show that a domain is C_+^2 iff it is C^2 and $\kappa > 0$. From transformation rule of the Jacobian, note that

$$(\mathbf{g}_K)_* \mathcal{H}^{n-1} \llcorner_{\partial K} = |\det(d \mathbf{g}_K^{-1})| \mathcal{H}^{n-1} \llcorner_{\mathbb{S}^{n-1}} = 1/(\kappa \circ \mathbf{g}_K^{-1}) \mathcal{H}^{n-1} \llcorner_{\mathbb{S}^{n-1}}$$

and from (2.30), for any $f \in L^1(\partial K, \mathcal{H}^{n-1})$, we have

$$\int_{\partial K} f(x) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{S}^{n-1}} f(\mathbf{g}_K^{-1}(\xi)) \det(\nabla^2 h_K(\xi) + h_K(\xi) \mathbb{I}) d\xi. \tag{2.31}$$

In general, since convex functions are locally Lipschitz, the boundary of a convex domain Ω can be locally written as a graph of a Lipschitz function and the Gauss map can be defined \mathcal{H}^{n-1} -a.e. on $\partial\Omega$. Precisely, for a small enough neighborhood U , if $\phi \in \mathcal{K}(\mathbb{R}^{n-1}) \cap C^{0,1}(\mathbb{R}^{n-1})$ is as in (2.19) then, for a.e. $x' \in \mathbb{R}^{n-1}$, ϕ is differentiable and

$$g_\Omega(\phi(x'), x') = (-1, \nabla \phi(x')) / \sqrt{1 + |\nabla \phi(x')|^2}.$$

Moreover, a classical theorem of Aleksandrov [6, 9] (see also [18]) is the following.

Theorem 2.32 (Aleksandrov) *If $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex, then ϕ is twice-differentiable a.e.*

Consequently, for a convex domain Ω and $K = \bar{\Omega}$, we have $\mathcal{W}_K(x)$ and $\kappa(x) \geq 0$ well-defined for \mathcal{H}^{n-1} -a.e. $x \in \partial\Omega$ and the surface measure S_Ω on \mathbb{S}^{n-1} induced by the almost everywhere defined Gauss map is defined by

$$S_\Omega(E) := \mathcal{H}^{n-1}\left(\{x \in \partial\Omega : \mathbf{g}_\Omega(x) \text{ is well-defined, } \mathbf{g}_\Omega(x) \in E\}\right) \tag{2.33}$$

for any measurable $E \subseteq \mathbb{S}^{n-1}$ and we denote $S_\Omega = S_K$ if $K = \bar{\Omega}$. For any $w \in L^1(\mathbb{S}^{n-1}, \mathcal{H}^{n-1})$, the integral formula (2.31) generalizes to

$$\int_{\partial\Omega} w(\mathbf{g}_\Omega(x)) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{S}^{n-1}} w(\xi) dS_\Omega(\xi). \tag{2.34}$$

It is evident that if g_Ω is defined everywhere then $S_\Omega = (\mathbf{g}_\Omega)_*\mathcal{H}^{n-1}$ and if Ω is C^2_+ , then $dS_\Omega(\xi) = \det(\nabla^2 h_\Omega(\xi) + h_\Omega(\xi)\mathbb{I})d\xi$ as earlier.

For a general convex set K the Gauss map is re-defined as a set-valued map at $x \in \partial K$ as

$$\mathbf{g}_K(x) = \{\xi \in \mathbb{S}^{n-1} : \langle x' - x, \xi \rangle \leq 0, \forall x' \in K\}, \tag{2.35}$$

i.e. the set of all normal vectors $\xi \in \mathbb{S}^{n-1}$ of every supporting hyperplane $H_K[x]$ at $x \in \partial K$. The push forward $(\mathbf{g}_K)_*$ can be re-defined with respect to the re-defined inverse image

$$\mathbf{g}_K^{-1}(E) = \{x \in \partial K : \mathbf{g}_K(x) \cap E \neq \emptyset\}, \tag{2.36}$$

so that we can re-define $S_K = (\mathbf{g}_K)_*\mathcal{H}^{n-1}$ as $S_K(E) = \mathcal{H}^{n-1}(\mathbf{g}_K^{-1}(E))$ for any convex domain K and measurable $E \subseteq \mathbb{S}^{n-1}$. If $K = \bar{\Omega}$ for a convex domain Ω then, the set (2.35) is a singleton for \mathcal{H}^{n-1} -a.e. $x \in \partial K$. More generally, the set (2.35) can be a cone at non-smooth boundary points or a discrete set if K is of lower dimension. Henceforth, according to the regularity of the convex set K , the notions \mathbf{g}_K and S_K shall be assumed at the appropriate level of generality, hereafter.

For any $A \subset \mathbb{R}^n$, let us denote $C_+(A)$ as the class of positive, continuous functions $f : A \rightarrow \mathbb{R}$. If $0 \in \Omega$ then from (2.12) it is clear that $h_\Omega \in C_+(\mathbb{S}^{n-1})$. Conversely, any $h \in C_+(\mathbb{S}^{n-1})$ coincides with the support function of a domain Ω_h except possibly for a set of zero surface measure on \mathbb{S}^{n-1} ; the domain Ω_h , called the *Wulff shape* of h , is defined by

$$\bar{\Omega}_h = \bigcap_{\xi \in \mathbb{S}^{n-1}} \{x \in \mathbb{R}^n : \langle x, \xi \rangle \leq h(\xi)\}. \tag{2.37}$$

Indeed, since h is positive and continuous, it is evident that Ω_h is a bounded convex domain with $0 \in \Omega_h$. One can extend h to be a 1-homogeneous function on $\mathbb{R}^n \setminus \{0\}$ as $y \mapsto |y|h(y/|y|)$ and letting $K = \bar{\Omega}_h$, clearly $h_K \leq h$. Moreover, if $x \in \partial K$ then (2.37) implies $\langle x, \xi \rangle = h(\xi)$ for some $\xi \in \mathbb{S}^{n-1}$ and (2.12) implies $\langle x, \xi' \rangle = h_K(\xi')$ for some $\xi' \in \mathbb{S}^{n-1}$; if $h(\xi) > h_K(\xi)$ then $\xi \neq \xi'$ and hence $H_{\xi, h(\xi)}$ and $H_{\xi', h_K(\xi')}$ are two different supporting hyperplanes at x with normals ξ, ξ' making x a non-smooth point of ∂K . Thus Aleksandrov's theorem enforces $h_K(\xi) = h(\xi)$ for S_K -a.e. $\xi \in \mathbb{S}^{n-1}$ and thus we have a generalization of Theorem 2.11 as follows.

Theorem 2.38 *Given any $h \in C_+(\mathbb{S}^{n-1})$, there exists a bounded convex domain $\Omega = \Omega_h$ as in (2.37) such that $0 \in \Omega$ and $h_\Omega(\xi) = h(\xi)$ holds for S_Ω -a.e. $\xi \in \mathbb{S}^{n-1}$.*

Let Ω be a compact convex domain with $0 \in \Omega$ and $K = \bar{\Omega}$. Then, integral formulae on ∂K can be obtained independently using the *radial map* $r_K : \mathbb{R}^n \setminus \{0\} \rightarrow [0, \infty)$ defined by

$$r_K(y) = \sup \{r \geq 0 : ry \in K\}. \tag{2.39}$$

It is clear that for any $\alpha > 0$, we have $r_{\alpha K}(y) = \alpha r_K(y)$ and $r_K(\alpha y) = r_K(y)/\alpha$. Also, it is not hard to see that $r_K(y)y \in \partial K$ for any $y \in \mathbb{R}^n \setminus \{0\}$. This gives rise to the *radial projection* $\rho_K : \mathbb{S}^{n-1} \rightarrow \partial K$ defined by $\rho_K(\theta) = r_K(\theta)\theta$, which is the polar coordinate image on ∂K . From the polar coordinate formula (2.2) we have the following, for any $F \in L^1(K)$

$$\int_K F(x)dx = \omega_n \int_{\mathbb{S}^{n-1}} \left[\int_0^{r_K(\theta)} F(r\theta)r^{n-1} dr \right] d\theta; \tag{2.40}$$

in particular, we have $|K| = (\omega_n/n) \int_{\mathbb{S}^{n-1}} r_K(\theta)^n d\theta$. Hence, from the divergence theorem, we have that $\int_{\partial\Omega} \langle x, \mathbf{g}_K(x) \rangle d\mathcal{H}^{n-1} = \int_{\mathbb{S}^{n-1}} |\rho_K(\theta)|^n d\theta$, and

$$\mathcal{H}^{n-1} \llcorner_{\partial K} = (\rho_K)_* \left(|x|^n / \langle x, \mathbf{g}_K(x) \rangle \theta \llcorner_{\mathbb{S}^{n-1}} \right)$$

so that, $d\mathcal{H}^{n-1} = |\rho_K(\theta)|^n / \langle \rho_K(\theta), \mathbf{g}_K(\rho_K(\theta)) \rangle d\theta = |r_K(\theta)|^{n-1} / \langle \theta, \mathbf{g}_K(\rho_K(\theta)) \rangle d\theta$, in infinitesimal form. Thus we have the following due to Aleksandrov (see [7, Lemma 1]).

Theorem 2.41 *Let Ω be a bounded convex domain with $0 \in \Omega$, $K = \bar{\Omega}$ and \mathbf{g}_K be the Gauss map defined \mathcal{H}^{n-1} -a.e. on ∂K . For any $f \in L^1(\partial K, \mathcal{H}^{n-1})$, we have the following;*

$$\int_{\partial K} f(x) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{S}^{n-1}} f(\rho_K(\theta)) \frac{r_K(\theta)^{n-1}}{\langle \mathbf{g}_K(\rho_K(\theta)), \theta \rangle} d\theta. \tag{2.42}$$

Furthermore, as \mathcal{H}^{n-1} -a.e. points on ∂K are smooth, hence recalling (2.20), we can also re-write the above formulae using $d\mathcal{H}^{n-1} = |\rho_K(\theta)|^n / h_K(\mathbf{g}_K(\rho_K(\theta))) d\theta$. Unlike (2.30), (2.31) above, this density can be used for any convex body that are not necessarily smooth.

2.3 Hausdorff distance and approximations

Here we provide some details on properties of distance functions on convex sets, results on compactness and approximations.

The *Hausdorff distance* between Borel sets $E, E' \subset \mathbb{R}^n$ is defined as

$$d_{\mathcal{H}}(E, E') = \max \left(\sup\{\text{dist}(y, E) : y \in E'\}, \sup\{\text{dist}(y, E') : y \in E\} \right);$$

equivalently, we have $d_{\mathcal{H}}(E, E') = \sup_{y \in \mathbb{R}^n} |\text{dist}(y, E) - \text{dist}(y, E')|$. The Hausdorff distance of convex sets can be characterized by support functions as

$$d_{\mathcal{H}}(E, F) = d_{\mathcal{H}}(\partial E, \partial F) = \|h_E - h_F\|_{L^\infty(\mathbb{S}^{n-1})}, \tag{2.43}$$

see [45, Lemmas 1.8.1, 1.8.14]. Any compact convex set K can be approximated by polytopes with respect to $d_{\mathcal{H}}$, since, given any $\varepsilon > 0$, the compactness implies there exists points x_1, \dots, x_N such that K contained in the union of $B_\varepsilon(x_j)$'s and then the polytope $P = \text{conv}(\{x_1, \dots, x_N\})$ satisfies $P \subset K \subset P + \varepsilon \mathbb{S}^{n-1}$ and $d_{\mathcal{H}}(K, P) < \varepsilon$. More generally, the space of bounded convex bodies is locally compact with respect to the Hausdorff distance.

Theorem 2.44 (*Blaschke selection theorem*) *Given any bounded sequence of convex sets $\{K_m\}$, there exists a subsequence $\{K_{m_j}\}$ and a convex set K such that $d_{\mathcal{H}}(K_{m_j}, K) \rightarrow 0^+$ as $j \rightarrow \infty$.*

If $d_{\mathcal{H}}(K_j, K) \rightarrow 0^+$ as $j \rightarrow \infty$, (2.43) and Aleksandrov's theorem implies $\mathbf{g}_{K_j} \rightarrow \mathbf{g}_K$ \mathcal{H}^{n-1} a.e. Moreover, since $\text{dist}(x_0, \partial K) \leq \text{dist}(x_0, \partial K_j) + d_{\mathcal{H}}(K_j, K)$, it is not hard to see from local compactness, that then any subsequential limit x_0 of radial projections $\rho_{K_j}(\theta)$

lies on ∂K , leading to uniform convergence of the projections, hence $r_{K_j} \rightarrow r_K$ uniformly as $j \rightarrow \infty$ and the densities of \mathcal{H}^{n-1} with respect to radial projections

$$\frac{|\rho_{K_j}(\theta)|^n}{h_{K_j}(\mathbf{g}_{K_j}(\rho_{K_j}(\theta)))} \rightarrow \frac{|\rho_K(\theta)|^n}{h_K(\mathbf{g}_K(\rho_K(\theta)))} \quad \mathcal{H}^{n-1} - a.e. \theta \in \mathbb{S}^{n-1}. \tag{2.45}$$

We have the following approximation theorem for convex domains, see Weil [47, 48].

Theorem 2.46 *Given any convex set K with \mathcal{H}^{n-1} -a.e. defined Gaussian curvature κ , there exists a sequence $\{K_j\}$ of convex sets of class C_+^2 such that as $j \rightarrow \infty$ we have $d_{\mathcal{H}}(K_j, K) \rightarrow 0^+$ and if κ_j is the Gaussian curvature of K_j then the following holds:*

- (1) $1/\kappa_j \rightarrow 1/\kappa$ pointwise \mathcal{H}^{n-1} -a.e. as $j \rightarrow \infty$;
- (2) $1/\kappa_j \rightarrow 1/\kappa$ in $L^1(\mathbb{S}^{n-1}, \mathcal{H}^{n-1})$ as $j \rightarrow \infty$;
- (3) $\int (1/\kappa_j) d\xi \rightarrow \int (1/\kappa) d\xi$ as $j \rightarrow \infty$.

Thus, from Theorem 2.46, we can conclude that given a convex domain Ω with Gaussian curvature κ , there exists C_+^2 (or $C_+^{2,\alpha}$) domains Ω_j with Gaussian curvature κ_j , such that as $j \rightarrow \infty$, we have $d_{\mathcal{H}}(\Omega_j, \Omega) \rightarrow 0^+$ and $(\mathbf{g}_{\Omega_j})_* \mathcal{H}^{n-1} \rightarrow (\mathbf{g}_{\Omega})_* \mathcal{H}^{n-1}$, i.e. for any $w \in C(\mathbb{S}^{n-1})$,

$$\lim_{j \rightarrow \infty} \int_{\mathbb{S}^{n-1}} \frac{w(\xi)}{\kappa_j(\mathbf{g}_{K_j}^{-1}(\xi))} d\xi = \int_{\mathbb{S}^{n-1}} \frac{w(\xi)}{\kappa(\mathbf{g}_K^{-1}(\xi))} d\xi. \tag{2.47}$$

For general convex sets, we have the weak convergence of surface measures with respect to Hausdorff distance, i.e. as $j \rightarrow \infty$,

$$S_{K_j} \rightharpoonup S_K \quad \text{if } d_{\mathcal{H}}(K_j, K) \rightarrow 0^+, \tag{2.48}$$

when the measures are defined with respect to (2.35) and (2.36), see [45, Theorem 4.2.1]. In general, there are results of weak convergence with respect to the Hausdorff distance for any support measures and curvature measures for general convex sets, the proofs involve approximation of convex sets by convex domains, in particular, polytopes with non-empty interiors that have combinatorial formulae for support measures. We refer to [45, Chaper 4,5] for further details.

2.4 p -harmonic functions and measures

Here we provide some preliminaries and some well-known classical results for p -harmonic functions and p -harmonic measures. We refer to [25, 29, 30, 33, 36, 46], etc. for more details.

Just as a harmonic function in a domain $\Omega \subset \mathbb{R}^n$ being minimizer of the Dirichlet energy $\int_{\Omega} |\nabla w|^2 dx$ with $w|_{\partial\Omega} = f$, is the unique solution of the Dirichlet problem $\Delta v = 0$ in Ω and $v = f$ on $\partial\Omega$, the minimizers of the p -Dirichlet energy $\int_{\Omega} |\nabla w|^p dx$ are weak solutions to the p -Laplacian equation $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0$ in Ω , i.e.

$$\int_{\Omega} |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = 0, \quad \forall \phi \in C_0^\infty(\Omega),$$

and are called p -harmonic functions for $1 \leq p < \infty$. They coincide with harmonic functions for $p = 2$. The existence of weak solution $u \in W^{1,p}(\Omega)$ is classical and follows from direct methods of calculus of variations. We have the following monotonicity inequality for any

measurable $E \subseteq \mathbb{R}^n$, functions $u, v \in W^{1,p}(E)$ and a constant $c = c(p) > 0$, given by

$$\begin{aligned} & \int_E (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v, \nabla u - \nabla v) \, dx \\ & \geq c \begin{cases} \int_E |\nabla u - \nabla v|^p \, dx & \text{if } p \geq 2, \\ \frac{(\int_E |\nabla u - \nabla v|^p \, dx)^{2/p}}{(\int_E (|\nabla u| + |\nabla v|)^p \, dx)^{2/p-1}} & \text{if } 1 < p < 2, \end{cases} \end{aligned} \tag{2.49}$$

which can be used to show the comparison principle for the p -Laplacian, i.e. $\Delta_p v \leq \Delta_p u$ on Ω and $u \leq v$ on $\partial\Omega$ in the trace sense implies $u \leq v$ on Ω ; we refer to [25] for a proof. The uniqueness of weak solutions of $\Delta_p u = 0$ in Ω and $u = f$ on $\partial\Omega$ follows easily from the comparison principle. The regularity theory of p -harmonic functions is more involved. It has been shown by DiBenedetto [17], Lewis [35] and Tolksdorf [46] that the weak solutions of $\Delta_p u = 0$ in Ω for $1 < p < \infty$ are locally $C^{1,\beta}$, i.e. there exists $\beta = \beta(n, p) \in (0, 1)$ such that for any $B \subset\subset \Omega$, we have

$$\begin{aligned} \|u\|_{C^{1,\beta}(B)} &= \|u\|_{L^\infty(B)} + \|\nabla u\|_{L^\infty(B, \mathbb{R}^n)} + \sup_{\substack{x,y \in B \\ x \neq y}} \frac{|\nabla u(x) - \nabla u(y)|}{|x - y|^\beta} \\ &\leq c \|u\|_{W^{1,p}(\Omega)}, \end{aligned} \tag{2.50}$$

for some $c = c(n, p, \text{diam}(B)) > 0$. For $p > 2$ the regularity is optimal. Furthermore, the continuity of the gradient implies that if $\nabla u \neq 0$ in $\Omega' \subset \Omega$, then we can conclude $u \in C^\infty(\Omega')$ from Schauder estimates, see [21]. The boundary regularity is also known for $\Delta_p u = 0$ in Ω and $u = f$ on $\partial\Omega$ if Ω is of class $C^{1,\alpha}$ and $f \in C^{1,\alpha}(\partial\Omega)$, see Lieberman [39]; in this case $u \in C^{1,\beta}(\bar{\Omega})$ for some $\beta = \beta(n, p, \alpha) \in (0, 1)$ along with the following global estimate

$$\|u\|_{C^{1,\beta}(\bar{\Omega})} \leq C(n, p, \alpha, \|f\|_{C^{1,\alpha}(\partial\Omega)}, \|u\|_{W^{1,p}(\Omega)}, \Omega). \tag{2.51}$$

Given a bounded and sufficiently regular domain $\Omega \subset \mathbb{R}^n$, for any $f \in C(\partial\Omega)$ and v being the solution of the Dirichlet problem $\Delta v = 0$ in Ω and $v = f$ on $\partial\Omega$, we have

$$v(x) = \int_{\partial\Omega} f(y) d\omega^x(y) \tag{2.52}$$

from maximum principle and Riesz representation theorem, where ω^x is a measure on $\partial\Omega$ referred to as the *harmonic measure* at $x \in \Omega$. The measure can be prescribed by the Green's function with a pole at x , i.e. the weak (distributional) solution of the Dirichlet problem $\Delta G^x = \delta_x$ in Ω and $G^x = 0$ on $\partial\Omega$. If $\partial\Omega$ is smooth enough with unit normal ν , it is not difficult to see that $d\omega^x = (\partial G^x / \partial \nu) d\mathcal{H}^{n-1} = \langle \nabla G^x, \nu \rangle d\mathcal{H}^{n-1}$. This notion can be generalized to the case for the p -Laplacian for $1 < p < \infty$. Given a neighbourhood N of $\partial\Omega$, if $u \in W^{1,p}(\Omega \cap N)$ is a positive weak solution to the p -Laplacian in $\Omega \cap N$, then upon zero extension $u \in W^{1,p}(N)$. Since p -superharmonic functions form a non-negative distribution as shown in [25], from a theorem of Schwartz we have that there exists a non-negative Radon measure ω_p on $\partial\Omega$ such that

$$\int_{\Omega} |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle \, dx = - \int_{\partial\Omega} \phi \, d\omega_p,$$

for any $\phi \in C_0^\infty(N)$; the measure ω_p is called the *p -harmonic measure* associated to u . Such measures can also be defined for more general \mathcal{A} -harmonic functions that are referred as Riesz measures, see [25, 28]. If $\partial\Omega$ is Lipschitz with unit normal $\nu_{\partial\Omega}$ defined almost

everywhere on $\partial\Omega$, then $d\omega_p = -|\nabla u|^{p-2} \langle \nabla u, \nu_{\partial\Omega} \rangle d\mathcal{H}^{n-1}$; moreover, since $\{u > 0\}$ in Ω we have $\partial\Omega = \partial\{u > 0\}$ and $\nu_{\partial\Omega} = -\nabla u/|\nabla u|$ and hence

$$d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega}. \tag{2.53}$$

Evidently, for the case $p = 2$, if $N = \mathbb{R}^n \setminus \{x\}$ and $u = G^x$ then $\omega_2 = \omega^x$. More generally, the function u may have a similar blow-up inside $\Omega \setminus N$ but the p -harmonicity in $\Omega \cap N$ implies that $u \in C_{\text{loc}}^{1,\beta}(\Omega \cap N)$ for some $\beta \in (0, 1)$ from (2.50). Given such a neighbourhood, it is evident that we can select finer reduced neighborhood N' of $\partial\Omega$ i.e. $\partial\Omega \subset N' \subseteq N$, having the same properties. Such neighborhood N (without relabelling) when chosen up to possible reduction, we can assume without loss of generality that $\nabla u \neq 0$ in $\Omega \cap N$ so that $u \in C^\infty(\Omega \cap N)$ as stated above, and furthermore, we can assume

$$\|u\|_{L^\infty(\partial N \cap \Omega)} + \|\nabla u\|_{L^\infty(\partial N \cap \Omega)} < \infty;$$

in other words, all possible singularities of u are strictly in the interior of $\Omega \setminus \bar{N}$. Any such reductions can be chosen as long as it contains $\partial\Omega$ without changing any properties of the measure since the support of the measure is within $\partial\Omega$.

The fundamental solution of the p -Laplacian, given by

$$\Phi(x) = \begin{cases} 1/|x|^{\frac{n-p}{p-1}} & \text{for } p \neq n; \\ \log(1/|x|) & \text{for } p = n, \end{cases} \tag{2.54}$$

solves $\Delta_p \Phi = 0$ in $\mathbb{R}^n \setminus \{0\}$ and can be used to form examples of p -harmonic measures.

The above notion of p -harmonic measure is defined for any open connected domains Ω , including convex domains. The notion extends to a convex set K of non-empty interior in the same way taking Ω as the interior of K so that given a neighbourhood N of ∂K and a function $u \in W^{1,p}(K \cap N)$, we have the notion of a p -harmonic measure ω_p associated to u given by $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial K}$. Then, this can be further extended naturally to the case of K being a convex set of empty interior i.e. $K = \partial K$; given N is a neighborhood of K and $u \in W^{1,p}(N)$ being a positive weak solution to the p -Laplacian in $N \setminus K$ and u vanish on K , there exists a non-negative Radon measure ω_p on K such that

$$\int_N |\nabla u|^{p-2} \langle \nabla u, \nabla \phi \rangle dx = - \int_K \phi d\omega_p,$$

for any $\phi \in C_0^\infty(N)$ similarly as above which is called the p -harmonic measure associated to u , and $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_K$ if K is Lipschitz. We refer to [37] for further details on quasi-linear equations on low dimensional sets. In general, since for all convex sets K we have ∂K locally Lipschitz, hence $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial K}$ for $u = u_K$.

2.5 Minkowski problem

Given a positive finite Borel measure μ on \mathbb{S}^{n-1} that satisfies the conditions (1.1), the Minkowski problem can be regarded as the inquiry for existence of a convex domain Ω satisfying

$$\mu_\Omega = \mu, \tag{2.55}$$

where $\mu_\Omega = (\mathbf{g}_\Omega)_* \eta$ for some prescribed measure η on $\partial\Omega$ and \mathbf{g}_Ω is the Gauss map. The prescribed measure η is typically absolutely continuous with respect to $\mathcal{H}^{n-1} \llcorner_{\partial\Omega}$ and with

respect to the Gauss map, we can express the density of the induced measure on \mathbb{S}^{n-1} as

$$d\mu_\Omega(\xi) = \mathcal{F}[h_\Omega](\xi)d\xi \tag{2.56}$$

for a functional $\mathcal{F} : \mathcal{K}(\mathbb{S}^{n-1}) \rightarrow \mathbb{R}$. Existence of domains for the Minkowski problem involves computation of the first variation of measures corresponding to $\Omega^t = \Omega + t\Omega_0$ with the support functions $h_{\Omega^t} = h_\Omega + th_{\Omega_0}$,

$$\mathcal{L}_\Omega[v] := \left. \frac{d}{dt} \right|_{t=0} \mathcal{F}[h_{\Omega^t}] = \left. \frac{d}{dt} \right|_{t=0} \mathcal{F}[h_\Omega + th_{\Omega_0}]. \tag{2.57}$$

This is used to show the existence of domains using continuity method or variational techniques of constrained minimization problems.

In case of the original Minkowski problem, we have $\eta = \mathcal{H}^{n-1}$ and $\mu_\Omega = (\mathbf{g}_\Omega)_*\mathcal{H}^{n-1}$. The case of the given measure being discrete was considered in [41, 42], where the corresponding convex domains are polytopes. The continuous case has been shown in [5, 7, 19] and the smooth case in [13]. If Ω is of class C^2_+ and the covariant derivatives are as in (2.26) defined by the charts as shown in the previous subsection, then recalling (2.30) and (2.31), we note that the density (2.56) in this case, is the reciprocal of the Gauss curvature, i.e.

$$d\mu_\Omega(\xi) = \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I})d\xi = \frac{d\xi}{\kappa(\mathbf{g}_\Omega^{-1}(\xi))}.$$

Furthermore, if $\mu_\Omega = (\mathbf{g}_\Omega)_*\eta$ where $d\eta = fd\mathcal{H}^{n-1}$ for a function $f : \partial\Omega \rightarrow \mathbb{R}$, then we have

$$\mathcal{F}[h_\Omega](\xi) = \frac{(f \circ \mathbf{g}_\Omega^{-1})(\xi)}{\kappa(\mathbf{g}_\Omega^{-1}(\xi))} = f(\mathbf{g}_\Omega^{-1}(\xi)) \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}). \tag{2.58}$$

The above formulae hold for S_Ω -a.e. $\xi \in \mathbb{S}^{n-1}$ for a general convex domain. As examples of the above, the prescribed measure is the harmonic measure ω at origin in [26] where we have $\mu_\Omega = (\mathbf{g}_\Omega)_*\omega$ and $f = (\partial G/\partial v)$ where G is the Green’s function with pole at 0; in the case of capacity measures in [15], we have $f = |\nabla U|^p$ where U is the capacity function.

In our case, we consider $\mu_\Omega = (\mathbf{g}_\Omega)_*\omega_p$ where ω_p a the p harmonic measure with respect to a function u and the domain Ω being convex, we have $f = |\nabla u|^{p-1}$ for $1 < p < \infty$.

3 Existence for p -harmonic Minkowski problem

In this section, we consider the Minkowski problem (2.55) for p -harmonic measures. We establish the existence of solutions of the problem. Henceforth, we shall denote

$$\mu_\Omega = (\mathbf{g}_\Omega)_*\omega_p, \tag{3.1}$$

where ω_p is the p harmonic measure with respect to a function $u = u_\Omega \in W^{1,p}(\Omega \cap N)$ given by $d\omega_p = |\nabla u|^{p-1}d\mathcal{H}^{n-1} \llcorner_{\partial\Omega}$ where u is p -harmonic in $\Omega \cap N$ and satisfies

$$\begin{cases} \operatorname{div}(|\nabla u|^{p-2}\nabla u) = 0 & \text{in } \Omega \cap N; \\ u > 0 & \text{in } \Omega; \\ u = 0 & \text{on } \partial\Omega, \end{cases} \tag{3.2}$$

where N is a neighbourhood of $\partial\Omega$; thus, $u \in W^{1,p}(N)$ upon zero extension. Up to possible reduction, the choice of N is made so that $\nabla u \neq 0$ in $\Omega \cap N$ and

$$\|u\|_{L^\infty(\bar{N} \cap \Omega)} + \|\nabla u\|_{L^\infty(\bar{N} \cap \Omega)} < \infty, \tag{3.3}$$

and we also assume that ∂N is C^∞ . For a general bounded convex set K , if $K = \bar{\Omega}$ for a convex domain Ω then we define $\mu_K = \mu_\Omega$; if K is of empty interior, i.e. $K = \partial K$, we define $\mu_K = (\mathbf{g}_K)_* \omega_p$ in the sense of (2.35) and (2.36), where ω_p is the p -harmonic measure with respect to a function $u = u_K \in W^{1,p}(N)$ with N being a neighbourhood of K , given by $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1} \llcorner_K$ where u is p -harmonic in N and satisfies

$$\begin{cases} \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0 & \text{in } N \setminus K; \\ u \geq 0 & \text{in } N; \\ u = 0 & \text{on } K. \end{cases} \tag{3.4}$$

Thus, in general, we have μ_K defined on \mathbb{S}^{n-1} associated to $u = u_K \in W^{1,p}(N)$ as

$$\mu_K(E) = \int_{\mathbf{g}_K^{-1}(E)} |\nabla u|^{p-1} d\mathcal{H}^{n-1}, \quad \text{for any measurable } E \subseteq \mathbb{S}^{n-1}, \tag{3.5}$$

where \mathbf{g}_K and $\mathbf{g}_K^{-1}(E)$ are in the sense of (2.35) and (2.36). If $K = \bar{\Omega}$ for a convex domain Ω then (3.5) coincides with the standard push forward measure (3.1).

3.1 First variation of density of p -harmonic measures

Here, we consider Ω as strongly convex domain of class $C_+^{2,\alpha}$ so that $\mathbf{g}_\Omega : \partial\Omega \rightarrow \mathbb{S}^{n-1}$ is a diffeomorphism. Let its support function be h_Ω . Using the same notations as Sect. 2, from (3.1) and (2.53), we have

$$d\mu_\Omega = |\nabla u(F_\Omega(\xi))|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega} = \mathcal{F}[h_\Omega](\xi) d\xi,$$

where $F_\Omega(\xi) := \mathbf{g}_\Omega^{-1}(\xi) = \nabla h_\Omega(\xi)$ and recalling (2.58), we have

$$\mathcal{F}[h_\Omega](\xi) = |\nabla u(F_\Omega(\xi))|^{p-1} \det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}).$$

Henceforth, we shall denote $h = h_\Omega$ and $F = F_\Omega$. Thus, from (2.20) and (2.23), we have $h(\xi) = h_\Omega(\xi) = \langle \xi, \mathbf{g}_\Omega^{-1}(\xi) \rangle = \langle x, \mathbf{g}_\Omega(x) \rangle$ and

$$F(\xi) = F_\Omega(\xi) = \mathbf{g}_\Omega^{-1}(\xi) = \nabla h(\xi).$$

As in Sect. 2, we shall consider the orthonormal frame field $\{e^1, \dots, e^{n-1}\}$ of \mathbb{S}^{n-1} so that for any $\xi \in \mathbb{S}^{n-1}$, the tangent space $T_\xi(\mathbb{S}^{n-1})$ is spanned by $\{e^i(\xi)\}$; furthermore, we denote covariant derivatives with respect to the frame as in (2.26) with respect to the coordinate charts as in Sect. 2. We have $\det(\nabla^2 h_\Omega + h_\Omega \mathbb{I}) = 1/(\kappa \circ \mathbf{g}_\Omega^{-1})$ from (2.30) and (2.31), and recalling (2.53), for any integrable function $f : \partial\Omega \rightarrow \mathbb{R}$, we have

$$\begin{aligned} & \int_{\partial\Omega} f(x) d\omega_p(x) \\ &= \int_{\mathbb{S}^{n-1}} f(\mathbf{g}_\Omega^{-1}(\xi)) |\nabla u(\mathbf{g}_\Omega^{-1}(\xi))|^{p-1} \det(\nabla^2 h_\Omega(\xi) + h_\Omega(\xi) \mathbb{I}) d\xi. \end{aligned} \tag{3.6}$$

We consider a convex domain $\Omega_0 \subset \mathbb{R}^n$, also of class $C_+^{2,\alpha}$ to find the first variation for p -harmonic measures on the following family of domains,

$$\Omega^t = \Omega + t\Omega_0 \quad \text{with support functions } h_{\Omega^t} = h_\Omega + th_{\Omega_0} = h + tv. \tag{3.7}$$

Without loss of generality, we assume $0 \in \Omega_0$ so that $v \geq 0$ and hence we have $\Omega \subseteq \Omega^t$ for $t \geq 0$ and $\Omega^t \subseteq \Omega$ for $t \leq 0$, more generally $\Omega^s \subseteq \Omega^t$ if $s \leq t$. We take a small enough

$$\tau = \tau(n, d_{\mathcal{H}}(\partial\Omega, \partial N), d_{\mathcal{H}}(\partial\Omega_0, \partial N), \|u\|_{W^{1,p}(N)}) > 0, \tag{3.8}$$

so that for all $|t| \leq \tau$ the domain Ω^t is also of class $C^{2,\alpha}_+$ and $\partial\Omega^t \subset N$ and $\partial N \cap \Omega = \partial N \cap \Omega^t$. Also, note that, from (2.43) we have $d_{\mathcal{H}}(\Omega^t, \Omega) = |t|\|v\|_{L^\infty(\mathbb{S}^{n-1})}$ and hence $d_{\mathcal{H}}(\Omega^t, \Omega) \rightarrow 0^+$ as $t \rightarrow 0$. Moreover, from Steiner’s formula for volumes and mixed volumes, we have

$$t \mapsto |\Omega + t\Omega_0|, \mathcal{H}^{n-1}(\partial\Omega + t\partial\Omega_0) \text{ are smooth;} \tag{3.9}$$

in fact, the above maps are polynomials, see [45, Chapter 5].

We consider the p -harmonic measures corresponding to $u(\cdot, t) \in W^{1,p}(\Omega^t \cap N)$ defined as the weak solution of the following Dirichlet problem

$$\begin{cases} \operatorname{div}(|\nabla u(\cdot, t)|^{p-2}\nabla u(\cdot, t)) = 0, & \text{in } \Omega^t \cap N; \\ u(x, t) = 0, & \forall x \in \partial\Omega^t \cap N; \\ u(x, t) = u\left(\frac{x}{1+t}\right), & \forall x \in \partial N \cap \Omega^t; \end{cases} \tag{3.10}$$

for $t \in [-\tau, \tau]$ for $\tau > 0$ small enough, so that upon zero extension, $u(\cdot, t) \in W^{1,p}(N)$. Then the corresponding measure is defined by

$$d\mu_{\Omega^t} = |\nabla u(F(\xi, t), t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega^t} = \mathcal{F}[h_{\Omega^t}](\xi) d\xi$$

where $F(\xi, t) := F_{\Omega^t}(\xi) = \mathbf{g}_{\Omega^t}^{-1}(\xi) = \nabla h(\xi) + t\nabla v(\xi)$ and

$$\mathcal{F}[h + tv](\xi) = |\nabla u(F(\xi, t), t)|^{p-1} \det(\nabla^2 h + h\mathbb{I} + t(\nabla^2 v + v\mathbb{I})). \tag{3.11}$$

We need to establish some properties of the first variation as in (2.57). The behavior with respect to dilations is provided in the following lemma.

Lemma 3.12 *Let \mathcal{F} be as in (3.11) and $\mathcal{L}_h[v] = \frac{d}{dt}\big|_{t=0} \mathcal{F}[h + tv]$, then we have*

$$\mathcal{F}[(1+t)h] = (1+t)^{n-p} \mathcal{F}[h] \text{ and } \mathcal{L}_h[h] = (n-p)\mathcal{F}[h], \tag{3.13}$$

for all $|t| \leq \tau$ with a small enough $\tau > 0$ as in (3.8).

Proof Let $\lambda = 1 + t$ and consider the case of $\Omega_0 = \Omega$ and $v = h$ of the above. Then, we have $\Omega^t = \lambda\Omega$ and $F_{\Omega^t} = \mathbf{g}_{\Omega^t}^{-1} = \lambda\nabla h$. Let us denote $u_\lambda(\cdot) = u(\cdot, \lambda - 1)$ as the weak solution of the problem (3.10) for this case, i.e.

$$\begin{cases} \operatorname{div}(|\nabla u_\lambda|^{p-2}\nabla u_\lambda) = 0, & \text{in } \lambda\Omega \cap N; \\ u_\lambda(x) = 0, & \forall x \in \partial\lambda\Omega \cap N; \\ u_\lambda(x) = u(x/\lambda), & \forall x \in \partial N \cap \lambda\Omega; \end{cases} \tag{3.14}$$

for $|\lambda - 1| \leq \tau$ being small enough. Using these in (3.11), we can conclude

$$\begin{aligned} \mathcal{F}[\lambda h](\xi) &= |\nabla u_\lambda(\lambda\nabla h(\xi))|^{p-1} \lambda^{n-1} \det(\nabla^2 h + h\mathbb{I}) \\ &= \left(\frac{|\nabla u_\lambda(\lambda\nabla h(\xi))|}{|\nabla u(\nabla h(\xi))|}\right)^{p-1} \lambda^{n-1} \mathcal{F}[h](\xi). \end{aligned}$$

Now, note that (3.2) trivially implies that $u(x/\lambda)$ is also a solution of (3.14) and hence, from uniqueness $u_\lambda(x) = u(x/\lambda)$ in $\lambda\Omega$. Therefore, using $\nabla u_\lambda(x) = \frac{1}{\lambda}\nabla u(x/\lambda)$ on the above, we get (3.13) (the second part of (3.13) follows immediately from the first) and the proof is finished. \square

Towards the computation of the first variation, we need to establish some basic properties. From (2.50), (2.51), we know that weak solutions $u(\cdot, t) \in W^{1,p}(\Omega^t \cap N)$ of (3.10) are $C^{1,\beta}$ in Ω^t for any $t \in [-\tau, \tau]$ and some $\beta = \beta(n, p) \in (0, 1)$.

First, we show uniform continuity of $t \mapsto u(\cdot, t)$.

Lemma 3.15 *For any $t \in [-\tau, \tau]$, if $u(\cdot, t) \in C_{loc}^{1,\beta}(\Omega^t \cap N)$ is the solution of the Dirichlet problem (3.10), then $t \mapsto u(\cdot, t), \nabla u(\cdot, t)$ are uniformly continuous on compact subsets of N .*

Proof From (3.10), note that for any $\eta \in W^{1,p}(N)$, we have

$$\begin{aligned} & \int_{\Omega^t \cap N} |\nabla u(x, t)|^{p-2} \langle \nabla u(x, t), \nabla \eta(x) \rangle dx \\ &= \int_{\partial(\Omega^t \cap N)} \eta(x) |\nabla u(x, t)|^{p-2} \langle \nabla u(x, t), \nu_{\partial(\Omega^t \cap N)}(x) \rangle d\mathcal{H}^{n-1}(x) \\ &= \frac{1}{|1+t|^{p-2}(1+t)} \int_{\Omega^t \cap \partial N} \eta(x) |\nabla u(\frac{x}{1+t})|^{p-2} \langle \nabla u(\frac{x}{1+t}), \nu_{\Omega^t \cap \partial N}(x) \rangle d\mathcal{H}^{n-1}(x) \\ & \quad + \int_{\partial\Omega^t} \eta(x) |\nabla u(x, t)|^{p-2} \langle \nabla u(x, t), \mathbf{g}_{\Omega^t}(x) \rangle d\mathcal{H}^{n-1}(x) \end{aligned} \tag{3.16}$$

for any $t \in [-\tau, \tau]$. From boundary conditions of (3.10) and comparison principle, we can conclude $u(x, t) > 0$ in $\Omega^t \cap N$, hence $\mathbf{g}_{\Omega^t}(x) = -\nabla u(x, t)/|\nabla u(x, t)|$ for all $x \in \partial\Omega^t$, $|t| < \tau$ for a small enough τ in (3.8). Using (3.16) for $t, s \in [-\tau, \tau]$ and taking their difference, we obtain

$$\begin{aligned} & \int_{\Omega^t \cap N} |\nabla u(x, t)|^{p-2} \langle \nabla u(x, t), \nabla \eta(x) \rangle dx - \int_{\Omega^s \cap N} |\nabla u(x, s)|^{p-2} \langle \nabla u(x, s), \nabla \eta(x) \rangle dx \\ &= \frac{1}{|1+t|^{p-2}(1+t)} \int_{\Omega^t \cap \partial N} \eta(x) |\nabla u(\frac{x}{1+t})|^{p-2} \langle \nabla u(\frac{x}{1+t}), \nu_{\Omega^t \cap \partial N}(x) \rangle d\mathcal{H}^{n-1}(x) \\ & \quad - \frac{1}{|1+s|^{p-2}(1+s)} \int_{\Omega^s \cap \partial N} \eta(x) |\nabla u(\frac{x}{1+s})|^{p-2} \langle \nabla u(\frac{x}{1+s}), \nu_{\Omega^s \cap \partial N}(x) \rangle d\mathcal{H}^{n-1}(x) \\ & \quad - \left(\int_{\partial\Omega^t} \eta(x) |\nabla u(x, t)|^{p-1} d\mathcal{H}^{n-1}(x) - \int_{\partial\Omega^s} \eta(x) |\nabla u(x, s)|^{p-1} d\mathcal{H}^{n-1}(x) \right) \end{aligned}$$

Then taking $\eta(x) = u(x, t) - u(x, s)$ on the above and using (2.49), it is not difficult to see that we can obtain

$$\int_{\Omega^{\min\{t,s\}} \cap N} |\nabla u(x, t) - \nabla u(x, s)|^p dx \leq cM^p \omega(|t-s|),$$

where $c = c(n, p, \tau) > 0$, $M = \|u\|_{L^\infty(\Omega \cap N)} + \|\nabla u\|_{L^\infty(\Omega \cap N)} + \max_{|t| \leq \tau} \|u(\cdot, t)\|_{W^{1,p}(N)} < \infty$ and $\omega(|t-s|) \rightarrow 0^+$ as $t \rightarrow s$, since ∇u is continuous in $\Omega \cap N$ and $\nabla u(\cdot, t)$ is continuous in $\Omega^t \cap N$ from (2.50) for all t , and furthermore $|\Omega^t| - |\Omega^s| \rightarrow 0$ and $\mathcal{H}^{n-1}(\partial\Omega^t) - \mathcal{H}^{n-1}(\partial\Omega^s) \rightarrow 0$ as $t \rightarrow s$ from (3.9). The continuity of $x \mapsto \nabla u(x, t)$ for every t also implies that all points are Lebesgue points and hence the above is enough to conclude the following pointwise convergences

$$|u(x, t) - u(x, s)| + |\nabla u(x, t) - \nabla u(x, s)| \rightarrow 0^+ \text{ as } t \rightarrow s.$$

From (2.50), we also have the following local estimates for any $B \subset\subset N$ with M as above and $c = c(n, p, \tau, \text{diam}(B)) > 0$, given by

$$|u(x, t) - u(y, t)| \leq cM|x-y|, \quad |\nabla u(x, t) - \nabla u(y, t)| \leq cM|x-y|^\beta, \quad \forall x, y \in B,$$

for every $t \in [-\tau, \tau]$, that affirms equicontinuity of $\{u(\cdot, t), \nabla u(\cdot, t)\}_{0 \leq |t| \leq \tau}$. Uniform continuity follows from the Arzelà-Ascoli theorem and the proof is finished. \square

Remark 3.17 Notice that, $u(\cdot, t)$ being the solution of (3.10), from uniqueness and (3.2) we have $u(x, 0) = u(x)$ and hence, from the Lemma (3.15), $u(x, t) \rightarrow u(x)$ uniformly as $t \rightarrow 0$. Hence, $\nabla u(\cdot, t) \neq 0$ in $\Omega^t \cap N$ and $\|u(\cdot, t)\|_{L^\infty(\bar{N} \cap \Omega^t)} + \|\nabla u(\cdot, t)\|_{L^\infty(\bar{N} \cap \Omega^t)} < \infty$ for all $|t| \leq \tau$.

Now we need to investigate the differentiability of $t \mapsto u(\cdot, t)$. This is shown in the following that involve some standard regularity results and adaptations of the techniques in Colesanti et al. [15]. For any function $f(\cdot, t)$ related to (3.10), let us denote

$$\dot{f}(x) = \left. \frac{\partial}{\partial t} \right|_{t=0} f(x, t),$$

if the derivative exists. Thus, we have $\dot{F}(\xi) = \nabla v(\xi)$ and in the following proposition, we find \dot{u} . Note that if $t \mapsto u(\cdot, t)$ is differentiable, then $\partial_t(\operatorname{div}(|\nabla u|^{p-2} \nabla u)) = \operatorname{div}(A(\nabla u) \nabla \partial_t u) = 0$, where we henceforth denote

$$A(\nabla u) := |\nabla u|^{p-2} \left(\mathbb{I} + (p-2) \frac{\nabla u \otimes \nabla u}{|\nabla u|^2} \right), \tag{3.18}$$

which is uniformly elliptic modulo the weight $|\nabla u|^{p-2}$. Indeed, it is not hard to see that

$$\min\{1, p-1\} |\nabla u|^{p-2} |\zeta|^2 \leq \langle A(\nabla u) \zeta, \zeta \rangle \leq \max\{1, p-1\} |\nabla u|^{p-2} |\zeta|^2, \quad \forall \zeta \in \mathbb{R}^n. \tag{3.19}$$

However, the differentiability of $t \mapsto u(\cdot, t)$ being unknown a priori, difference quotients are to be used. This is shown in the following.

Proposition 3.20 *If $u(\cdot, t) \in W^{1,p}(\Omega^t \cap N)$ is the solution of (3.10) and A is as in (3.18), then the following holds:*

- (1) $t \mapsto u(\cdot, t)$ is differentiable at $t = 0$ for all $x \in \bar{\Omega} \cap N$ and $\dot{u} \in C^{2,\beta}(\overline{\Omega \cap N})$;
- (2) \dot{u} is a solution of the equation $\operatorname{div}(A(\nabla u) \nabla \dot{u}) = 0$ in $\Omega \cap N$;
- (3) $\dot{u}(x) = -\langle \nabla u(x), x \rangle$ for all $x \in \partial N \cap \Omega$;
- (4) $\dot{u}(x) = |\nabla u(x)| v(\mathbf{g}_\Omega(x))$ for all $x \in \partial \Omega$.

Proof For $t \neq 0$, let $w(x, t) = (u(x, t) - u(x, 0))/t$ and for any $s \in [0, 1]$, let us denote

$$u_s(x, t) = su(x, t) + (1-s)u(x, 0).$$

Then, $u'_s := du_s/ds = tw(x, t)$ and for any $|t| \leq \tau$, note that we have

$$\begin{aligned} 0 &= \Delta_p u(x, t) - \Delta_p u(x, 0) = \int_0^1 \frac{d}{ds} (\Delta_p u_s(x, t)) ds \\ &= \int_0^1 \operatorname{div} (A(\nabla u_s(x, t)) \nabla u'_s(x, t)) ds = t \int_0^1 \operatorname{div} (A(\nabla u_s(x, t)) \nabla w(x, t)) ds \end{aligned} \tag{3.21}$$

where A is as in (3.18). Thus, for $t \neq 0$, we have that $w(x, t)$ is a solution of the equation $\operatorname{div}(B_u(x, t) \nabla w(x, t)) = 0$, where entries of B_u are given by

$$B_u(x, t)_{i,j} = \langle B_u(x, t) e_j, e_i \rangle = \int_0^1 \langle A(\nabla u_s(x, t)) e_j, e_i \rangle ds. \tag{3.22}$$

From Lemma 3.15 and Remark 3.17, note that $\nabla u_s(x, t) \rightarrow \nabla u(x, 0) = \nabla u(x)$ uniformly as $t \rightarrow 0$, which together with (3.22) implies $B_u(\cdot, t) \rightarrow A(\nabla u)$ uniformly as well. Also, from Remark 3.17, $\nabla u(\cdot, t) \neq 0$ in $\Omega^t \cap N$ for all $|t| \leq \tau$. Therefore, there exists $0 < \lambda \leq \Lambda < \infty$ possibly dependent on $n, p, \min_{|t| \leq \tau} \inf_{x \in \Omega^t \cap N} |\nabla u(x, t)|$ and $\max_{|t| \leq \tau} \sup_{x \in \Omega^t \cap N} |\nabla u(x, t)|$ such that from (3.19) and (3.22), we can conclude

$$\lambda|y|^2 \leq \langle B_u(x, t)y, y \rangle \leq \Lambda|y|^2, \quad \forall x \in (\Omega \cap \Omega^t) \cap N, \quad y \in \mathbb{R}^n,$$

for any $t \in [-\tau, \tau]$, thereby making the equations uniformly elliptic. Hence, for $t \neq 0$, we have $w(\cdot, t) \in C^\infty(\Omega \cap \Omega^t \cap N)$. The limit of $w(x, t)$ needs to be established as $t \rightarrow 0$ for the proof.

To this end, note that for $B \subset\subset \Omega \cap N$ there exists $\tau' = \tau'(B) \leq \tau$ such that $B \cap \Omega^t \neq \emptyset$ for $t \in [-\tau', \tau']$. From (2.50) used on the Eq. (3.10) implies that $u(\cdot, t) \in C_{\text{loc}}^{1,\beta}(\Omega^t \cap N)$ for $\beta = \beta(n, p) \in (0, 1)$, hence (3.19) and (3.22) gives $\|B_u(\cdot, t)_{i,j}\|_{C^{0,\sigma}(B \cap \Omega^t)} \leq C'(n, p, M, \text{diam}(B))$ for some $\sigma = \sigma(n, p, \beta) \in (0, 1)$, where $M = \max_{|t| \leq \tau} \|u(\cdot, t)\|_{W^{1,p}(N)}$. Then, from a standard compactness argument

$$\|B_u(\cdot, t)_{i,j}\|_{C^{0,\sigma}(B)} \leq C(n, p, M, \text{diam}(B)), \quad \forall t \in [-\tau', \tau'].$$

Therefore, we can invoke interior Schauder estimates, see [21, Theorem 6.2], for the equation $\text{div}(B_u(x, t)\nabla w(x, t)) = 0$ on $\Omega \cap N$ to conclude that there exists $0 < \tau' = \tau'(B) \leq \tau$ for any $B \subset\subset \Omega \cap N$ such that

$$\|w(\cdot, t)\|_{C^{2,\beta}(B)} \leq \Lambda', \quad \forall 0 < |t| \leq \tau', \tag{3.23}$$

for some $\Lambda' = \Lambda'(n, p, \lambda, M, \text{diam}(B)) > 0$ and $\beta \in (0, 1)$. It is not hard to see that (3.23) implies uniform boundedness and equicontinuity of the family $\{w(\cdot, t)\}_{0 < |t| \leq \tau'}$ and hence Arzelà-Ascoli theorem yields the existence of a convergent subsequence. For a sequence $t_k \rightarrow 0$ as $k \rightarrow \infty$, let $w \in W^{1,p}(\Omega \cap N)$ be such a subsequential limit, i.e.

$$w(x) := \lim_{k \rightarrow \infty} w(x, t_k) = \lim_{k \rightarrow \infty} \frac{u(x, t_k) - u(x, 0)}{t_k}, \quad \forall x \in \Omega \cap N; \tag{3.24}$$

then taking $t_k \rightarrow 0$ on (3.23) we get $w \in C_{\text{loc}}^{2,\alpha}(\Omega \cap N)$ and since $B_u(\cdot, t_k) \rightarrow A(\nabla u)$ uniformly, we have that w is a solution of $\text{div}(A(\nabla u)\nabla w) = 0$ in $\Omega \cap N$.

Now we look into the boundary behavior of w . Since Ω^t 's are of class $C^{2,\alpha}$ and $u \in C^\infty(\Omega^t \cap N)$ for all $|t| \leq \tau$, (2.51) used for the Eq. (3.10) implies that $u(\cdot, t) \in C^{1,\gamma}(\overline{\Omega^t \cap N})$ for some $\gamma = \gamma(n, p, \alpha) \in (0, 1)$ and $\|u(\cdot, t)\|_{C^{1,\gamma}(\overline{\Omega^t \cap N})} \leq C(n, p, \alpha, M, N)$ with $M > 0$ as above. This, together with (3.19), (3.22) and (3.9), further imply

$$\|B_u(\cdot, t)_{i,j}\|_{C^{0,\sigma}(\overline{\Omega \cap N})} \leq C(n, p, \alpha, M, N), \quad \forall t \in [-\tau, \tau],$$

for some $\sigma = \sigma(n, p, \gamma) \in (0, 1)$. Recalling $\partial N \cap \Omega^t = \partial N \cap \Omega$ for all $|t| \leq \tau$ and from boundary condition of (3.10), note that we have

$$\begin{aligned} w(x, t) &= \frac{u\left(\frac{x}{1+t}\right) - u(x)}{t} \\ &= -\frac{\langle \nabla u(x), x \rangle}{(1+t)} + O(|t|^\gamma), \quad \forall x \in \partial N \cap \Omega^t = \partial N \cap \Omega, t \neq 0, \end{aligned} \tag{3.25}$$

which implies $\|w(\cdot, t)\|_{L^\infty(\partial N \cap \Omega)} \leq C(n, p, \tau, \gamma, M, N)$. Also from boundary condition of (3.10), we can conclude that for some $\beta = \beta(n, p, \gamma) \in (0, 1)$,

$$\|w(\cdot, t)\|_{C^{2,\beta}(\partial(\Omega \cap \Omega^t \cap N))} \leq C(n, p, \tau, \gamma, M, N), \quad \forall 0 < |t| \leq \tau. \tag{3.26}$$

Therefore, we can invoke global Schauder estimates up to the boundary, see [21, Theorem 6.6], for the equation $\operatorname{div}(B_u(x, t)\nabla w(x, t)) = 0$ on $(\Omega \cap \Omega^t) \cap N$ to conclude

$$\|w(\cdot, t)\|_{C^{2,\beta}(\Omega \cap \Omega^t \cap N)} \leq \Lambda'', \quad \forall 0 < |t| \leq \tau, \tag{3.27}$$

for some $\Lambda'' = \Lambda''(n, p, \alpha, \lambda, \tau, M, N) > 0$ and $\beta \in (0, 1)$. Taking $t_k \rightarrow 0$ on (3.26) and (3.27) and using (3.9), we get $w \in C^{2,\beta}(\overline{\Omega \cap N})$ and $\|w\|_{C^{2,\beta}(\Omega \cap N)} \leq \Lambda''$. Using this we can find the boundary values of w at $x \in \partial(\Omega \cap N)$ by taking sequences $x_j \in \Omega \cap N$ such that $x_j \rightarrow x$ as $j \rightarrow \infty$ and using (3.24) as

$$w(x) = \lim_{j \rightarrow \infty} w(x_j) = \lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} w(x_j, t_k) = \lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} \frac{u(x_j, t_k) - u(x_j, 0)}{t_k}.$$

Since we have $\partial N \cap \Omega^{t_k} = \partial N \cap \Omega$, the evaluation of the limit is clear on $\partial N \cap \Omega$ taking $t_k \rightarrow 0$ and using (3.25) and continuity of ∇u , to obtain

$$w(x) = -\langle \nabla u(x), x \rangle, \quad \forall x \in \partial N \cap \Omega.$$

However, finding w at $\partial\Omega$ is more involved since values at $\partial\Omega^{t_k}$'s can accumulate to that on $\partial\Omega$ as $t_k \rightarrow 0$ and the limits may not be interchangeable in general. Here, the strong convexity of the domain is used to obtain sequences $x_k \in \partial\Omega^{t_k}$ with a common normal to evaluate the limit diagonally. Indeed, since Ω^t 's are of class $C^{2,\alpha}_+$, hence $\mathbf{g}_{\Omega^t} : \partial\Omega^t \rightarrow \mathbb{S}^{n-1}$ is a diffeomorphism for every $|t| \leq \tau$. Hence, for any $x \in \partial\Omega$ with $\mathbf{g}_{\Omega}(x) = \xi \in \mathbb{S}^{n-1}$, let $x_k = \mathbf{g}_{\Omega^{t_k}}^{-1}(\xi) \in \partial\Omega^{t_k}$. Then, recalling (3.7) and (2.21), notice that

$$x_k = \nabla h(\xi) + t_k \nabla v(\xi) = x + t_k \nabla v(\xi), \tag{3.28}$$

and thus $x_k \rightarrow x$ as $k \rightarrow \infty$. Since (3.26) and (3.27) together with (3.28) imply

$$|w(x_k, t_k) - w(x, t_k)| \leq C|x_k - x| = C|\nabla v(\xi)||t_k|,$$

for $C = C(n, p, \alpha, \lambda, \tau, M, N) > 0$. This is used in (3.24) to obtain

$$w(x) = \lim_{k \rightarrow \infty} w(x_k, t_k) = \lim_{k \rightarrow \infty} \frac{u(x_k, t_k) - u(x_k, 0)}{t_k} = \lim_{k \rightarrow \infty} -\frac{u(x_k)}{t_k}, \tag{3.29}$$

since $u(x_k, t_k) = 0$ from boundary conditions of (3.10) as $x_k \in \partial\Omega^{t_k}$. Furthermore, since $x \in \partial\Omega$, from boundary conditions of (3.2) we also have $u(x) = 0$. Also, since $u \in C^{1,\gamma}(\overline{\Omega \cap N})$ with $\|u\|_{C^{1,\gamma}(\Omega \cap N)} \leq C(n, p, \alpha, \|u\|_{L^\infty(\bar{N} \cap \Omega)} + \|\nabla u\|_{L^\infty(\bar{N} \cap \Omega)}, N)$, therefore, using (3.28), note that

$$u(x_k) - u(x) = \langle \nabla u(x), x_k - x \rangle + O(|x_k - x|^{1+\gamma}) = t_k \langle \nabla u(x), \nabla v(\xi) \rangle + O(|t_k|^{1+\gamma}).$$

Using these on (3.29) we obtain

$$w(x) = \lim_{k \rightarrow \infty} \frac{u(x) - u(x_k)}{t_k} = -\langle \nabla u(x), \nabla v(\xi) \rangle, \quad \forall x \in \partial\Omega. \tag{3.30}$$

Recall that, since $\partial\Omega = \partial\{u > 0\}$, the outer normal $\nu_{\partial\Omega} = \xi = -\nabla u/|\nabla u|$ and since $v = h_{\Omega_0}$, from (2.21), $v(\xi) = \langle \xi, \nabla v(\xi) \rangle$. Using these on (3.30), we finally obtain $w(x) = |\nabla u(x)|v(\xi)$.

Thus, we established that any subsequential limit w as in (3.24) must be a solution of the equation $\operatorname{div}(A(\nabla u)\nabla w) = 0$ in $\Omega \cap N$ and moreover, $w(x) = -\langle \nabla u(x), x \rangle$ for all $x \in \partial N \cap \Omega$ and $w(x) = |\nabla u(x)|v(\mathbf{g}_{\Omega}(x))$ for all $x \in \partial\Omega \cap N$. This equation being uniformly elliptic in $\Omega \cap N$, w is unique for every subsequence leading to $\liminf_{t \rightarrow 0} w(\cdot, t) = w = \limsup_{t \rightarrow 0} w(\cdot, t)$. Therefore, the limit exists and hence $w = \dot{u}$, which completes the proof. \square

In view of the above lemma, let us define the operator $\Lambda_u : C(\partial\Omega) \rightarrow C(\partial\Omega)$ as

$$\Lambda_u[f](x) = \langle \nabla w(x), \mathbf{g}_\Omega(x) \rangle, \tag{3.31}$$

for all $x \in \partial\Omega$, $f \in C(\partial\Omega)$, where $w \in C^1(\overline{\Omega \cap N})$ is the solution of the Dirichlet problem

$$\begin{cases} \operatorname{div}(A(\nabla u)\nabla w) = 0 & \text{in } \Omega \cap N; \\ w(x) = -\langle \nabla u(x), x \rangle & \forall x \in \partial N \cap \Omega; \\ w(x) = f(x) & \forall x \in \partial\Omega \cap N; \end{cases} \tag{3.32}$$

with $A(\nabla u)$ as in (3.18). Notice that Proposition 3.20 implies that

$$\Lambda_u[|\nabla u|(v \circ \mathbf{g}_\Omega)](x) = \langle \nabla \dot{u}(x), \mathbf{g}_\Omega(x) \rangle, \quad \forall x \in \partial\Omega. \tag{3.33}$$

We have the following technical lemma.

Lemma 3.34 *Let $w \in C^1(\overline{\Omega \cap N})$ be the solution of (3.32). For any $\eta \in C^1(\Omega)$, we have*

$$\begin{aligned} & (p-1) \int_{\partial\Omega} \eta |\nabla u|^{p-2} \langle \nabla w(x), \mathbf{g}_\Omega(x) \rangle d\mathcal{H}^{n-1}(x) \\ &= \int_{\Omega \cap N} |\nabla u|^{p-2} \left(\langle \nabla w, \nabla \eta \rangle + \frac{(p-2)}{|\nabla u|^2} \langle \nabla w, \nabla u \rangle \langle \nabla u, \nabla \eta \rangle \right) dx \\ &+ \int_{\partial N \cap \Omega} \eta \langle A(\nabla u)(\nabla u + D^2u x), v_{\partial N \cap \Omega}(x) \rangle d\mathcal{H}^{n-1}(x). \end{aligned} \tag{3.35}$$

Proof If $w \in C^1(\overline{\Omega \cap N})$ is the solution of (3.32), then for any $\eta \in C^1(\Omega)$, we have

$$\int_{\Omega \cap N} \langle A(\nabla u)\nabla w, \nabla \eta \rangle dx = \int_{\partial(\Omega \cap N)} \eta \langle A(\nabla u)\nabla w, v_{\partial(N \cap \Omega)}(x) \rangle d\mathcal{H}^{n-1}(x). \tag{3.36}$$

Recalling (3.18), we note that

$$A(\nabla u)\nabla w = |\nabla u|^{p-2} \left(\nabla w + \frac{(p-2)}{|\nabla u|^2} \langle \nabla w, \nabla u \rangle \nabla u \right). \tag{3.37}$$

Since $\mathbf{g}_\Omega(x) = -\nabla u(x)/|\nabla u(x)|$ for $x \in \partial\Omega$, hence the right hand side of (3.36), together with (3.37), becomes

$$\begin{aligned} & \int_{\partial\Omega} \eta \langle A(\nabla u)\nabla w, \mathbf{g}_\Omega(x) \rangle d\mathcal{H}^{n-1}(x) + \int_{\partial N \cap \Omega} \eta \langle A(\nabla u)\nabla w, v_{\partial N \cap \Omega}(x) \rangle d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial\Omega} \eta |\nabla u|^{p-2} \left(\langle \nabla w, \mathbf{g}_\Omega(x) \rangle + \frac{(p-2)}{|\nabla u|^2} \langle \nabla w, \nabla u \rangle \langle \nabla u, \mathbf{g}_\Omega(x) \rangle \right) d\mathcal{H}^{n-1}(x) \\ &- \int_{\partial N \cap \Omega} \eta \langle A(\nabla u)(\nabla u + D^2u x), v_{\partial N \cap \Omega}(x) \rangle d\mathcal{H}^{n-1}(x) \\ &= (p-1) \int_{\partial\Omega} \eta |\nabla u|^{p-2} \langle \nabla w(x), \mathbf{g}_\Omega(x) \rangle d\mathcal{H}^{n-1}(x) \\ &- \int_{\partial N \cap \Omega} \eta \langle A(\nabla u)(\nabla u + D^2u x), v_{\partial N \cap \Omega}(x) \rangle d\mathcal{H}^{n-1}(x). \end{aligned}$$

Using (3.37) on the left-hand side of (3.36) together with the above, we obtain (3.35) and the proof is finished. □

Using the above, we show that Λ_u is self-adjoint on $L^2(\partial\Omega, |\nabla u|^{p-2}\mathcal{H}^{n-1})$ in the following.

Proposition 3.38 *Let $\Lambda_u : C(\partial\Omega) \rightarrow C(\partial\Omega)$ be as in (3.31). Then, for any $\psi_1, \psi_2 \in C(\partial\Omega)$, the following holds,*

$$\int_{\partial\Omega} \psi_1 \Lambda_u[\psi_2] |\nabla u|^{p-2} d\mathcal{H}^{n-1} = \int_{\partial\Omega} \psi_2 \Lambda_u[\psi_1] |\nabla u|^{p-2} d\mathcal{H}^{n-1}. \tag{3.39}$$

Proof Given any $\psi_1, \psi_2 \in C(\partial\Omega)$, there exists $w_1, w_2 \in C^1(\overline{\Omega \cap N})$ that are solutions of (3.32) with $w_1 = \psi_1$ and $w_2 = \psi_2$ on $\partial\Omega$ and $w_1 = -\langle \nabla u, x \rangle = w_2$ on $\partial N \cap \Omega$. Hence, using (3.35) of Lemma 3.34, note that

$$\begin{aligned} & \int_{\partial\Omega} \psi_1 \Lambda_u[\psi_2] |\nabla u|^{p-2} d\mathcal{H}^{n-1} \\ &= \int_{\partial\Omega} w_1 \langle \nabla w_2(x), \mathbf{g}_\Omega(x) \rangle |\nabla u|^{p-2} d\mathcal{H}^{n-1}(x) \\ &= \frac{1}{p-1} \int_{\Omega \cap N} |\nabla u|^{p-2} \left(\langle \nabla w_2, \nabla w_1 \rangle + \frac{(p-2)}{|\nabla u|^2} \langle \nabla w_2, \nabla u \rangle \langle \nabla u, \nabla w_1 \rangle \right) dx \\ &\quad - \frac{1}{p-1} \int_{\partial N \cap \Omega} \langle \nabla u, x \rangle \langle A(\nabla u)(\nabla u + D^2u x), \nu_{\partial N \cap \Omega}(x) \rangle d\mathcal{H}^{n-1}(x), \end{aligned}$$

which is symmetric with respect to w_1 and w_2 . This completes the proof. □

Remark 3.40 For the special case $p = 2$, notice that we have $A(\nabla u) = \mathbb{I}$ and $\Lambda_u = \Lambda$ corresponds to the Neumann operator as in [26, 27].

Our next goal is to compute the first variation explicitly to establish its self-adjointness. We shall denote $C[\nabla^2 h + h\mathbb{I}] =$ cofactor matrix of $(\nabla^2 h + h\mathbb{I})$ (transpose of adjoint) so that,

$$C\nabla^2 h + h\mathbb{I} = \det(\nabla^2 h + h\mathbb{I})\mathbb{I} = 1/(\kappa \circ \mathbf{g}_\Omega^{-1})\mathbb{I}. \tag{3.41}$$

Furthermore, since $\mathcal{W}_K^{-1} = \nabla^2 h + h\mathbb{I}$ for $K = \bar{\Omega}$, note that the above can be rewritten as

$$\mathcal{W}_K(x) = \kappa(x) C[\nabla^2 h + h\mathbb{I}](\mathbf{g}_\Omega(x)), \quad \forall x \in \partial\Omega. \tag{3.42}$$

The function u of (3.2) being p -harmonic is $C^{1,\beta}$ up to the boundary from (2.51), is uniformly elliptic in a neighborhood of $\partial\Omega$ since Ω is of class $C^{2,\alpha}_+$. Hence, from boundary Schauder estimates, see [21], D^2u is pointwise well-defined on $\partial\Omega$. The following lemma which prescribes the Hessian D^2u on $\partial\Omega$, has been shown for the case $p = 2$ in [26, Lemma A].

Remark 3.43 The sign in the following Lemma is reversed to that of [26, Lemma A] as the Green’s function $G \leq 0$ in [26], therefore our choice $u \geq 0$ corresponds to $u = -G$ for $p = 2$.

Lemma 3.44 *Let $u : \Omega \rightarrow \mathbb{R}$ be as in (3.2) and $\{e^1, \dots, e^{n-1}\}$ be an orthonormal frame field of \mathbb{S}^{n-1} such that for any $\xi \in \mathbb{S}^{n-1}$ the unit vectors $e^i = e^i(\xi) \in \mathbb{S}^{n-1}$ span the tangent space $T_\xi(\mathbb{S}^{n-1})$. The covariant derivatives being defined as in (2.26) with respect to the suitable local coordinate charts related to the frame, we have the following:*

- (1) $\langle D^2u(F(\xi))e^i, e^j \rangle = -\kappa(F(\xi))|\nabla u(F(\xi))| C_{i,j}[\nabla^2 h + h\mathbb{I}]$;
- (2) $\langle D^2u(F(\xi))\xi, e^i \rangle = -\kappa(F(\xi)) \sum_j C_{i,j}[\nabla^2 h + h\mathbb{I}] \nabla_j(|\nabla u(F(\xi))|)$;
- (3) $\langle D^2u(F(\xi))\xi, \xi \rangle = \frac{1}{(p-1)} \kappa(F(\xi))|\nabla u(F(\xi))| \text{Tr}(C[\nabla^2 h + h\mathbb{I}])$;

where $C_{i,j}[\cdot] = \langle C[\cdot]e^j, e^i \rangle$ are entries of the cofactor matrix as in (3.41) for $i, j \in \{1, \dots, n-1\}$ with respect to this frame and $F(\xi) = \mathbf{g}_\Omega^{-1}(\xi) = \nabla h(\xi)$.

Proof Let $K = \bar{\Omega}$ and the coordinate charts for the covariant derivatives be as in Sect. 2 in a neighborhood U of $\xi \in \mathbb{S}^{n-1}$, i.e. for $0 \in V \subset \mathbb{R}^{n-1}$ the coordinates $\chi : V \rightarrow U \subset \mathbb{S}^{n-1}$ satisfying $\chi(0) = \xi$, $\partial_j \chi(0) = e^j$ and $\partial_{i,j} \chi(0) = -\delta_{i,j} \chi(0)$ for all $i, j \in \{1, \dots, n-1\}$.

By differentiating $F \circ \chi = \mathbf{g}_\Omega^{-1} \circ \chi = \nabla h \circ \chi$, we note that $\partial_j(F \circ \chi) = (D^2 h \circ \chi) \partial_j \chi$ for any $j \in \{1, \dots, n-1\}$ and hence, at $\chi = \chi(z) \in U \subset \mathbb{S}^{n-1}$ for any $z \in V \subset \mathbb{R}^{n-1}$, we have $\nabla_j F(\chi) = D^2 h(\chi) \partial_j \chi$, in particular, at $0 \in V$, we have

$$\nabla_j F(\xi) = D^2 h(\xi) e^j = \mathcal{W}_K^{-1}(\xi) e^j = (\nabla^2 h(\xi) + h(\xi) \mathbb{I}) e^j. \tag{3.45}$$

Since any $\chi \in U \subset \mathbb{S}^{n-1}$ is the outer unit normal at $F(\chi) \in \partial\Omega$ as $F = \mathbf{g}_\Omega^{-1}$, hence we have $\chi = -\nabla u(F(\chi))/|\nabla u(F(\chi))|$. Using this together with the above, we get

$$\begin{aligned} \langle \nabla_j F(\chi), \nabla u(F(\chi)) \rangle &= -|\nabla u(F(\chi))| \langle \nabla_j F(\chi), \chi \rangle \\ &= -|\nabla u(F(\chi))| \langle D^2 h(\chi) \chi, \partial_j \chi \rangle = 0, \end{aligned} \tag{3.46}$$

since, recalling (2.27), $D^2 h(\chi) \chi = 0$. Now, we differentiate (3.46) by covariant derivative ∇_i at $\xi = \chi(0)$ for any $i \in \{1, \dots, n-1\}$ to obtain

$$\langle D^2 u(F(\xi)) \nabla_i F(\xi), \nabla_j F(\xi) \rangle + \langle \nabla_{i,j} F(\xi), \nabla u(F(\xi)) \rangle = 0. \tag{3.47}$$

We can take χ defined on the lower hemisphere without loss of generality and $\nabla_{i,j} F(\xi)$ can be computed transforming $z_i/\sqrt{1-\sum_{i=1}^{n-1} |z_i|^2} \mapsto y_i$ so that $\chi(z) \mapsto (y_1, \dots, y_{n-1}, -1)/\sqrt{1-\sum_{i=1}^{n-1} |y_i|^2}$, $\chi(0) \mapsto (0, -1)$ and $e^i \mapsto e_i$ as in [26]. Then, using (3.45) on (3.47), one can obtain

$$\langle D^2 u(F(\xi)) \mathcal{W}_K^{-1}(\xi) e^i, \mathcal{W}_K^{-1}(\xi) e^j \rangle = -|\nabla u(F(\xi))| \langle \mathcal{W}_K^{-1}(\xi) e^i, e^j \rangle, \tag{3.48}$$

(see the end of [26, p. 382]). We successively multiply the above entrywise with the entries of $\mathcal{W}_K(F(\xi)) = \kappa(F(\xi)) \mathcal{C}[\nabla^2 h + h \mathbb{I}]$ recalling (3.42), then sum over to obtain (1).

Next, we differentiate the equation $|\nabla u(F(\chi))| = -\langle \nabla u(F(\chi)), \chi \rangle$ at $\xi = \chi(0)$ to obtain

$$\nabla_j(|\nabla u(F(\xi))|) = -\langle D^2 u(F(\xi)) \nabla_j F(\xi), \xi \rangle - \langle \nabla u(F(\xi)), e^j \rangle = -\langle D^2 u(F(\xi)) \mathcal{W}_K^{-1}(\xi) e^j, \xi \rangle,$$

as $\langle \nabla u(F(\xi)), e^j \rangle = -|\nabla u(F(\xi))| \langle \xi, e^j \rangle = 0$ from orthonormality. Multiplying with the entries of $\mathcal{W}_K(F(\xi)) = \kappa(F(\xi)) \mathcal{C}[\nabla^2 h + h \mathbb{I}]$ from (3.42) and summing over, we can similarly obtain (2).

Finally, since u is a solution of the Eq. (3.2), we have

$$0 = \operatorname{div}(|\nabla u|^{p-2} \nabla u) = |\nabla u|^{p-2} \left(\Delta u + \frac{(p-2)}{|\nabla u|^2} \langle D^2 u \nabla u, \nabla u \rangle \right),$$

in $\Omega \cap N$ and $u \in C^\infty(\Omega \cap N)$. Taking pointwise limit, together with (1), we get

$$\begin{aligned} (p-2) \langle D^2 u(F(\xi)) \xi, \xi \rangle &= -\Delta u(F(\xi)) = -\operatorname{Tr}(D^2 u(F(\xi))) \\ &= -\sum_i \langle D^2 u(F(\xi)) e^i, e^i \rangle - \langle D^2 u(F(\xi)) \xi, \xi \rangle \\ &= \kappa(F(\xi)) |\nabla u(F(\xi))| \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h \mathbb{I}]) - \langle D^2 u(F(\xi)) \xi, \xi \rangle, \end{aligned}$$

which yields (3). Thus the proof is finished. □

Now we obtain the first variation of the density of p -harmonic measures in the following which has been shown previously for a special case of $p = 2$ by Jerison [26, Proposition 2].

Proposition 3.49 *Given support functions $h = h_\Omega$ and $v = h_{\Omega_0}$, the first variation, defined as $\mathcal{L}_h[v] = \frac{d}{dt} \Big|_{t=0} \mathcal{F}[h + tv]$ as in (2.57) with \mathcal{F} as in (3.11), is given by*

$$\begin{aligned} \mathcal{L}_h[v] &= \sum_{i,j=1}^{n-1} \nabla_j \left(C_{i,j} [\nabla^2 h + h\mathbb{I}] |\nabla u|^{p-1} \nabla_i v \right) \\ &\quad - \frac{(p-1)}{\kappa} |\nabla u|^{p-2} \Lambda_u [|\nabla u|(v \circ \mathbf{g}_\Omega)], \end{aligned} \tag{3.50}$$

where $C_{i,j}[\cdot] = \langle C[\cdot]e^j, e^i \rangle$ are entries of the cofactor matrix as in (3.41) for $i, j \in \{1, \dots, n-1\}$ with respect to a local orthonormal frame $\{e^i\}$ and Λ_u is as in (3.31).

Proof It is standard as shown in [13], that

$$\frac{d}{dt} \Big|_{t=0} \det(\nabla^2 h + h\mathbb{I} + t(\nabla^2 v + v\mathbb{I})) = \text{Tr}(C[\nabla^2 h + h\mathbb{I}](\nabla^2 v + v\mathbb{I})).$$

Recalling $\mathcal{F}[h + tv](\xi) = |\nabla u(F(\xi, t), t)|^{p-1} \det(\nabla^2 h + h\mathbb{I} + t(\nabla^2 v + v\mathbb{I}))$ and using the above, note that

$$\begin{aligned} \mathcal{L}_h[v] &= |\nabla u(F(\xi))|^{p-1} \frac{d}{dt} \Big|_{t=0} \det(\nabla^2 h + h\mathbb{I} + t(\nabla^2 v + v\mathbb{I})) \\ &\quad + \det(\nabla^2 h + h\mathbb{I}) \frac{d}{dt} \Big|_{t=0} |\nabla u(F(\xi, t), t)|^{p-1} \\ &= |\nabla u(F(\xi))|^{p-1} \text{Tr}(C[\nabla^2 h + h\mathbb{I}](\nabla^2 v + v\mathbb{I})) \\ &\quad + (p-1) |\nabla u(F(\xi))|^{p-2} \det(\nabla^2 h + h\mathbb{I}) \frac{d}{dt} \Big|_{t=0} |\nabla u(F(\xi, t), t)|, \end{aligned} \tag{3.51}$$

and we need to compute only the last term of the above. Notice that, from the boundary condition of (3.10), the outer unit normal at $F(\xi, t) \in \partial\Omega^t$ is given by

$$\xi = \mathbf{g}_{\Omega^t}(F(\xi, t)) = -\frac{\nabla u(F(\xi, t), t)}{|\nabla u(F(\xi, t), t)|},$$

hence, $|\nabla u(F(\xi, t), t)| = -\langle \nabla u(F(\xi, t), t), \xi \rangle$. Therefore, we obtain

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} |\nabla u(F(\xi, t), t)| &= -\frac{d}{dt} \Big|_{t=0} \langle \nabla u(F(\xi, t), t), \xi \rangle \\ &= -\left(\langle D^2 u(F(\xi)) \dot{F}(\xi), \xi \rangle + \langle \nabla \dot{u}(F(\xi)), \xi \rangle \right) = -(I_1 + I_2). \end{aligned} \tag{3.52}$$

Since, $\dot{F}(\xi) = \nabla v(\xi)$ and recalling (2.28), we have $\nabla v(\xi) = v(\xi)\xi + \sum_{i=1}^{n-1} \nabla_i v(\xi)e^i$. Using this together with Lemma 3.44 we compute the first term of (3.52) as

$$\begin{aligned} I_1 &= \langle D^2 u(F(\xi))\xi, \xi \rangle v(\xi) + \sum_{i=1}^{n-1} \langle D^2 u(F(\xi))\xi, e^i \rangle \nabla_i v(\xi) \\ &= \frac{1}{(p-1)} \kappa(F(\xi)) |\nabla u(F(\xi))| v(\xi) \text{Tr}(C[\nabla^2 h + h\mathbb{I}]) \\ &\quad - \kappa(F(\xi)) \sum_{i,j=1}^{n-1} C_{i,j} [\nabla^2 h + h\mathbb{I}] \nabla_j (|\nabla u(F(\xi))|) \nabla_i v(\xi). \end{aligned}$$

It has been shown in [13] that $\sum_j \nabla_j C_{i,j}[\nabla^2 h + h\mathbb{I}] = 0$ for any i and using this on the latter term of the above, we obtain

$$\begin{aligned}
 I_1 &= -\kappa(F(\xi)) \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|) \nabla_i v(\xi) \\
 &\quad + \frac{1}{(p-1)} \kappa(F(\xi)) |\nabla u(F(\xi))| v(\xi) \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h\mathbb{I}]).
 \end{aligned}
 \tag{3.53}$$

Using (3.53) on (3.52), we obtain

$$\begin{aligned}
 \frac{d}{dt} \Big|_{t=0} |\nabla u(F(\xi, t), t)| &= \kappa(F(\xi)) \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|) \nabla_i v(\xi) \\
 &\quad - \frac{\kappa(F(\xi))}{(p-1)} |\nabla u(F(\xi))| v(\xi) \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h\mathbb{I}]) - \langle \nabla \dot{u}(F(\xi)), \xi \rangle,
 \end{aligned}
 \tag{3.54}$$

Recalling $\det(\nabla^2 h + h\mathbb{I}) = 1/\kappa(F(\xi))$ from (2.30), we use (3.54) in (3.51) to get

$$\begin{aligned}
 \mathcal{L}_h[v] &= |\nabla u(F(\xi))|^{p-1} \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h\mathbb{I}](\nabla^2 v + v\mathbb{I})) \\
 &\quad + (p-1) |\nabla u(F(\xi))|^{p-2} \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|) \nabla_i v(\xi) \\
 &\quad - |\nabla u(F(\xi))|^{p-1} v(\xi) \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h\mathbb{I}]) \\
 &\quad - (p-1) \frac{|\nabla u(F(\xi))|^{p-2}}{\kappa(F(\xi))} \langle \nabla \dot{u}(F(\xi)), \xi \rangle.
 \end{aligned}
 \tag{3.55}$$

Now, we use the following identity to replace the second term of the above,

$$\begin{aligned}
 &\sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|^{p-1} \nabla_i v(\xi)) \\
 &= \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|^{p-1}) \nabla_i v(\xi) \\
 &\quad + \sum_{i,j=1}^{n-1} C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|^{p-1} \nabla_{j,i} v(\xi) \\
 &= (p-1) |\nabla u(F(\xi))|^{p-2} \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|) \nabla_i v(\xi) \\
 &\quad + |\nabla u(F(\xi))|^{p-1} \operatorname{Tr}(\mathcal{C}[\nabla^2 h + h\mathbb{I}]\nabla^2 v).
 \end{aligned}
 \tag{3.56}$$

Using (3.56) on (3.55), we observe that the first and third term in (3.55) get cancelled off and we are left with

$$\begin{aligned}
 \mathcal{L}_h[v] &= \sum_{i,j=1}^{n-1} \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|^{p-1} \nabla_i v(\xi)) \\
 &\quad - (p-1) \frac{|\nabla u(F(\xi))|^{p-2}}{\kappa(F(\xi))} \langle \nabla \dot{u}(F(\xi)), \xi \rangle,
 \end{aligned}$$

which completes the proof since $\Lambda_u[|\nabla u|(v \circ \mathbf{g}_\Omega)](F(\xi)) = \langle \nabla \dot{u}(F(\xi)), \xi \rangle$ from (3.33). \square

The following shows the self-adjointness of the first variation in $L^2(\mathbb{S}^{n-1}, \mathcal{H}^{n-1})$.

Corollary 3.57 *The first variation \mathcal{L}_h being as above, for any $v_1, v_2 \in C(\mathbb{S}^{n-1})$, we have*

$$\int_{\mathbb{S}^{n-1}} v_1 \mathcal{L}_h[v_2] d\xi = \int_{\mathbb{S}^{n-1}} v_2 \mathcal{L}_h[v_1] d\xi. \tag{3.58}$$

Proof From Proposition 3.49 and (2.30), we have

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} v_1(\xi) \mathcal{L}_h[v_2](\xi) d\xi \\ &= \int_{\mathbb{S}^{n-1}} \sum_{i,j=1}^{n-1} v_1(\xi) \nabla_j (C_{i,j}[\nabla^2 h + h\mathbb{I}]|\nabla u(F(\xi))|^{p-1} \nabla_i v_2(\xi)) d\xi \\ & \quad - (p-1) \int_{\mathbb{S}^{n-1}} v_1(\xi) \Lambda_u[|\nabla u|(v_2 \circ \mathbf{g}_\Omega)](F(\xi)) \frac{|\nabla u(F(\xi))|^{p-2}}{\kappa(F(\xi))} d\xi \\ &= J_1 + J_2. \end{aligned}$$

Note that, using integral by parts (Stoke’s theorem for compact manifold without boundary) on the first term of the above, we have

$$J_1 = - \int_{\mathbb{S}^{n-1}} \sum_{i,j=1}^{n-1} |\nabla u(F(\xi))|^{p-1} C_{i,j}[\nabla^2 h + h\mathbb{I}] \nabla_i v_2(\xi) \nabla_j v_1(\xi) d\xi$$

which is symmetric in v_1 and v_2 . For the second term, using (2.30) and (2.31), we have

$$J_2 = -(p-1) \int_{\partial\Omega} (v_1 \circ \mathbf{g}_\Omega) \Lambda_u[|\nabla u|(v_2 \circ \mathbf{g}_\Omega)] |\nabla u|^{p-2} d\mathcal{H}^{n-1},$$

which is also symmetric in v_1 and v_2 from self-adjointness (3.39) of Λ_u from Proposition 3.38. Combining both cases of the above, the proof is finished. \square

3.2 Convergence of p -harmonic measures

The goal here is to establish weak convergence of p -harmonic measures with respect to the Hausdorff distance. Given a convex domain $\Omega \subset \mathbb{R}^n$, a neighborhood N of $\partial\Omega$ and $u \in W^{1,p}(\Omega \cap N)$ as in (3.2), we consider a sequence of convex domains $\Omega_j \subset \mathbb{R}^n$ such that $d_{\mathcal{H}}(\Omega_j, \Omega) \rightarrow 0^+$ as $j \rightarrow \infty$; recalling (2.43), $d_{\mathcal{H}}(\partial\Omega_j, \partial\Omega) \rightarrow 0^+$.

First, we have the following lemma due to Jerison [26].

Lemma 3.59 *Given $\Omega \subset \mathbb{R}^n$ and $d_{\mathcal{H}}(\Omega_j, \Omega) \rightarrow 0^+$ as $j \rightarrow \infty$, if $0 \in \Omega \cap \Omega_j$ for all $j \geq j_0$ for some $j_0 \in \mathbb{N}$, then there exists $r > 0$ such that, up to a subsequence, $(1 - 1/j)\Omega_j/r \subset \Omega/r$.*

Proof Note that $|\text{dist}(0, \partial\Omega_j) - \text{dist}(0, \partial\Omega)| \leq d_{\mathcal{H}}(\partial\Omega_j, \partial\Omega) = d_{\mathcal{H}}(\Omega_j, \Omega) \leq 1/j$, for all $j \geq j_0$ with $j_0 \in \mathbb{N}$ large enough. Hence, we choose

$$0 < r < \min\{\text{dist}(0, \partial\Omega), \inf_{j \geq j_0} \text{dist}(0, \partial\Omega_j)\}$$

so that $B_r(0) \subset \Omega_j \cap \Omega$ for all $j \geq j_0$. Let $\tilde{\Omega} = \Omega/r$ and $\tilde{\Omega}_j = \Omega_j/r$. Hence $B_1(0) \subset \tilde{\Omega}_j \cap \tilde{\Omega}$ for all $j \geq j_0$ and $d_{\mathcal{H}}(\tilde{\Omega}_j, \tilde{\Omega}) \rightarrow 0^+$. Thus, $1 = h_{B_1(0)} \leq \min\{h_{\tilde{\Omega}_j}, h_{\tilde{\Omega}}\}$ and recalling (2.43), we have the convergence of support functions. Hence, up to a subsequence, $1 \leq h_{\tilde{\Omega}_j} \leq$

$h_{\bar{\Omega}} + 1/j$ for all $j \geq j_0$ which implies $(1 - 1/j)h_{\bar{\Omega}_j} \leq h_{\bar{\Omega}} - (h_{\bar{\Omega}} - 1)/j - 1/j^2 \leq h_{\bar{\Omega}}$, and the proof is complete. \square

Note that $\lim_{j \rightarrow \infty} d_{\mathcal{H}}((1 - 1/j)\Omega_j, \Omega) = \lim_{j \rightarrow \infty} d_{\mathcal{H}}(\Omega_j, \Omega) = 0$. Henceforth, in view of Lemma 3.59, we shall assume without loss of generality that

$$0 \in \Omega \cap \Omega_j \quad \text{and} \quad \Omega_j \subset \Omega, \quad \forall j \geq j_0, \tag{3.60}$$

for a large enough $j_0 \in \mathbb{N}$. We can also similarly assume without loss of generality that $\Omega \subset \Omega_j(1 + 1/j)$ for all $j \geq j_0$. Furthermore, since $d_{\mathcal{H}}(\partial\Omega_j, \partial\Omega) \rightarrow 0^+$, for any sequence of neighborhoods N_j of $\partial\Omega_j$ we have $d_{\mathcal{H}}(N_j, N) \rightarrow 0^+$ as $j \rightarrow \infty$. We can assume without loss of generality that N is a convex ring (see [34], e.g. annulus) so that the outer and inner segments of ∂N enclose convex domains. Hence, in view of (2.43) and Lemma 3.59, we can regard that being subject to a possible small extension and dilation, N is also a neighborhood of $\partial\Omega_j$ so that $\partial N \cap \Omega_j = \partial N \cap \Omega$ for all $j \geq j_0$, which shall be assumed hereafter.

There exists a unique $u_j \in W^{1,p}(\Omega_j \cap N)$ which is the weak solution of the Dirichlet problem

$$\begin{cases} \operatorname{div}(|\nabla u_j|^{p-2} \nabla u_j) = 0, & \text{in } \Omega_j \cap N; \\ u_j = 0, & \text{in } \partial\Omega_j \cap N; \\ u_j = u, & \text{in } \partial N \cap \Omega_j; \end{cases} \tag{3.61}$$

for all $j \geq j_0$ and upon zero extension, $u_j \in W^{1,p}(N)$. We consider the p -harmonic measure with respect to u_j as $\omega_{p,j}$ so that $d\omega_{p,j} = |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega_j}$.

Remark 3.62 In view of the above, given $d_{\mathcal{H}}(\Omega_j, \Omega) \rightarrow 0^+$ as $j \rightarrow \infty$, we can assume without loss of generality that $\Omega_j(1 - 1/j) \subset \Omega \subset \Omega_j(1 + 1/j)$ for $j \geq j_0$ large enough. Hence, (3.61) can be re-defined with dilations on Ω as $\Omega/(1 + 1/j)$ or $\Omega/(1 - 1/j)$ for which points in Ω are transformed as $x \mapsto x/(1 + 1/j)$ and $x \mapsto x/(1 - 1/j)$ respectively. In this sense, the approximants defined by (3.61) can be regarded as a generalization of (3.10) for $t > 0$ or $t < 0$, since for $\Omega^t = \Omega + t\Omega_0$, recall that $d_{\mathcal{H}}(\Omega^t, \Omega) = |t| \|h_{\Omega_0}\|_{L^\infty(\mathbb{S}^{n-1})} \rightarrow 0^+$ as $t \rightarrow 0$. All the following results corresponding to (3.61) remain unaffected with respect to these dilations.

Towards the weak convergence of p -harmonic measures, we make use of the following geometric lemma due to Jerison [27, Lemma 3.3], which says that most of the boundaries of convex domains close enough by Hausdorff distance, can be locally flattened, except for a small measure set. It can be shown using convexity and Vitali covering theorem.

Lemma 3.63 *For a convex domain Ω , given any $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$ and a finite collection of disjoint balls $B_{r_k}(x_k)$ for $1 \leq k \leq N_0$, with $r_k < \varepsilon$ and $x_k \in \partial\Omega$, such that for every Ω' with $d_{\mathcal{H}}(\Omega', \Omega) < \delta$, we have*

$$\mathcal{H}^{n-1} \left(\partial\Omega \setminus \bigcup_{k=1}^{N_0} \bar{B}_{r_k}(x_k) \right) < \varepsilon,$$

and after a possible rotation and translation, $\partial\Omega, \partial\Omega'$ are graphs of functions $\phi, \phi' \in C^{0,1}(\mathbb{R}^{n-1})$ in $B_{r_k/\varepsilon}(x_k)$, satisfying $|\nabla\phi(x)| + |\nabla\phi'(x)| \leq \varepsilon$ for all $x \in \mathbb{R}^n$ with $|x| < r_k/\varepsilon$.

The following is a part of [27, Lemma 3.7] and can be shown using the density of radial projection in (2.42), i.e. $d\mathcal{H}^{n-1} = |\rho_\Omega(\theta)|^n / h_\Omega(\mathbf{g}_\Omega(\rho_\Omega(\theta))) d\theta$ on Lemma 3.63.

Corollary 3.64 *For a convex domain Ω , given any $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$, $s_0 > 0$ and a family of balls \mathcal{B} on \mathbb{S}^{n-1} such that the following holds:*

- (1) *Every $B \in \mathcal{B}$ has radius s_0 .*
- (2) *If $B_r(0) \subset \Omega \subset B_R(0)$, then there exists $N_0 > 0$ depending on R/r such that any point in \mathbb{S}^{n-1} belongs to atmost N_0 balls of \mathcal{B} .*
- (3) *$\theta(\mathbb{S}^{n-1} \setminus F) < \varepsilon$ where $F = \bigcup_{B \in \mathcal{B}} B$ and $\theta = \mathcal{H}^{n-1} \llcorner_{\mathbb{S}^{n-1}}$ is the uniform measure.*

Now, we sketch the weak convergence of p -harmonic measures in the following.

Proposition 3.65 *Given a bounded convex domain $\Omega \subset \mathbb{R}^n$, for any sequence of convex domains $\Omega_j \subset \mathbb{R}^n$ with $d_{\mathcal{H}}(\Omega_j, \Omega) \rightarrow 0^+$ as $j \rightarrow \infty$, if $\omega_{p,j}$ is the p -harmonic measure with respect to the solution $u_j \in W^{1,p}(N)$ of (3.61), then up to a subsequence, $\nabla u_j \rightarrow \nabla u$ uniformly in N and we have the weak convergence $\omega_{p,j} \rightarrow \omega_p$ as $j \rightarrow \infty$, i.e. for any $\eta \in C(N)$, we have*

$$\lim_{j \rightarrow \infty} \int_{\partial\Omega_j} \eta |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} = \int_{\partial\Omega} \eta |\nabla u|^{p-1} d\mathcal{H}^{n-1}. \tag{3.66}$$

Proof From boundary conditions of (3.61) and comparison principle, we have $u_j > 0$ in Ω_j and hence, $\mathbf{g}_{\Omega_j} = -\nabla u_j / |\nabla u_j|$. Then, testing the Eq. (3.61) with any $\eta \in W^{1,p}(N)$, we have

$$\begin{aligned} & \int_{\Omega_j \cap N} |\nabla u_j|^{p-2} \langle \nabla u_j, \nabla \eta \rangle dx \\ &= \int_{\partial(\Omega_j \cap N)} \eta |\nabla u_j|^{p-2} \langle \nabla u_j, \nu_{\partial(\Omega_j \cap N)} \rangle d\mathcal{H}^{n-1} \\ &= \int_{\Omega_j \cap \partial N} \eta |\nabla u|^{p-2} \langle \nabla u, \nu_{\Omega_j \cap \partial N} \rangle d\mathcal{H}^{n-1} + \int_{\partial\Omega_j} \eta |\nabla u_j|^{p-1} d\mathcal{H}^{n-1}. \end{aligned}$$

Since $\partial N \cap \Omega_j = \partial N \cap \Omega$ for all $j \geq j_0$, we test the Eq. (3.2) with η and take the difference with the above to obtain

$$\begin{aligned} & \int_{\Omega_j \cap N} |\nabla u_j|^{p-2} \langle \nabla u_j, \nabla \eta \rangle dx - \int_{\Omega \cap N} |\nabla u|^{p-2} \langle \nabla u, \nabla \eta \rangle dx \\ &= - \left(\int_{\partial\Omega_j} \eta |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} - \int_{\partial\Omega} \eta |\nabla u|^{p-1} d\mathcal{H}^{n-1} \right). \end{aligned} \tag{3.67}$$

Now we show that $\nabla u_j \rightarrow \nabla u$ uniformly in N as $j \rightarrow \infty$. However, in this case $|\Omega \setminus \Omega_j|$ or $|\mathcal{H}^{n-1}(\partial\Omega) - \mathcal{H}^{n-1}(\partial\Omega_j)|$ are not generally dominated by $d_{\mathcal{H}}(\Omega_j, \Omega)$. Therefore, Lemma 3.63 and Corollary 3.64 can be used by taking radial projections of (2.42) and decompose the integral on \mathbb{S}^{n-1} into $F = \bigcup_{B \in \mathcal{B}} B$ and $\mathbb{S}^{n-1} \setminus F$ as in Corollary 3.64, then obtain for any $\varepsilon > 0$,

$$\left| \int_B |\nabla(u_j \circ \rho_j)|^{p-1} - |\nabla(u \circ \rho)|^{p-1} d\theta \right| \leq c\varepsilon \int_B |\nabla(u \circ \rho)|^{p-1} d\theta, \quad \forall j \geq j_0(\varepsilon), B \in \mathcal{B},$$

where $\rho_j \rightarrow \rho$ \mathcal{H}^{n-1} -a.e. in \mathbb{S}^{n-1} . This involves refined estimates of p -harmonic functions in [15, 31, 32], etc. used to obtain e.g. [15, Lemma 4.5] to which we refer to and omit the details here. The above together with $\theta(\mathbb{S}^{n-1} \setminus F) < \varepsilon$ from Corollary 3.64 allows us to estimate the integral on whole of \mathbb{S}^{n-1} and then using (2.45) and (2.42), we have

$$\left| \int_{\partial\Omega_j} |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} - \int_{\partial\Omega} |\nabla u|^{p-1} d\mathcal{H}^{n-1} \right| \leq c\varepsilon \|u\|_{W^{1,p}(N)}^{p-1}, \quad \forall j \geq j_0(\varepsilon).$$

An integral on $\Omega \setminus \Omega_j$ can be similarly estimated using (2.40), Corollary 3.64 since $r_{\Omega_j} \rightarrow r_\Omega$ uniformly as $j \rightarrow \infty$. Using these estimates together with $\eta = u_j - u$ on (3.67) and using the monotonicity (2.49), we can conclude

$$\int_{B_r} |\nabla u_j - \nabla u|^p dx \leq c\varepsilon \|u\|_{W^{1,p}(N)}^p, \quad \forall j \geq j_0(\varepsilon),$$

for any $B_r \subset \Omega_j \cap N$. The continuity of the gradients $\nabla u_j, \nabla u$ from (2.50) implies that all points are Lebesgue points and hence the above is enough to conclude the following pointwise convergence of $\nabla u_j - \nabla u$; the estimate of (2.50) for ∇u_j shows equicontinuity of the family $\{\nabla u_j\}_{j \geq j_0}$ and uniform convergence $\nabla u_j \rightarrow \nabla u$ in N as $j \rightarrow \infty$ follows from the Arzelà-Ascoli theorem. This used in (3.67) directly leads us to conclude

$$\lim_{j \rightarrow \infty} \int_{\partial \Omega_j} \eta |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} = \int_{\partial \Omega} \eta |\nabla u|^{p-1} d\mathcal{H}^{n-1}$$

for any $\eta \in C^1(\bar{N})$ and hence also for any $\eta \in C(\bar{N})$ by taking approximations. The proof is finished. \square

Remark 3.68 The proof of weak convergence of p -harmonic measures above is easier compared to other measures with a higher exponent on the gradient because in our case, the exponent $(p - 1)$ is precisely what appears in the weak form of the equations and so we do not require a reverse Hölder inequality as in [15]. It is so remarked in [27] that the weak convergence in case of harmonic measure in [26] is easier than that of capacitary measures in [27] for $p = 2$.

Let $\mu_{\Omega_j} = (\mathbf{g}_{\Omega_j})_* \omega_{p,j}$ be the corresponding measure as an approximation to μ_Ω . As $\mathbf{g}_{\Omega_j} \rightarrow \mathbf{g}_\Omega$ \mathcal{H}^{n-1} -a.e. we have the weak convergence $\mu_{\Omega_j} \rightarrow \mu_\Omega$ as a consequence of the above. Thus,

$$\lim_{j \rightarrow \infty} \int_{\mathbb{S}^{n-1}} w d\mu_{\Omega_j} = \int_{\mathbb{S}^{n-1}} w d\mu_\Omega, \quad \forall w \in C(\mathbb{S}^{n-1}). \tag{3.69}$$

More generally, we consider $u = u_K \in W^{1,p}(N)$ for a general convex set K and a neighborhood N of ∂K , where either u satisfies (3.2) if $K = \bar{\Omega}$ or u satisfies (3.4) if $K = \partial K$. Then ω_p defined by $d\omega_p = |\nabla u|^{p-1} d\mathcal{H}^{n-1}$ on ∂K and μ_K is as in (3.5) and defined with respect to (2.35) and (2.36) in general. We consider $\{K_j\}$ such that $d_{\mathcal{H}}(K_j, K) \rightarrow 0^+$. If $K_j = \bar{\Omega}_j$ are of non-empty interior then we define $u_j \in W^{1,p}(\Omega_j \cap N)$ similarly as (3.61), otherwise if $K_j = \partial K_j$ and $K = \partial K$ then we define $u_j \in W^{1,p}(N)$ as the unique solution of

$$\begin{cases} \operatorname{div} (|\nabla u_j|^{p-2} \nabla u_j) = 0, & \text{in } N \setminus K_j; \\ u_j = 0, & \text{in } K_j \cap N; \\ u_j = u, & \text{in } \partial N \setminus K_j. \end{cases} \tag{3.70}$$

The p -harmonic measures are defined by $d\omega_{p,j} = |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial K_j}$ and the weak convergence $\omega_{p,j} \rightarrow \omega_p$ can be obtained similarly as Proposition 3.65. Therefore, the weak convergence for measures for the p -harmonic Minkowski problem on general convex sets,

$$\mu_{K_j} \rightarrow \mu_K, \quad \text{if } d_{\mathcal{H}}(K_j, K) \rightarrow 0^+, \tag{3.71}$$

when the measures are defined as in (3.5) with respect to (2.35) and (2.36), can also be obtained using the above and approximation of general convex sets by polytopes, in ways similar to that of (2.48). We refer to [45, Chapter 4,5] for details of such arguments.

In the following, given a convex set K , we denote the functional

$$\Gamma(K) := \int_{\mathbb{S}^{n-1}} h_K(\xi) d\mu_K(\xi) \tag{3.72}$$

and $\Gamma(\Omega) = \Gamma(K)$ if $\bar{\Omega} = K$. It is easy to see from (2.43) and (3.71), that $\Gamma(K_j) \rightarrow \Gamma(K)$ uniformly if $d_{\mathcal{H}}(K_j, K) \rightarrow 0^+$ as $j \rightarrow \infty$.

Remark 3.73 For a compact convex set K of non-empty interior note that

$$\Gamma(K) = \int_{\mathbb{S}^{n-1}} h_K(\xi) |\nabla u(\mathbf{g}_K^{-1}(\xi))|^{p-1} dS_K(\xi)$$

closely resembles the well known formula of volumes given by

$$|K| = \frac{1}{n} \int_{\mathbb{S}^{n-1}} h_K(\xi) dS_K(\xi).$$

Hence we can conclude that if $K \mapsto \mu_K$ is defined by a common p -harmonic function u for C^2_+ domains then Γ is a measure absolutely continuous with respect to the Lebesgue measure. For non-smooth domains or convex sets of empty interior, the measure Γ can be singular.

We complete this subsection with the following.

Lemma 3.74 *Given convex domains $\Omega_j, \Omega_{j,0} \subset \mathbb{R}^n$ with $d_{\mathcal{H}}(\Omega_j, \Omega), d_{\mathcal{H}}(\Omega_{j,0}, \Omega_0) \rightarrow 0^+$ as $j \rightarrow \infty$, let $\Omega^t = \Omega + t\Omega_0$ and $\Omega_j^t = \Omega_j + t\Omega_{j,0}$ for $|t| < \tau$ small enough. Then, let*

$$d\mu_{\Omega^t} = |\nabla u(\cdot, t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega^t},$$

where $u(\cdot, t)$ is the unique solution of the Dirichlet problem (3.10) and let $d\mu_{\Omega_j^t}$ be defined similarly; then with Γ as (3.72), if $t \mapsto \Gamma(\Omega_j^t)$ is differentiable at $t = 0$ for all $j \geq j_0$ then $t \mapsto \Gamma(\Omega^t)$ is also differentiable at $t = 0$ and we have

$$\lim_{j \rightarrow \infty} \frac{d}{dt} \Big|_{t=0} \Gamma(\Omega_j^t) = \frac{d}{dt} \Big|_{t=0} \Gamma(\Omega^t) = \frac{d}{dt} \Big|_{t=0} \lim_{j \rightarrow \infty} \Gamma(\Omega_j^t). \tag{3.75}$$

Proof From (2.16), we note that

$$\begin{aligned} \Gamma(\Omega_j^t) - \Gamma(\Omega_j) &= \int_{\mathbb{S}^{n-1}} h_{\Omega_j^t} d\mu_{\Omega_j^t} - \int_{\mathbb{S}^{n-1}} h_{\Omega_j} d\mu_{\Omega_j} \\ &= \int_{\mathbb{S}^{n-1}} h_{\Omega_j} d\mu_{\Omega_j^t} - \int_{\mathbb{S}^{n-1}} h_{\Omega_j} d\mu_{\Omega_j} + t \int_{\mathbb{S}^{n-1}} h_{\Omega_{j,0}} d\mu_{\Omega_j^t}. \end{aligned}$$

Recalling (2.43), we know that $h_{\Omega_j}, h_{\Omega_{j,0}} \rightarrow h_{\Omega}, h_{\Omega_0}$ uniformly as $j \rightarrow \infty$. Hence for each $t \neq 0$, we have $(\Gamma(\Omega_j^t) - \Gamma(\Omega_j))/t \rightarrow (\Gamma(\Omega^t) - \Gamma(\Omega))/t$ uniformly (independent of t) as $j \rightarrow \infty$. Also from (2.43), note that

$$d_{\mathcal{H}}(\Omega_j^t, \Omega_j) = |t| \|h_{\Omega_{j,0}}\|_{L^\infty} \quad \text{and} \quad d_{\mathcal{H}}(\Omega^t, \Omega) = |t| \|h_{\Omega_0}\|_{L^\infty},$$

which converge to zero uniformly (independent of j) as $t \rightarrow 0$ as well. From (3.69) and above, $\mu_{\Omega_j^t} \rightarrow \mu_{\Omega^t}$ as $j \rightarrow \infty$ for all t and $\mu_{\Omega_j^t} \rightarrow \mu_{\Omega_j}$ as $t \rightarrow 0$ for all $j \geq j_0$. This is enough to conclude the proof. □

3.3 Proof of existence

We shall prove the existence of p -harmonic measures prescribing measures on spheres in two steps; first, for discrete measures and then for the general case thereby leading to the proof of Theorem 1.2 in the end.

The following proposition resembles the Hadamard’s variational formula.

Proposition 3.76 *Given convex domains $\Omega^t = \Omega + t\Omega_0$ with $|t| < \tau$ small enough, $v = h_{\Omega_0} > 0$ and Γ as in (3.72) with $d\mu_{\Omega^t} = |\nabla u(\cdot, t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega^t}$ where $u(\cdot, t)$ is the unique solution of the Dirichlet problem (3.10), we have*

$$\frac{d}{dt} \Big|_{t=0} \Gamma(\Omega^t) = (n - p + 1) \int_{\mathbb{S}^{n-1}} v \, d\mu_{\Omega}. \tag{3.77}$$

Proof First, we assume that Ω, Ω_0 are of class $C_+^{2,\alpha}$. Then, recalling $d\mu_{\Omega^t}(\xi) = \mathcal{F}[h + tv](\xi) \, d\xi$ where $h_{\Omega} = h$, observe that

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \Gamma(\Omega^t) &= \frac{d}{dt} \Big|_{t=0} \int_{\mathbb{S}^{n-1}} h_{\Omega^t}(\xi) \, d\mu_{\Omega^t}(\xi) = \frac{d}{dt} \Big|_{t=0} \int_{\mathbb{S}^{n-1}} (h + tv) \mathcal{F}[h + tv](\xi) \, d\xi \\ &= \int_{\mathbb{S}^{n-1}} v(\xi) \mathcal{F}[h](\xi) \, d\xi + \int_{\mathbb{S}^{n-1}} h(\xi) \frac{d}{dt} \Big|_{t=0} \mathcal{F}[h + tv](\xi) \, d\xi \\ &= \int_{\mathbb{S}^{n-1}} v(\xi) \, d\mu_{\Omega}(\xi) + \int_{\mathbb{S}^{n-1}} h(\xi) \mathcal{L}_h[v](\xi) \, d\xi \\ &= \int_{\mathbb{S}^{n-1}} v(\xi) \, d\mu_{\Omega}(\xi) + \int_{\mathbb{S}^{n-1}} v(\xi) \mathcal{L}_h[h](\xi) \, d\xi, \end{aligned} \tag{3.78}$$

where the self-adjointness of Corollary 3.57 is used to obtain the last equality of the above. Now, recalling (3.13) of Lemma 3.12, we have $\mathcal{L}_h[h](\xi) = (n - p) \mathcal{F}[h](\xi)$. Using this on (3.78), we obtain

$$\frac{d}{dt} \Big|_{t=0} \Gamma(\Omega^t) = (n - p + 1) \int_{\mathbb{S}^{n-1}} v(\xi) \, d\mu_{\Omega}(\xi),$$

for the case of $C_+^{2,\alpha}$ domains. The general case follows from approximation of convex domains by $C_+^{2,\alpha}$ domains using Theorem 2.46 and (2.47) along with (3.69) and (3.75). \square

In the following lemma, we consider the possibility of having a lower dimensional convex set in order to be able to assert the existence theorem for convex sets of non-empty interior.

Lemma 3.79 *Given a compact convex set K and a measure μ on \mathbb{S}^{n-1} , we have the following:*

- (1) *If $\dim_{\mathcal{H}}(K) = n - 1$, then K is contained in a hyperplane; precisely, there exists $\xi_0 \in \mathbb{S}^{n-1}$ and $c \in \mathbb{R}$ such that $K \subset \{x : \langle x, \xi_0 \rangle = c\}$ and μ_K as in (3.5) is given by*

$$\mu_K = \left(\int_K |\nabla u|^{p-1} d\mathcal{H}^{n-1} \right) (\delta_{\xi_0} + \delta_{-\xi_0}).$$

- (2) *If $\mu = a(\delta_{\xi_0} + \delta_{-\xi_0})$ for some $\xi_0 \in \mathbb{S}^{n-1}$ and $a > 0$, then μ does not satisfy (i) of (1.1).*

Proof To prove (1), recalling (2.12) since $K = \bigcap_{\xi \in \mathbb{S}^{n-1}} H_{\xi, h_K}^-$, observe that $\dim_{\mathcal{H}}(K) = n - 1$ can only occur if at least two half-spaces share a common boundary and K is degenerated to a 1-codimensional subset of a hyperplane, i.e. $H_K(\xi) = H_K(\xi')$ for some $\xi \neq \xi' \in \mathbb{S}^{n-1}$

which can happen only if $\xi' = -\xi$ and $h_K(\xi') = -h_K(\xi)$. These together imply $h_K(-\xi) = -h_K(\xi)$; but, from the definition of support function, since

$$-h_K(-\xi) \leq \langle x, \xi \rangle \leq h_K(\xi), \quad \forall x \in K, \xi \in \mathbb{S}^{n-1},$$

therefore equality has to hold. Thus, we have $K \subset \{x : \langle x, \xi_0 \rangle = c\}$ where $\xi_0 = \xi$ and $c = h_K(\xi)$. Then, we observe that, in the sense of (2.35), we have $\mathbf{g}_K(x) = \{\xi_0, -\xi_0\}$ for \mathcal{H}^{n-1} -a.e. $x \in K$. Precisely, if $F \subset K$ is the lower dimensional boundary of K within the hyperplane then $\mathbf{g}_K(x) = \{\xi_0, -\xi_0\}$ for all $x \in K \setminus F$ and $\mathcal{H}^{n-1}(F) = 0$. Also in this case $K = \partial K$. Hence, for any measurable $E \subseteq \mathbb{S}^{n-1}$, note that, $g_K^{-1}(E)$ in the sense of (2.36) is of full \mathcal{H}^{n-1} -measure only if $\xi_0 \in E$ or $-\xi_0 \in E$; to be precise, letting $F' = \{x \in F : \mathbf{g}_K(x) \cap E \neq \emptyset\} \subseteq F$, we have that $g_K^{-1}(E) = (K \setminus F) \cup F'$ if $\{\xi_0, -\xi_0\} \cap E \neq \emptyset$ and otherwise $g_K^{-1}(E) = F'$. Thus,

$$S_K = \mathcal{H}^{n-1}(K)(\delta_{\xi_0} + \delta_{-\xi_0}) \quad \text{and} \quad \mu_K = \left(\int_K |\nabla u|^{p-1} d\mathcal{H}^{n-1} \right) (\delta_{\xi_0} + \delta_{-\xi_0}),$$

from (3.5), which completes the first part.

To prove (2), we observe that if $\mu = a(\delta_{\xi_0} + \delta_{-\xi_0})$ for some $\xi_0 \in \mathbb{S}^{n-1}$ and $a > 0$, then

$$\int_{\mathbb{S}^{n-1}} |\langle \zeta, \xi \rangle| d\mu(\xi) = a(|\langle \zeta, \xi_0 \rangle| + |\langle \zeta, -\xi_0 \rangle|) = 2a|\langle \zeta, \xi_0 \rangle|,$$

which vanishes for any $\zeta \in \mathbb{S}^{n-1}$ normal to ξ_0 . This completes the second part and the proof. \square

The following theorem is a discrete version of Theorem 1.2, which is an essential step towards the general existence theorem. Here we require a technical assumption of an antipodal condition and $p \neq n + 1$, which shall be ultimately removed.

Theorem 3.80 *Let μ be a discrete positive Borel measure on \mathbb{S}^{n-1} satisfying conditions (1.1) and has no antipodal pair of point masses i.e. if $\mu(\{\xi\}) > 0$ for some $\xi \in \mathbb{S}^{n-1}$ then $\mu(\{-\xi\}) = 0$. Then there exists a polytope P with non-empty interior such that $\mu_P = \mu$ and $\Gamma(P) = 1$ with $\mu_P = (\mathbf{g}_P)_* \omega_p$ as in (3.5), where ω_p is any p -harmonic measure on ∂P and Γ is as in (3.72), for any $1 < p < \infty$ with $p \neq n + 1$.*

Proof By virtue of the duality relation of Theorem 2.38 between convex domains and positive continuous functions on \mathbb{S}^{n-1} , the existence of the polytope P shall be obtained from existence of minimizer of the following constrained minimization problem

$$b = \inf \left\{ \int_{\mathbb{S}^{n-1}} h d\mu : h \in C_+(\mathbb{S}^{n-1}), \Gamma(\Omega_h) \geq 1 \right\}, \tag{3.81}$$

where Ω_h is the Wulff shape of h as in (2.37), hence $\bar{\Omega}_h = \{x \in \mathbb{R}^n : \langle x, \xi \rangle \leq h(\xi) \forall \xi \in \mathbb{S}^{n-1}\}$, and Γ is as in (3.72) for a suitably defined measure μ_{Ω_h} , to be mentioned below.

Since μ is discrete, there exists $\xi_1, \dots, \xi_m \in \mathbb{S}^{n-1}$ such that we have

$$\mu = \sum_{i=1}^m c_i \delta_{\xi_i}$$

where, the condition (1.1) implies for all $\xi \in \mathbb{S}^{n-1}$, we have

$$\sum_{i=1}^m c_i |\langle \xi, \xi_i \rangle| > 0 \quad \text{and} \quad \sum_{i=1}^m c_i \xi_i = 0, \tag{3.82}$$

and in particular, $c_1, \dots, c_m > 0$. Let us define $P_1 = \{x \in \mathbb{R}^n : \langle x, \xi_i \rangle \leq h(\xi_i) \forall i \in \{1, \dots, m\}\}$ for some $h \in C_+(\mathbb{S}^{n-1})$, which is a polytope from Theorem 2.18. We note that $\bar{\Omega}_h \subseteq P_1$ and $h_{P_1}(\xi_i) = \sup_{x \in P_1} \langle x, \xi_i \rangle \leq h(\xi_i)$ from the definition of P_1 and $\mathcal{H}^{n-1}(\partial\Omega_h) \leq \mathcal{H}^{n-1}(\partial P_1)$. Thus, we have

$$\int_{\mathbb{S}^{n-1}} h_{P_1} d\mu = \sum_{i=1}^m c_i h_{P_1}(\xi_i) \leq \sum_{i=1}^m c_i h(\xi_i) = \int_{\mathbb{S}^{n-1}} h d\mu. \tag{3.83}$$

We also note that $0 \in \Omega_h$ if $h > 0$ and from Theorem 2.38, for \mathbb{S}^{Ω_h} -a.e. $\xi \in \mathbb{S}^{n-1}$, we have $h(\xi) = h_{\Omega_h}(\xi) \leq h_{P_1}(\xi)$ that may or may not include the set $\{\xi_1, \dots, \xi_m\}$.

For any bounded sequence $h_j \in C_+(\mathbb{S}^{n-1})$, let us denote the corresponding sequence of polytopes as above, i.e. $P_j = \{x \in \mathbb{R}^n : \langle x, \xi_i \rangle \leq h_j(\xi_i) \forall i \in \{1, \dots, m\}\}$. Then, $\bar{\Omega}_{h_j} \subseteq P_j$ and (3.83) leads to

$$\sum_{i=1}^m c_i h_{P_j}(\xi_i) \leq \limsup_{j \rightarrow \infty} \int_{\mathbb{S}^{n-1}} h_j d\mu =: M < \infty$$

and since $c_i > 0$, this implies for all $x \in P_j$ we have $\langle x, \xi_i \rangle \leq h_{P_j}(\xi_i) \leq M / (\min_{1 \leq i \leq m} c_i)$. Thus, we can conclude that the sequence of polytopes $\{P_j\}$ is bounded and hence, from Blaschke selection theorem (Theorem 2.44), there exists P such that, up to subsequence, $d_{\mathcal{H}}(P_j, P) \rightarrow 0^+$ as $j \rightarrow \infty$. From Theorem 2.18, P is also a polytope with normals ξ_1, \dots, ξ_m .

Let $u = u_P$ be the p -harmonic function corresponding to any fixed p -harmonic measure of P , defined in the neighborhood of ∂P that contains ∂P_j for all $j \geq j_0$, since from (2.43) $d_{\mathcal{H}}(\partial P_j, \partial P) \rightarrow 0^+$ as $j \rightarrow \infty$. With respect to this u , we choose u_j uniquely as solutions of the Dirichlet problem (3.61) as in the previous subsection, so that we have $\mu_{P_j} \rightarrow \mu_P$ from (3.71). We shall define $\mu_{\Omega_{h_j}}$ with respect to u_j similarly as μ_{P_j} on neighborhoods of ∂P_j that contain $\partial\Omega_{h_j}$ for $j \geq j_0$ large enough. In other words, we have defined

$$\mu_{P_j} = (\mathbf{g}_{P_j})_* (|\nabla u_j|^{p-1} \mathcal{H}^{n-1} \llcorner \partial P_j) \quad \text{and} \quad \mu_{\Omega_{h_j}} = (\mathbf{g}_{\Omega_{h_j}})_* (|\nabla u_j|^{p-1} \mathcal{H}^{n-1} \llcorner \partial\Omega_{h_j}).$$

Now, let $h_j \in C_+(\mathbb{S}^{n-1})$ be a minimizing sequence of the infima (3.81) where Γ is as in (3.72) with $\mu_{\Omega_{h_j}}$ as above. Since $\mu_{P_j} \rightarrow \mu_P$, we have hence $\Gamma(P_j) \rightarrow \Gamma(P)$ as $j \rightarrow \infty$ and since $\Omega_{h_j} \subseteq P_j$, hence from (3.81) note that $1 \leq \Gamma(\Omega_{h_j}) \leq \Gamma(P_j)$ and therefore,

$$\int_{\mathbb{S}^{n-1}} h_P d\mu_P = \Gamma(P) = \lim_{j \rightarrow \infty} \Gamma(P_j) \geq 1.$$

Suppose, P is of empty interior, hence $P = \partial P$ and Hausdorff dimension $\dim_{\mathcal{H}}(P) < n$. Notice that the above enforces $\dim_{\mathcal{H}}(P) \geq n - 1$ (otherwise, $\mathcal{H}^{n-1}(P) = 0 = \mathcal{H}^{n-1}(\partial P)$ which, from (3.5), implies $\mu_P \equiv 0$ and hence $\Gamma(P) = 0$ contradicting the above). Then, $\dim_{\mathcal{H}}(P) = n - 1$. As in (1) of Lemma 3.79, this can occur only if at least two half-spaces of P share a common boundary and degenerate the whole polytope to a 1-codimensional face; if those are the half spaces corresponding to $\xi_i \neq \xi_j \in \mathbb{S}^{n-1}$, it enforces $\xi_j = -\xi_i$. But since $c_i, c_j > 0$, this contradicts the antipodal condition. Thus, P is of non-empty interior.

Therefore, let $P_0 \neq \emptyset$ be the interior of P and hence $\Gamma(P_0) \geq 1$, h_{P_0} is a minimizer of (3.81) and from (ii) of (1.1), the function $\xi \mapsto h_{P_0}(\xi) + \gamma\xi$ is also a minimizer for any $\gamma > 0$. Hence, without loss of generality, we can regard $h_{P_0}(\xi) > 0$ for all $\xi \in \mathbb{S}^{n-1}$. Recalling Theorem 2.38, for any arbitrary $v \in C_+(\mathbb{S}^{n-1})$, let Ω_v is the Wulff shape of v as

in (2.37) and then $h_{\Omega_v} = v$ holds S_{Ω_v} -a.e. Let $h_0 = h_{P_0}$ and

$$\Omega^t = P_0 + t\Omega_v \text{ so that } h_{\Omega^t} = h_{P_0} + th_{\Omega_v} = h_0 + tv, \quad S_{\Omega_v} - a.e.$$

With respect to $u = u_P$ denoted above that is p -harmonic in a neighborhood of $\partial P = \partial P_0$, we choose μ_{Ω^t} to denote $\Gamma(\Omega^t)$ as in (3.72), defined by $d\mu_{\Omega^t} = |\nabla u(\cdot, t)|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial\Omega^t}$ where $u(\cdot, t)$ is the unique solution of the Dirichlet problem (3.10). Without loss of generality, we can replace P_j and Ω_{h_j} above with infinitesimal dilations by $(1+t)$ with $|t| < 1/j$ which does not affect the Hausdorff convergence. Then, μ_{Ω^t} with respect to $u(\cdot, t)$ for $|t| < 1/j$ forms a special class of approximants among $\mu_{\Omega_{h_j}}$ with respect to u_j of the above (see Remark 3.62). Therefore, the minimization at h_0 leads to the existence of Lagrange multiplier λ such that we have

$$\frac{d}{dt} \Big|_{t=0} \int_{S^{n-1}} (h_0 + tv) d\mu = \lambda \frac{d}{dt} \Big|_{t=0} \Gamma(\Omega^t). \tag{3.84}$$

Now (3.77) of Proposition 3.76 together with (3.84), lead to

$$\int_{S^{n-1}} v d\mu = \lambda(n - p + 1) \int_{S^{n-1}} v d\mu_{P_0}$$

for any $v \in C_+(\mathbb{S}^{n-1})$. Then, the above is also obtained for any $v \in C(\mathbb{S}^{n-1})$ using $v = v^+ - v^-$, which is enough to conclude $\mu = \lambda(n - p + 1)\mu_{P_0}$. Since h_0 is the minimizer of (3.81), we have

$$b = \int_{S^{n-1}} h_0 d\mu = \lambda(n - p + 1) \int_{S^{n-1}} h_{P_0} d\mu_{P_0} = \lambda(n - p + 1)\Gamma(P_0),$$

so that $\lambda = b/(n - p + 1)\Gamma(P_0)$ and hence, $\mu = \frac{b}{\Gamma(P_0)}\mu_{P_0}$. In other words, we have

$$\frac{\mu}{\int_{S^{n-1}} h_{P_0} d\mu} = \frac{\mu_{P_0}}{\int_{S^{n-1}} h_{P_0} d\mu_{P_0}},$$

which is equivalent to $\mu = c\mu_{P_0}$ for some $c > 0$. By rescaling P_0 , we can ensure $\mu_{P_0} = \mu$ and $\Gamma(P_0) = 1$. The proof is complete. \square

Finally, we are ready to prove the existence theorem for general Borel measures.

Proof of Theorem 1.2 Given the measure μ on \mathbb{S}^{n-1} satisfying (1.1), note that from (i) of (1.1), we have

$$\delta = \inf_{\zeta \in \mathbb{S}^{n-1}} \int_{\mathbb{S}^{n-1}} |\langle \zeta, \xi \rangle| d\mu(\xi) > 0.$$

There exists discrete measures μ_j on \mathbb{S}^{n-1} such that $\mu_j \rightarrow \mu$ as $j \rightarrow \infty$ and we can assume that there exists $c = c(\delta) > 0$ such that for all $j \in \mathbb{N}$, we have

$$\inf_{\zeta \in \mathbb{S}^{n-1}} \int_{\mathbb{S}^{n-1}} |\langle \zeta, \xi \rangle| d\mu_j(\xi) \geq c. \tag{3.85}$$

Without loss of generality we can assume that μ_j 's have no antipodal pair of point masses. Indeed, if $\mu_j(\{\xi_j\}), \mu_j(\{-\xi_j\}) > 0$ then $\mu_j = c(\delta_{\xi_j} + \delta_{-\xi_j}) + \nu_j$ for some other discrete measure ν_j and we can replace μ_j with $\mu_j^{\varepsilon_j} = c(\delta_{\xi_j'} + \delta_{-\xi_j}) + \nu_j$ where $0 < |\xi_j' - \xi_j| < \varepsilon_j$ for $\varepsilon_j > 0$ small enough, so that $\mu_j^{\varepsilon_j}(\{\xi_j\}) = 0$ while $\mu_j^{\varepsilon_j}(\{-\xi_j\}) > 0$ and $\mu_j^{\varepsilon_j} \rightarrow \mu$ as $j \rightarrow \infty$.

After finitely many replacements by such small perturbations on all antipodal points, we can regard μ_j 's have no antipodal point masses for each $j \in \mathbb{N}$.

Now we can use Theorem 3.80 above to conclude for all $j \in \mathbb{N}$ there exists a polytope P_j of non-empty interior such that $\mu_{P_j} = \mu_j$ and $\Gamma(P_j) = 1$ defined as in (3.72) with respect to $\mu_{P_j} = (\mathbf{g}_{P_j})_*\omega_{p,j}$ where $\omega_{p,j}$ is any p -harmonic measure on ∂P_j for $p \neq n + 1$. For the case $p = n + 1$, we denote $\mu_{P_j}^{\varepsilon_j}$ and Γ^{ε_j} corresponding to $p = n + 1 + \varepsilon_j$ and using Theorem 3.80, take P_j such that $\mu_{P_j}^{\varepsilon_j} = \mu_j$ and $\Gamma^{\varepsilon_j}(P_j) = 1$ where $\varepsilon_j \neq 0$ is small enough; the limits remain the same since $\|\mu_{P_j}^{\varepsilon_j} - \mu_{P_j}\|, |\Gamma^{\varepsilon_j}(\cdot) - \Gamma(\cdot)| \rightarrow 0^+$ as $\varepsilon_j \rightarrow 0$. We show that the sequence of polytopes $\{P_j\}$ is bounded. Indeed, for any $x \in \bar{P}_j$ we have

$$\int_{\mathbb{S}^{n-1}} \langle x, \xi \rangle d\mu_j(\xi) \leq \int_{\mathbb{S}^{n-1}} h_{P_j}(\xi) d\mu_j(\xi) = \int_{\mathbb{S}^{n-1}} h_{P_j} d\mu_{P_j} = \Gamma(P_j) = 1.$$

Up to a suitable translation, we have $0 \in P_j$ and $x_j, -x_j \in \partial P_j$ such that $\text{diam}(P_j) = 2|x_j|$. Therefore, we have $\langle \pm x_j, \xi \rangle \leq h_{P_j}(\xi)$ and hence $|\langle x_j, \xi \rangle| \leq h_{P_j}(\xi)$. Using this on the above together with (3.85), we get

$$c|x_j| \leq \int_{\mathbb{S}^{n-1}} |\langle x_j, \xi \rangle| d\mu_j(\xi) \leq \int_{\mathbb{S}^{n-1}} h_{P_j}(\xi) d\mu_j(\xi) = 1,$$

which implies $\text{diam}(P_j)/2 = |x_j| \leq 1/c$. From Blaschke selection theorem (Theorem 2.44), there exists a convex set K such that, up to subsequence, $d_{\mathcal{H}}(P_j, K) \rightarrow 0^+$ as $j \rightarrow \infty$. Let ω_p be any p -harmonic measure on ∂K with respect to which we have μ_K as in (3.5) and $u = u_K$ be the corresponding p -harmonic function in a neighborhood of ∂K . Then, we choose the p -harmonic measures $\omega_{p,j}$ as $d\omega_{p,j} = |\nabla u_j|^{p-1} d\mathcal{H}^{n-1} \llcorner_{\partial P_j}$ where, with respect to this $u = u_K$, we choose u_j 's uniquely as the solutions of the Dirichlet problem (3.61) as in the previous subsection, so that we have $\mu_{P_j} \rightarrow \mu_K$ from (3.71) and hence $\Gamma(P_j) \rightarrow \Gamma(K)$ as $j \rightarrow \infty$, leading to $\Gamma(K) = 1$ and from uniqueness of the weak limit, $\mu_K = \mu$. Now we show that K has a non-empty interior.

Suppose K is of empty interior so that $K = \partial K$ then, the Hausdorff dimension $\dim_{\mathcal{H}}(K) < n$. Now if $\dim_{\mathcal{H}}(K) < n - 1$ then $\mathcal{H}^{n-1}(K) = 0 = \mathcal{H}^{n-1}(\partial K)$ would imply $\mu_K \equiv 0$ from (3.5) and hence $\Gamma(K) = 0$ which contradicts $\Gamma(K) = 1$. Therefore, as $\dim_{\mathcal{H}}(K) \in \mathbb{N}$, we conclude $\dim_{\mathcal{H}}(K) = n - 1$. Since $\mu_K = \mu$ satisfy (1.1), we invoke Lemma 3.79 for a contradiction.

Therefore, there exists a domain Ω such that $K = \bar{\Omega}$ and $\mu_{\Omega} = \mu$. The proof is complete. □

Acknowledgements The authors are thankful to Peter Van Hintum and the anonymous referee for their helpful remarks on a previous version of this paper.

Data availability No data was used for the research described in the article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Akman, M., Gong, J., Hineman, J., Lewis, J., Vogel, A.: The Brunn–Minkowski inequality and a Minkowski problem for nonlinear capacity. *Mem. Am. Math. Soc.* **275**(1348), vi+115 (2022)
2. Akman, M., Lewis, J., Saari, O., Vogel, A.: The Brunn–Minkowski inequality and a Minkowski problem for \mathcal{A} -harmonic Green's function. *Adv. Calc. Var.* **14**(2), 247–302 (2021)
3. Akman, M., Lewis, J., Vogel, A.: σ -finiteness of elliptic measures for quasilinear elliptic PDE in space. *Adv. Math.* **309**, 512–557 (2017)
4. Akman, M., Lewis, J., Vogel, A.: Note on an eigenvalue problem with applications to a Minkowski type regularity problem in \mathbb{R}^n . *Calc. Var. Partial Differ. Equ.* **59**(2), 47 (2020)
5. Aleksandrov, A.D.: On the theory of mixed volumes. iii. Extension of two theorems of Minkowski on convex polyhedra to arbitrary convex bodies. *Mat. Sb. (N.S.)* **3**, 27–46 (1938)
6. Aleksandrov, A.D.: Almost everywhere existence of the second differential of a convex function and some properties of convex surfaces connected with it. *Leningrad State Univ. Annals [Uchenye Zapiski] Math. Ser.*, pp. 3–35 (1939)
7. Aleksandrov, A.D.: On the surface area measure of convex bodies. *Mat. Sb. (N.S.)* **6**, 167–174 (1939)
8. Böröczky, K.J., Lutwak, E., Yang, D., Zhang, G.: The logarithmic Minkowski problem. *J. Am. Math. Soc.* **26**(3), 831–852 (2013)
9. Busemann, H., Feller, W.: Krümmungseigenschaften Konvexer Flächen. *Acta Math.* **66**(1), 1–47 (1936)
10. Caffarelli, L.A.: A localization property of viscosity solutions to the Monge–Ampère equation and their strict convexity. *Ann. Math. (2)* **131**(1), 129–134 (1990)
11. Caffarelli, L.A.: Interior $W^{2,p}$ estimates for solutions of the Monge–Ampère equation. *Ann. Math. (2)* **131**(1), 135–150 (1990)
12. Caffarelli, L.A.: Some regularity properties of solutions of Monge Ampère equation. *Commun. Pure Appl. Math.* **44**(8–9), 965–969 (1991)
13. Cheng, S.Y., Yau, S.T.: On the regularity of the solution of the n -dimensional Minkowski problem. *Commun. Pure Appl. Math.* **29**(5), 495–516 (1976)
14. Chou, K.-S., Wang, X.-J.: The L_p -Minkowski problem and the Minkowski problem in centroaffine geometry. *Adv. Math.* **205**(1), 33–83 (2006)
15. Colesanti, A., Nyström, K., Salani, P., Xiao, J., Yang, D., Zhang, G.: The Hadamard variational formula and the Minkowski problem for p -capacity. *Adv. Math.* **285**, 1511–1588 (2015)
16. Dahlberg, B.E.J.: Estimates of harmonic measure. *Arch. Ration. Mech. Anal.* **65**(3), 275–288 (1977)
17. DiBenedetto, E.: $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. *Nonlinear Anal.* **7**(8), 827–850 (1983)
18. Evans, L.C., Gariepy, R.F.: *Measure Theory and Fine Properties of Functions*. Studies in Advanced Mathematics. CRC Press, Boca Raton (1992)
19. Fenchel, W., Jessen, B.: Mengenfunktionen und konvexe körper, danske vid. selsk. Mat.-Fys. Medd **16**, 1–31 (1938)
20. Gardner, R.J.: The Brunn–Minkowski inequality. *Bull. Am. Math. Soc. (N.S.)* **39**(3), 355–405 (2002)
21. Gilbarg, D., Trudinger, N.S.: *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition
22. Guan, B., Guan, P.: Convex hypersurfaces of prescribed curvatures. *Ann. Math. (2)* **156**(2), 655–673 (2002)
23. Guan, P., Li, J., Li, Y.: Hypersurfaces of prescribed curvature measure. *Duke Math. J.* **161**(10), 1927–1942 (2012)
24. Guan, P., Ma, X.-N.: The Christoffel–Minkowski problem. I. Convexity of solutions of a Hessian equation. *Invent. Math.* **151**(3), 553–577 (2003)
25. Heinonen, J., Kilpeläinen, T., Martio, O.: *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Dover Publications Inc., Mineola (2006)
26. Jerison, D.: Prescribing harmonic measure on convex domains. *Invent. Math.* **105**(2), 375–400 (1991)
27. Jerison, D.: A Minkowski problem for electrostatic capacity. *Acta Math.* **176**(1), 1–47 (1996)
28. Kilpeläinen, T., Zhong, X.: Growth of entire \mathcal{A} -subharmonic functions. *Ann. Acad. Sci. Fenn. Math.* **28**(1), 181–192 (2003)
29. Lewis, J., Lundström, N., Nyström, K.: Boundary Harnack inequalities for operators of p -Laplace type in Reifenberg flat domains. In: *Perspectives in partial differential equations, harmonic analysis and applications*, volume 79 of Proc. Sympos. Pure Math., pages 229–266. Amer. Math. Soc., Providence, RI (2008)
30. Lewis, J., Nyström, K.: Boundary behaviour for p harmonic functions in Lipschitz and starlike Lipschitz ring domains. *Ann. Sci. École Norm. Sup. (4)* **40**(5), 765–813 (2007)

31. Lewis, J., Nyström, K.: Regularity and free boundary regularity for the p Laplacian in Lipschitz and C^1 domains. *Ann. Acad. Sci. Fenn. Math.* **33**(2), 523–548 (2008)
32. Lewis, J., Nyström, K.: Regularity and free boundary regularity for the p -Laplace operator in Reifenberg flat and Ahlfors regular domains. *J. Am. Math. Soc.* **25**(3), 827–862 (2012)
33. Lewis, J., Nyström, K., Vogel, A.: On the dimension of p -harmonic measure in space. *J. Eur. Math. Soc. (JEMS)* **15**(6), 2197–2256 (2013)
34. Lewis, J.L.: Capacitary functions in convex rings. *Arch. Ration. Mech. Anal.* **66**(3), 201–224 (1977)
35. Lewis, J.L.: Regularity of the derivatives of solutions to certain degenerate elliptic equations. *Indiana Univ. Math. J.* **32**(6), 849–858 (1983)
36. Lewis, J.L.: Note on p -harmonic measure. *Comput. Methods Funct. Theory* **6**(1), 109–144 (2006)
37. Lewis, J.L., Nyström, K.: Quasi-linear PDEs and low-dimensional sets. *J. Eur. Math. Soc. (JEMS)* **20**(7), 1689–1746 (2018)
38. Lewy, H.: On differential geometry in the large. I. Minkowski's problem. *Trans. Am. Math. Soc.* **43**(2), 258–270 (1938)
39. Lieberman, G.M.: Boundary regularity for solutions of degenerate elliptic equations. *Nonlinear Anal.* **12**(11), 1203–1219 (1988)
40. Lutwak, E., Oliker, V.: On the regularity of solutions to a generalization of the Minkowski problem. *J. Differ. Geom.* **41**(1), 227–246 (1995)
41. Minkowski, H.: Allgemeine lehrsätze über die convexen polyeder. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* **198–220**, 1897 (1897)
42. Minkowski, H.: Volumen und Oberfläche. *Math. Ann.* **57**(4), 447–495 (1903)
43. Nirenberg, L.: The Weyl and Minkowski problems in differential geometry in the large. *Commun. Pure Appl. Math.* **6**, 337–394 (1953)
44. Pogorelov, A.V.: The Minkowski multidimensional problem. *Scripta Series in Mathematics*. V. H. Winston & Sons, Washington, DC; Halsted Press [John Wiley & Sons], New York-Toronto-London, 1978. Translated from the Russian by Vladimir Oliker, Introduction by Louis Nirenberg
45. Schneider, R.: *Convex Bodies: The Brunn–Minkowski Theory*. *Encyclopedia of Mathematics and its Applications*, vol. 44. Cambridge University Press, Cambridge (1993)
46. Tolksdorf, P.: Regularity for a more general class of quasilinear elliptic equations. *J. Differ. Equ.* **51**(1), 126–150 (1984)
47. Weil, W.: Über die Projektionkörper konvexer Polytope. *Arch. Math. (Basel)* **22**, 664–672 (1971)
48. Weil, W.: Ein Approximationssatz für konvexe Körper. *Manuscr. Math.* **8**, 335–362 (1973)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.