



OPEN ACCESS

Inequality in green space distribution and its association with preventable deaths across urban neighbourhoods in the UK, stratified by Index of Multiple Deprivation

Tran Thu Ngan ,¹ Ruoyu Wang,^{1,2} Christopher Tate,¹ Mark Green ,³ Richard Mitchell,⁴ Ruth F Hunter ,¹ Ciaran O'Neill¹

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/jech-2024-222485>).

¹Centre for Public Health, Queen's University Belfast, Belfast, UK

²Institute of Public Health and Wellbeing, University of Essex, Essex, UK

³Department of Geography & Planning, University of Liverpool, Liverpool, UK

⁴School of Health and Wellbeing, University of Glasgow, Glasgow, UK

Correspondence to

Dr Tran Thu Ngan;
n.t.tran@qub.ac.uk

Received 13 May 2024

Accepted 18 September 2024

Published Online First

12 November 2024

ABSTRACT

Background This study investigated inequalities in the distribution of green space (GS) and the association between inequalities in amounts of GS and preventable deaths across urban neighbourhoods with different Index of Multiple Deprivation (IMD) scores in the UK.

Methods Data on preventable deaths, IMD, percentage of grassland and woodland, urban/rural, population size, and density were sourced for each of 6791 middle-layer super output areas (MSOAs) in England, 410 MSOAs in Wales, 1279 intermediate zones (IZs) in Scotland, and 890 super output areas (SOAs) in Northern Ireland (NI). While appreciating the potential for ecological fallacy we related area-based measures of deprivation to deaths. Concentration curves, Lorenz dominance tests, and negative binomial regression models were used to analyse the data.

Results In urban areas of England, Scotland, and NI, the percentage of grassland was significantly lower among the more deprived neighbourhoods (Lorenz test, $p < 0.0001$). In England, a 1% increase in grassland area was associated with a 37% reduction in annual preventable deaths among the most deprived urban MSOAs (incidence rate ratio (IRR) 0.63, 95% CI 0.52 to 0.76). In NI and Scotland, a 1% increase in grassland area was associated with a 37% (IRR 0.63, 95% CI 0.43 to 0.91) and 41% (IRR 0.59, 95% CI 0.42 to 0.81) reduction in 5-year accumulated preventable deaths in the most deprived urban SOAs/IZs, respectively.

Conclusions Results suggest that investment in GS in urban areas may be an important public health prevention strategy. There is evidence that investments in the most deprived urban neighbourhoods where the highest inequality currently exists would see the largest effect on preventable deaths.

BACKGROUND

Investment in green space (GS) (green in the sense of being predominantly covered with vegetation¹) in urban areas may help reduce preventable deaths, and generate various benefits that have been well-documented.^{2–5} Exposure to GS is associated with reduction of mortality and morbidity from chronic diseases, improvement of mental health and cognitive functioning, and reduction of obesity.⁶ These benefits might be realised through the provision of opportunities for physical activity/exercise and

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Investment in green space (GS) may help reduce preventable deaths.
- ⇒ GS in urban areas tends to be more accessible within less deprived neighbourhoods than in neighbourhoods of lower socioeconomic position.
- ⇒ There is a paucity of UK-wide data investigating inequalities in the distribution of GS by deprivation levels as well as the association between GS and preventable deaths among neighbourhoods of different deprivation levels.

WHAT THIS STUDY ADDS

- ⇒ Except for Wales, there was significant inequality in the amount of GS across urban neighbourhoods with different Index of Multiple Deprivation (IMD) scores across the UK, to the advantage of those in more affluent areas.
- ⇒ In urban areas of England, NI, and Scotland, a higher percentage of grassland was significantly associated with a lower number of preventable deaths among the most deprived group of IMD quintiles.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Increased investment in GS in urban areas is urgently needed and should be prioritised in the most deprived neighbourhoods where provision is currently lowest and preventable deaths are highest.
- ⇒ Investing in GS in the most deprived urban neighbourhoods is important for public health prevention strategies, particularly in light of the ongoing cost-of-living crisis and growing NHS issues in the UK.

settings for relaxation and restoration from stress and fatigue; facilitating social interaction; and promoting social cohesion.^{6,7} GS also provides environmental benefits (eg, mitigate urban heat island effect, reduce air and noise pollution, function as biodiversity and nature conservation) and economic benefits (eg, reduced energy costs of cooling buildings and increase property values) which may benefit health outcomes.^{8,9}



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

To cite: Ngan TT, Wang R, Tate C, et al. *J Epidemiol Community Health* 2025;**79**:102–109.

While investment in GS in urban areas may help reduce preventable deaths, where to prioritise investment is an important question for both equity and efficiency reasons. In European cities, accessibility to GS (within a 300 m linear distance of residences) tends to be higher within less deprived neighbourhoods than in neighbourhoods of lower socioeconomic position.¹⁰ This trend has been observed in Stockholm (Sweden),¹¹ Porto (Portugal),¹² Debrecen (Hungary),¹³ and in cities across the Netherlands.¹⁴ Similarly, a study conducted in England reported that local authorities with poorer GS provision had larger Black, Asian and minority ethnic populations and higher proportions of low-income households¹⁵—two population groups that already carry an excessively high burden of co-morbidities attributable to ill-health. UK data comparing the percentage of GS across urban neighbourhoods of different deprivation levels is absent. Further, research on the association between GS and preventable deaths among urban neighbourhoods of different deprivation levels is needed to identify geographic and socioeconomic priorities for interventions to improve health.

The aim of this study was to investigate: (1) the inequality in GS distribution in both urban and rural areas (for comparison); and (2) the association between availability of GS and preventable deaths across urban neighbourhoods in the UK, stratified by the Index of Multiple Deprivation (IMD).

METHODS

Variables, measurements, and data sources

Statistical output geographies (from small to large) are: output areas (OAs), lower-layer super output areas (LSOAs), and middle-layer super output areas (MSOAs) for England and Wales; data zones (DZs) and intermediate zones (IZs) for Scotland; and small areas (SAs) and super output areas (SOAs) for Northern Ireland (NI) (see online supplemental table 1 in supplementary materials for details). Due to data availability, the primary geographic unit of analyses were the smallest geographic areas on which data on GS and preventable deaths were available. As such, it is MSOA for England and Wales, IZ for Scotland, and SOA for NI. There were 6791 MSOAs in England and 410 MSOAs in Wales; average MSOA population size is 7600 and 7200, respectively. There were 1279 IZs in Scotland with average population

size ~4000, and 890 SOAs in NI with average population size ~2000. MSOA/IZ/SOA boundaries are mutually exclusive. Table 1 shows the sources of data.

Outcome/dependent variable: preventable deaths

Preventable death refers to ‘causes of death that can be mainly avoided through effective public health and primary prevention interventions’ (that is, before the onset of diseases or injuries, to reduce incidence)¹⁶—for example, tuberculosis, ischaemic heart diseases, cerebrovascular diseases, cancer (depends on types), and chronic obstructive pulmonary disease.

Data on preventable deaths were provided by equivalent statistics offices in Scotland and NI (table 1) or calculated from data on deaths by all causes using the Organisation for Economic Co-operation and Development (OECD) lists of preventable and treatable causes of mortality (2022 version)¹⁶ in the cases of England and Wales.

Independent variables

Index of Multiple Deprivation (IMD) is an overall relative measure of deprivation calculated by combining the weighted indices of deprivation from several domains (eg, income, employment, education, health, crime, access to services, and living environment). Each of these domains is based on a set of indicators. The number of domains and indicators are slightly different across each constituent country in the UK (see online supplemental table 2 for details).

IMD was originally measured and reported at LSOAs (England, Wales), DZs (Scotland), and SOAs (NI). As preventable deaths data are available at MSOAs for England/Wales, IZs for Scotland and SOAs for NI, the IMD was used as it is in the analysis for NI, while for the remaining countries IMDs were aggregated from the lower statistical output geographies. First, average rank and score for each MSOA/IZ were calculated by averaging all of the LSOA/DZ ranks and scores within each MSOA/IZ after these LSOA/DZ ranks and scores have been population weighted. These population weighted average ranks and average scores were then ranked, where the rank of one (most deprived) was given to the area with the ‘lowest average

Table 1 Variables, measurements, and data sources

Variables	Type	Data sources			
		England	Wales	Scotland	Northern Ireland
Preventable deaths	Number (integer)	<ul style="list-style-type: none"> ▶ 2019 data of deaths by all causes (ICD-10) were sourced from ONS¹⁸ ▶ Preventable deaths were calculated using OECD 2022 definition¹⁶ 		2015–2019* data were provided by the Vital Events Team at National Records of Scotland using OECD 2022 definition ¹⁶	2016–2020* data were provided by Public Health Information and Research Branch (PHIRB), Department of Health using OECD 2022 definition ¹⁶
Index of Multiple Deprivation	<ul style="list-style-type: none"> ▶ Ranking (rank first–most deprived) ▶ Quintiles 	English Index of Multiple Deprivation (IMD) 2019 ³⁵	Welsh Index of Multiple Deprivation (WIMD) 2019 ³⁶	Scottish Index of Multiple Deprivation 2020 (SIMD) v2 ³⁷	Northern Ireland Multiple Deprivation Measure (MDM) 2017 ³⁸
Percentage of grassland or woodland	Proportion to total land area	UK Land Cover Map 2019 (20 m classified pixels imagery) ¹⁷			
Percentage of green space areas	Proportion to total land area	Calculated by summing the areas of grassland and woodland (UK Land Cover Map 2019) ¹⁷			
Population size	Number of people	Mid-2015 population estimates			
Population density	Number of people per km ²	Mid-2017 population estimates (Scotland)			
Settlement category	Categories (urban/rural)	Census 2011 ¹⁸			

*Due to the small number of deaths in these countries, the 5-year accumulated data were used. While time periods differ slightly, this is unlikely to affect results materially given the minor differences involved.

ICD-10, International Classification of Diseases, 10th revision; OECD, Organisation for Economic Co-operation and Development; ONS, Office for National Statistics.

rank' or 'highest average score'. Due to data availability, rank of average rank and rank of average score were used in the analysis for England/Scotland and Wales, respectively.

Percentage of grassland and percentage of woodland were calculated by dividing the area of land covered by grass/wood by the total land area of a given geographic area. Data from the UK Centre for Ecology & Hydrology (UKCEH) Land Cover Map 2019 (20 m classified pixels imagery)¹⁷ were used for this calculation. In this dataset, there were 19 mutually exclusive classes of 'land cover' (see online supplemental table 3 for details). Grassland was made up of class 4–7 and 10 (improve grassland, neutral grassland, calcareous grassland, acid grassland, and heather grassland) while woodland comprised class 1 and 2 (deciduous woodland and coniferous woodland). Percentage of GS was calculated as the sum of percentage of grassland and woodland.

Settlement category (urban/rural) is a measure taken from Census 2011 data on the Office for National Statistics (ONS) Nomis website¹⁸ (see the ONS leaflet in online supplemental materials for details on how rural-urban classification was assigned to each MSOAs).

Population and population density measures are from mid-2015 (England, Wales, and NI) and mid-2017 (Scotland) population estimates which were also the data point used in the calculation of IMD.

Median age was also calculated for each MSOA/IZ/SOA using the above population estimates by age.

Data analysis

Descriptive statistics (median and IQR) were used to describe the characteristics of the samples. Concentration curves and concentration index (C-index) were used to examine the inequality in the amount of GS among MSOAs/IZs/SOAs across the quintile of IMD. The concentration curve plots the cumulative percentage of GS areas (y-axis) against the cumulative percentage of the MSOA/IZ/SOA, ranked by IMD, beginning with the most deprived, and ending with the least deprived (x-axis). The line of equality is the 45° line, showing that irrespective of the geographic areas' IMD, they have the same percentage of GS areas to total land areas. Lorenz (concentration) dominance tests were used to test if a concentration curve departs significantly from an equal distribution.¹⁹ The concentration index (bounded between -1 and +1), defined as twice the area between the concentration curve and the line of equality, would be zero if there is no socioeconomic-related inequality.¹⁹ The negative/positive values indicate a distribution of GS that favours more/less deprived areas, respectively.

The factors that influenced preventable deaths were investigated using a negative binomial regression model, given the count nature of preventable deaths, and its distribution significantly differed from a Poisson distribution. In each country, the model was run only among the urban MSOAs/IZs/SOAs and stratified by each quintile of IMD to assess how associations varied by level of deprivation. The models have 'count of preventable deaths' as the outcome; 'percentage of grassland/woodland', 'IMD', 'median age', and 'population density' as independent variables; and 'population size' as an offset variable. The various statistical procedures were conducted in STATA version 15.0.

RESULTS

Characteristics of the data are presented in table 2. The median percentage of GS was highest in Wales (45%), followed by NI and England (24% and 21%, respectively), and then Scotland

Table 2 Characteristics of the data

	Rural	Urban	Total
England, number of MSOAs (%)	1192 (17.6)	5599 (82.4)	6791 (100.0)
	<i>Median (IQR)</i>	<i>Median (IQR)</i>	<i>Median (IQR)</i>
Number of preventable deaths	19 (12)	18 (12)*	18 (12)
Percentage of grassland†	33 (32)	12 (22)*	15 (26)
Percentage of woodland†	7 (9)	2 (8)*	3 (9)
Percentage of green space area†	43 (36)	17 (28)*	21 (33)
Population (mid-2015 population estimates)	7355 (2198)	7904 (2319)*	7834 (2343)
Population density (mid-2015) (per km ²)	120 (164)	3271 (3494)*	2623 (3926)
Median age of the population (mid-2015)	46 (5)	38 (10)*	40 (10)
Percentage of population in age group 60–74‡	20 (4)	14 (6)*	15 (7)
Wales, number of MSOAs (%)	134 (32.7)	276 (67.3)	410 (100.0)
	<i>Median (IQR)</i>	<i>Median (IQR)</i>	<i>Median (IQR)</i>
Number of preventable deaths	21 (12)	21 (12)	21 (12)
Percentage of grassland†	66 (24)	27 (38)*	41 (46)
Percentage of woodland†	2 (8)	0* (1)*	0.02 (3)
Percentage of green space area	71 (23)	28 (43)*	45 (53)
Population (mid-2015 population estimates)	7120 (1960)	7413 (2405)*	7328 (2159)
Population density (mid-2015) (per km ²)	98 (188)	1294 (2570)*	735 (2035)
Median age of the population (mid-2015)	46 (6.25)	40 (7)*	41.50 (7)
Percentage of population in age group 60–74‡	21 (5)	16 (4)*	17 (5)
Scotland, number of IZs (%)	288 (22.5)	991 (77.5)	1279 (100.0)
	<i>Median (IQR)</i>	<i>Median (IQR)</i>	<i>Median (IQR)</i>
Number of preventable deaths 2015–2019 (accumulated)	32 (17)	39 (28)*	38 (25)
Percentage of grassland†	28 (38)	4 (13)*	6 (22)
Percentage of woodland†	11 (19)	5 (12)*	7 (14)
Percentage of green space area†	45 (50)	11 (22)*	16 (34)
Population (mid-2017 population estimates)	3830 (1633)	4170 (1551)*	4101 (1645)
Population density (mid-2017) (per km ²)	100 (0)	3200 (2200)*	27 (36)
Median age of the population (mid-2017)	46 (6)	41 (8)*	42 (9)
Percentage of population in age group 60–74‡	20 (5)	16 (5)*	17 (6)
Northern Ireland, number of SOAs (%)	340 (38.2)	550 (61.8)	890 (100.0)
	<i>Median (IQR)</i>	<i>Median (IQR)</i>	<i>Median (IQR)</i>
Number of preventable deaths 2016–2020 (accumulated)	13 (7)	15 (12)*	14 (10)
Percentage of grassland†	70 (38)	4 (20)*	17 (62)
Percentage of woodland†	7 (6)	0.3 (4)*	3 (7)
Percentage of green space area†	80 (30)	7 (25)*	24 (70)
Population (mid-2015 population estimates)	2291 (944)	1808 (707)*	1952 (870)
Population density (mid-2015) (per km ²)	100 (100)	3000 (2900)*	13 (34)
Median age of the population (mid-2015)	29 (13)	30 (14)*	29 (13)

*Significant difference with $p < 0.001$ (Mann-Whitney test between urban and rural).

†To total land areas.

‡Data about percentage of population in age group 60–74 are not available for Northern Ireland.

IZs, intermediate zones; MSOAs, middle-layer super output areas; SOAs, super output areas.

(16%). In all four countries, the median percentage of grassland/woodland/GS in urban areas was significantly lower than that of rural areas (Mann Whitney test, $p < 0.001$).

Concentration curves for urban and rural MSOAs/IZs/SOAs of the four countries are plotted in figure 1a–d. In England, Scotland, and NI, the concentration curves of GS for urban areas depart significantly from an equal distribution (Lorenz test, $p < 0.0001$). These curves lie below the line of equality, indicating that the amount of GS is lower in more deprived areas.




Figure 1 Concentration curves for percentage of green space areas to total land areas in England, Wales, Scotland, and Northern Ireland, by urban and rural areas.

In contrast, the concentration curves of GS for rural areas do not depart significantly from the line of equality (Lorenz test, $p > 0.05$), indicating that there is an equal amount of GS in rural areas of differing deprivation levels.

In Wales, both the concentration curves for urban and rural areas do not depart significantly from the line of equality (Lorenz test, $p > 0.05$). As such, GS is distributed more equally across neighbourhoods with different levels of deprivation in both urban and rural areas of Wales.

The C-index and detailed results of the Lorenz dominance tests are presented in table 3. The magnitude of the index reflects the strength of the relationship between GS distribution and IMD as well as the degree of variability in the distribution. As such, the variability in distribution of GS among neighbourhoods with different IMD is largest in England (C-index=0.17), followed by NI (C-index=0.13) and Scotland (C-index=0.09).

Table 4 presents results of the negative binomial regression models investigating factors that influenced preventable deaths in urban areas, stratified by quintiles of IMD.

In urban areas of England, Scotland, and NI, higher percentage of grassland was significantly associated with a lower number of preventable deaths among the most deprived group of IMD quintiles. In England, with every 1% increase of grassland area in an MSOA, the incidence rate ratio (IRR) suggests that annual preventable deaths count in that MSOAs were lower by 37%

(IRR 0.63, 95% CI 0.52 to 0.76). In NI and Scotland, with every 1% increase of grassland area in an SOA/IZ, the IRRs suggested that 5-year accumulated preventable deaths count in that SOA/IZ were lower by approximately 37% (IRR 0.63, 95% CI 0.43 to 0.91) and 41% (IRR 0.59, 95% CI 0.42 to 0.81), respectively.

In Wales, the percentage of grassland was not significantly associated with the number of preventable deaths. There were also no significant associations found between woodland area and the number of preventable deaths in Wales, Scotland, and NI. In England, the associations were observed among the third and least deprived group of IMD quintiles.

DISCUSSION

This study reported two key findings. First, except for Wales, there was significant inequality in the amount of GS among urban neighbourhoods with different IMD across the UK, to the advantage of those in more affluent areas. Second, in urban areas of England, NI, and Scotland, a higher percentage of grassland was significantly associated with a lower number of preventable deaths among the most deprived group of IMD quintiles.

Inequality in the amount of GS in urban areas

The amount of GS was lowest among the most deprived areas of England, Scotland, and NI. This result echoes a previous

Table 3 Concentration index of grassland, woodland, and green space in UK, by urban and rural

	Grassland*		Woodland*		Green space*	
	C-index (SE)	P value†	C-index (SE)	P value†	C-index (SE)	P value†
England						
Urban (n=5599)	0.17 (0.01)	<0.0001‡	0.16 (0.01)	<0.0001‡	0.17 (0.01)	<0.0001‡
Rural (n=1192)	0.002 (0.01)	0.81	0.06 (0.02)	0.0002‡	0.01 (0.01)	0.1093
Wales						
Urban (n=276)	-0.05 (0.03)	0.0703	0.26 (0.10)	0.0109‡	-0.02 (0.03)	0.3785
Rural (n=134)	-0.02 (0.01)	0.1711	0.33 (0.08)	0.0001‡	0.01 (0.01)	0.4617
Scotland						
Urban (n=991)	0.13 (0.03)	<0.0001‡	0.10 (0.02)	0.0151‡	0.09 (0.02)	<0.0001‡
Rural (n=288)	0.02 (0.03)	0.4681	0.10 (0.03)	0.1084	0.03 (0.02)	0.2072
Northern Ireland						
Urban (n=550)	0.12 (0.03)	0.0003‡	0.16 (0.05)	0.0008‡	0.13 (0.03)	<0.0001‡
Rural (n=340)	0.01 (0.01)	0.6389	-0.08 (0.02)	0.001‡	-0.003 (0.01)	0.7932

*Percentage of area to total land area (green space area=grassland area+woodland area).
†Results of the Lorenz (concentration) dominance tests (to test if a concentration curve departs significantly from an equal distribution).
‡Statistically significant at p<0.05.
C-index, concentration index value; SE, Standard error.

2008 study in England which reported that those with greater exposure to GS were more likely to be less deprived ($r^2 = -0.28$, $p < 0.0001$).²⁰ This trend is also similar in Australian and European cities. For example, GS accounted for 20% of land area in the most affluent areas compared with 12% among the most low-income neighbourhoods of Adelaide (Australia).²¹ In Porto (Portugal), the number of accessible GS was smallest in the most deprived neighbourhoods and the distance to GS increased with neighbourhood deprivation.¹²

In the Netherlands, lower presence and quality of GS were observed in neighbourhoods with a low socioeconomic status compared with those with a high socioeconomic status.¹⁴ In Germany, neighbourhoods with high multiple deprivation index scores, including low average income, low levels of educational attainment and high unemployment rates, tended to have access to smaller areas of GS than their more affluent counterparts.²²⁻²³ With the known health benefits of GS, this discrepancy may help to explain the wide health inequalities

Table 4 Negative binomial regression model investigating factors that influenced preventable deaths in urban areas, stratified by quintiles of multiple deprivation

	Most deprived IRR (95% CI)	Second IRR (95% CI)	Third IRR (95% CI)	Fourth IRR (95% CI)	Least deprived IRR (95% CI)
England* (MSOA level)	n=1335	n=1216	n=1082	n=915	n=1053
Percentage of grassland	0.63 (0.52 to 0.76)†	0.81 (0.68 to 0.97)†	0.84 (0.70 to 1.02)	0.76 (0.63 to 0.93)†	0.84 (0.68 to 1.03)
Percentage of woodland	0.85 (0.60 to 1.20)	0.91 (0.66 to 1.26)	0.63 (0.43 to 0.91)†	0.71 (0.46 to 1.08)	0.56 (0.41 to 0.77)†
Median age of the population	1.05 (1.04 to 1.05)†	1.05 (1.04 to 1.05)†	1.05 (1.04 to 1.05)†	1.04 (1.04 to 1.05)†	1.05 (1.04 to 1.05)†
Population density (per 1000 m ²)	0.95 (0.94 to 0.96)†	0.95 (0.95 to 0.96)†	0.95 (0.94 to 0.96)†	0.94 (0.93 to 0.96)†	0.95 (0.93 to 0.97)†
Wales* (MSOA level)	n=50	n=38	n=55	n=63	n=70
Percentage of grassland	1.15 (0.39 to 3.43)	1.98 (0.94 to 4.16)	0.96 (0.60 to 1.53)	0.91 (0.53 to 1.58)	0.74 (0.51 to 1.07)
Percentage of woodland	0.11 (0.00 to 1.69)	0.29 (0.01 to 16.6)	0.58 (0.16 to 2.18)	2.06 (0.74 to 5.72)	0.79 (0.38 to 1.63)
Median age of the population	1.06 (1.02 to 1.10)†	1.05 (1.03 to 1.07)†	1.04 (1.02 to 1.06)†	1.04 (1.02 to 1.07)†	1.03 (1.01 to 1.04)†
Population density (per 1000 m ²)	1.00 (0.87 to 1.15)	1.08 (0.98 to 1.18)	0.97 (0.91 to 1.03)	1.02 (0.94 to 1.09)	0.97 (0.93 to 1.01)
Scotland‡ (IZ level)	n=245	n=214	n=165	n=144	n=223
Percentage of grassland	0.59 (0.42 to 0.81)†	0.77 (0.63 to 0.94)†	0.89 (0.71 to 1.10)	0.93 (0.72 to 1.20)	0.94 (0.72 to 1.21)
Percentage of woodland	1.20 (0.85 to 1.71)	0.84 (0.62 to 1.15)	0.90 (0.59 to 1.37)	0.73 (0.44 to 1.22)	0.99 (0.62 to 1.56)
Median age of the population	1.01 (1.00 to 1.02)†	1.01 (1.00 to 1.02)†	1.01 (1.00 to 1.01)†	1.02 (1.01 to 1.02)†	1.02 (1.01 to 1.03)†
Population density (per 1000 m ²)	0.98 (0.97 to 0.998)†	0.98 (0.96 to 0.996)†	0.99 (0.98 to 1.01)	0.99 (0.97 to 1.01)	0.99 (0.97 to 1.01)
Northern Ireland‡ (SOA level)	n=151	n=106	n=82	n=81	n=130
Percentage of grassland	0.63 (0.43 to 0.91)†	1.35 (0.92 to 1.96)	0.97 (0.60 to 1.55)	1.34 (0.83 to 2.17)	0.75 (0.48 to 1.18)
Percentage of woodland	0.49 (0.11 to 2.26)	0.83 (0.35 to 1.96)	0.53 (0.15 to 1.87)	0.44 (0.11 to 1.62)	0.65 (0.24 to 1.71)
Median age of the population	1.02 (1.01 to 1.03)†	1.02 (1.01 to 1.03)†	1.02 (1.01 to 1.03)†	1.02 (1.01 to 1.03)†	1.02 (1.01 to 1.03)†
Population density (per 1000 m ²)	1.01 (0.99 to 1.04)†	1.01 (0.98 to 1.04)	0.98 (0.94 to 1.02)	1.00 (0.95 to 1.04)	0.96 (0.91 to 1.01)

*Outcome variable is annual preventable deaths.

†P<0.05.

‡Outcome variable is 5-year accumulated preventable deaths.

CI, Confidence interval; IRR, incidence rate ratio; IZ, intermediate zones (average population size ~4000); MSOA, middle layer super output areas (average population size ~7000); SOA, super output areas (average population size ~2000).

in urban areas in which the poorest and most vulnerable are most impacted.

The unequal distribution of GS in urban areas demonstrates the need to target interventions at more deprived urban areas. Studies have shown that GS brings about greater benefit to those of lower socioeconomic position than those who belong to the more privileged groups, particularly in mental health and social integration.^{24–26} This suggests that GS distribution mirrors the ‘inverse care law’,²⁷ in that it is least available to the communities who need it most, and for whom it might have greatest impact. Knowledge of where the greatest need for investment in GS exists can, therefore, help increase the return on investment, and cost-effectiveness, of such interventions.⁵ In addition, community involvement in the development and management of GS should be carried out as it can increase the sense of ownership, and potentially encourage use of the facilities.^{5 28–30}

GS and preventable deaths in urban areas

The analysis found a statistically significant association between the percentage of grassland and number of preventable deaths among the most deprived urban areas in three of the four countries of the UK. It is perhaps notable that the exception, Wales, was where inequalities in the amount of GS were not significant (in either urban or rural areas). We observed that a 1% increase in grassland area was significantly associated with a 37–41% reduction in the number of preventable deaths. However, these associations were not consistent across other quintiles of multiple deprivation. This finding adds to the evidence from Ruoyu *et al* (NI)³¹ and Mitchell and Popham²⁰ (England) which reported that among the least green areas, incidence rates for all-cause mortality of the most income deprived quartile was 1.93 (95% CI 1.86 to 2.01) times higher than that of the least deprived.²⁰ These findings suggest that a greater impact can be achieved from investment in GS targeting the most deprived urban areas. It also echoes the fact that preventable deaths are massively socially patterned, and therefore influenced by socioeconomic factors rather than just a lack of healthcare intervention.

Although the study results show the association between lower number of preventable deaths and higher percentage of GS area, investment should not solely focus on increasing the existence of GS but also on the accessibility and quality of such GS.^{5 30 32} In reality, the land area covered by GS is often bigger than the area of GS that is accessible to the public. A European Environment Agency study from 38 countries reported that area of publicly available GS only accounted for 3% of the total city area (though GS of all kinds made up 42% of the total land areas).³³ Further research investigating the link between health outcomes and accessibility to GS should be explored.

Strength and limitations

To the best of our knowledge, this is the first UK-wide study that used data at the small area level to look at inequalities in the amount of GS among neighbourhoods and the associations between levels of GS in urban areas and preventable deaths across levels of deprivation. Data on different parameters were sourced from various government public databases or directly from the national statistic agencies before linking with each other in order to create a robust dataset for secondary analysis.

There were several limitations to the present study. First, we relate area-based measures of deprivation to deaths assuming implicitly that those living and dying in deprived areas are themselves more likely to be deprived. This may not be the case—being rich in a deprived area may be positively correlated with

avoidable death—however, in the absence of data on access, preventable death and deprivation at the individual level, we are obliged to proceed recognising the potential for ecological fallacies. We also aggregated IMD to larger statistical output geographies (from LSOA/DZ to MSOA/IZ) for which it was not designed; thus, it might introduce ecological fallacies into our analyses. The ecological nature of our analyses mean that we cannot rule out residual confounding (eg, greener areas may just pick up neighbourhood deprivation indirectly). Besides, the use of statistical output geographies as a unit of analysis carries an inherent risk of error with respect to individuals, especially those living on the boundary of the MSOAs/IZs/SOAs as very nearby GS in the adjacent unit may be more relevant (for actual use) than GS at the other side of their own administrative unit.

Second, the study used ‘land cover’ data which show the visible surface of land (eg, crops, grass, broad-leaved forest or built-up area). ‘Land use’ data might be a better measurement for analysis since they indicate how people are using land (eg, agriculture, forestry, recreation, or residential use) which is part of the mechanism on how GS may benefit health. Data showing frequency of access to GS, activities performed, and length of each visit could have improved on these measures, although such national level data are rarely available to researchers. As such data are not available, we had to resort to ‘land cover’ data under the assumption that land cover and use are positively correlated with each other.

Thirdly, the negative binomial models could be adjusted further for quality of GS (using, for example, the normalised difference vegetation index) which as an indicator may influence the use and activity within GS and, in turn, the number of preventable deaths. Such data are, again, not commonly available to researchers and represent an important research gap. It may be reasonable to assume that the most deprived neighbourhoods not only experience a lower quantity, but also a lower quality of GS compared with less deprived areas.³² It is plausible that the impact on preventable deaths may be even greater than the current estimates if targeted investment was directed towards increasing the area of GS as well as the quality.

Apart from the quality of GS, we appreciate a range of potential confounding variables could have been added to our analysis. These might include economic indicators, indicators of healthcare provision, crime rates and pollution, and perhaps include lagged values of these. However, many of these would be expected to be correlated with the IMD (which was calculated by combining the weighted indices of deprivation from several domains including income, employment, education, health, crime, access to services, and living environment). Each of these domains is based on a set of indicators and issues of endogeneity may arise in estimation.

Finally, this is a secondary analysis of cross-sectional data so the interpretation of the results should be treated with caution. We cannot conclude that the relationship between GS and preventable deaths is causal. It is conceivable that GS and preventable deaths are related through their relationships with other unobserved variables. Future research should consider extending our work using longitudinal data, including study designs that can identify cause and effect.³⁴ Work examining the return on investments in GS could further examine this.

CONCLUSIONS

Inequality in the amount of GS in urban areas exists across three of the four countries of the UK, with the tendency that the most deprived urban areas have the least GS provision. The strongest

association between GS area and the number of preventable deaths was observed in the most urban deprived neighbourhoods. These results support the argument for increased investment in GS, and that this should be prioritised in the most deprived neighbourhoods where provision is currently lowest and preventable deaths are highest. Investing in GS in the most deprived neighbourhoods, therefore, is important for prevention, particularly in light of the ongoing cost-of-living crisis and growing NHS issues in the UK.

X Mark Green @markalngreen and Ruth F Hunter @Ruth_HunterQUB

Acknowledgements The UK Prevention Research Partnership is funded by the British Heart Foundation, Cancer Research UK, Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Health and Social Care Research and Development Division (Welsh Government), Medical Research Council, National Institute for Health Research, Natural Environment Research Council, Public Health Agency (Northern Ireland), The Health Foundation and Wellcome. This work was conducted by GroundsWell consortium which is an interdisciplinary consortium involving researchers, policy, implementers and communities. It is led by Queen's University Belfast, University of Edinburgh and University of Liverpool in partnership with Cranfield University, University of Exeter, University of Glasgow, University of Lancaster and Liverpool John Moores University. We would like to acknowledge our partners including: Belfast, Edinburgh and Liverpool City Councils, Public Health Agencies of Scotland and Northern Ireland, Green space Scotland, Scottish Forestry, Edinburgh and Lothians Health Foundation, Department for Infrastructure Northern Ireland, Belfast Healthy Cities, Climate Northern Ireland, Health Data Research UK, Administrative Data Research Centre, NatureScot, Mersey Care NHS Foundation Trust, Liverpool City Region Combined Authority, Liverpool Health Partners, NHS Liverpool Clinical Commissioning Group, the Scottish Government, Edinburgh Health and Social Care Partnership, HSC Research and Development Office Northern Ireland, EastSide Partnership, Ashton Centre, Regenerus, Sustrans, Cycling UK, CHANGES, The Mersey Forest, Translink, Anaeko, AECOM Ltd, The Paul Hogarth Company and Moai Digital Ltd. RM is funded by the MRC and CSO (MC_UU_00022/4, SPHSU19)

Contributors TTN: conceptualisation, data curation, methodology, formal analysis, writing – original draft, writing – review and editing. RW: data curation, writing – review and editing. MG, RM, and CT: writing – review and editing. RH: funding acquisition, writing – review and editing, supervision. CON: conceptualisation, funding acquisition, writing – review and editing, supervision. TTN acted as guarantor.

Funding This work was supported by the UK Prevention Research Partnership (MR/V049704/1). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. All data generated or analysed during this study are included in this published article or downloadable from public resources (URLs in Reference list).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Tran Thu Ngan <http://orcid.org/0000-0003-2771-9878>
Mark Green <http://orcid.org/0000-0002-0942-6628>
Ruth F Hunter <http://orcid.org/0000-0001-7315-0382>

REFERENCES

- Taylor L, Hochuli DF. Defining greenspace: multiple uses across multiple disciplines. *Landsc Urban Plan* 2017;158:25–38.
- Twohig-Bennett C, Jones A. The health benefits of the great outdoors: a systematic review and meta-analysis of greenspace exposure and health outcomes. *Environ Res* 2018;166:628–37.
- World Health Organization. Regional Office for Europe. Assessing the value of urban green and blue spaces for health and well-being. Copenhagen World Health Organization. Regional Office for Europe; 2023.
- WHO. Regional Office for Europe. Urban Green Space Interventions and Health: A Review of Impacts and Effectiveness. Copenhagen: World Health Organization. Regional Office for Europe, 2017.
- Hunter RF, Cleland C, Cleary A, et al. Environmental, health, wellbeing, social and equity effects of urban green space interventions: a meta-narrative evidence synthesis. *Environ Int* 2019;130:104923.
- European Environment Agency (EEA). Healthy Environment, Healthy Lives: How the Environment Influences Health and Well-Being in Europe. Luxembourg: Publications Office of the European Union, 2020.
- Lee ACK, Jordan HC, Horsley J. Value of urban green spaces in promoting healthy living and wellbeing: prospects for planning. *Risk Manag Healthc Policy* 2015;8:131–7.
- Haq SMA. Urban Green Spaces and an Integrative Approach to Sustainable Environment. *Journal of Environmental Protection* 2011;02:601–8.
- Tate C, O'Neill C, Tran N, et al. The social return on investment of an urban regeneration project using real-world data: the Connswater Community Greenway, Belfast, UK. *Cities & Health* 2023;7:699–718.
- Barboza EP, Cirach M, Khomenko S, et al. Green space and mortality in European cities: a health impact assessment study. *Lancet Planet Health* 2021;5:e718–30.
- Goldenberg R, Kalantari Z, Destouni G. Increased access to nearby green–blue areas associated with greater metropolitan population well-being. *Land Degrad Dev* 2018;29:3607–16.
- Hoffmann E, Barros H, Ribeiro AI. Socioeconomic Inequalities in Green Space Quality and Accessibility-Evidence from a Southern European City. *Int J Environ Res Public Health* 2017;14:916.
- Csomós G, Farkas JZ, Kovács Z. Access to urban green spaces and environmental inequality in post-socialist cities. *HunGeoBull* 2020;69:191–207.
- de Vries S, Buijs AE, Snel RPH. Environmental Justice in The Netherlands: Presence and Quality of Greenspace Differ by Socioeconomic Status of Neighbourhoods. *Sustainability* 2020;12:5889.
- Public Health England. Improving Access to Greenspace: A New Review for 2020. London: PHE Publications, 2020.
- Organisation for Economic Co-Operation and Development (OECD), Eurostat. Avoidable Mortality: OECD/Eurostat Lists of Preventable and Treatable Causes of Death. Paris: OECD, 2022.
- UKCEH Land Cover Maps. UK Centre for Ecology & Hydrology (UKCEH), Available: <https://www.ceh.ac.uk/data/ukceh-land-cover-maps> [Accessed 18 May 2023].
- Official census and labour market statistics, Available: <https://www.nomisweb.co.uk>
- The World Bank. Analyzing Health Equity Using Household Survey Data: A Guide to Techniques and Their Implementation. Washington, D.C: World Bank, 2008.
- Mitchell R, Popham F. Effect of exposure to natural environment on health inequalities: an observational population study. *Lancet* 2008;372:1655–60.
- Astell-Burt T, Feng X, Mavoa S, et al. Do low-income neighbourhoods have the least green space? A cross-sectional study of Australia's most populous cities. *BMC Public Health* 2014;14:292.
- Schüle SA, Gabriel KMA, Bolte G. Relationship between neighbourhood socioeconomic position and neighbourhood public green space availability: an environmental inequality analysis in a large German city applying generalized linear models. *Int J Hyg Environ Health* 2017;220:711–8.
- Wüstemann H, Kalisch D. Towards a national indicator for urban green space provision and environmental inequalities in Germany: method and findings. Berlin Humboldt University of Berlin; 2016.
- Ward Thompson C, Aspinall P, Roe J, et al. Mitigating Stress and Supporting Health in Deprived Urban Communities: The Importance of Green Space and the Social Environment. *Int J Environ Res Public Health* 2016;13:440.
- Marselle MR, Bowler DE, Watzema J, et al. Urban street tree biodiversity and antidepressant prescriptions. *Sci Rep* 2020;10:22445.
- Camps-Calvet M, Langemeyer J, Calvet-Mir L, et al. Ecosystem services provided by urban gardens in Barcelona, Spain: insights for policy and planning. *Environmental Science & Policy* 2016;62:14–23.
- Tudor Hart J. The inverse care law. *Lancet* 1971;297:405–12.
- Wilk B, Latinos V, Hanania S, et al. Guidelines for Co-Designing and Co-Implementing Green Infrastructure in Urban Regeneration Processes, D2.10, proGReg. Horizon 2020 Grant Agreement No 776528. European Commission, 2020.
- Hansen R, Rall E, Chapman E, et al. Urban green infrastructure planning: a guide for practitioners. 2017. Available: <http://greensurge.eu/working-packages/wp5/>

- 30 Hunter RF, Christian H, Veitch J, *et al.* The impact of interventions to promote physical activity in urban green space: a systematic review and recommendations for future research. *Soc Sci Med* 2015;124:246–56.
- 31 Wang R, Grekousis G, Maguire A, *et al.* Examining the spatially varying and interactive effects of green and blue space on health outcomes in Northern Ireland using multiscale geographically weighted regression modeling. *Environ Res Commun* 2023;5:035007.
- 32 Nguyen PY, Astell-Burt T, Rahimi-Ardabili H, *et al.* Green Space Quality and Health: A Systematic Review. *Int J Environ Res Public Health* 2021;18:11028.
- 33 European Environment Agency (EEA). Urban green infrastructure. 2018. Available: https://www.eea.europa.eu/ds_resolveuid/398ff344641549dcb3bd429894b30910
- 34 Hunter RF, Nieuwenhuijsen M, Fabian C, *et al.* Advancing urban green and blue space contributions to public health. *Lancet Public Health* 2023;8:e735–42.
- 35 English indices of deprivation 2019. Min of Hous, Comm & Loc Govt, 2019. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/845345/File_7_-_All_10D2019_Scores__Ranks__Deciles_and_Population_Denominators_3.csv
- 36 Welsh Government. Welsh index of multiple deprivation 2019. 2020. Available: <https://www.gov.wales/sites/default/files/statistics-and-research/2022-02/wimd-2019-index-and-domain-scores-by-small-area.ods>
- 37 Scottish Government. Scottish index of multiple deprivation 2020. 2020. Available: <https://www.gov.scot/publications/scottish-index-of-multiple-deprivation-2020v2-ranks> [Accessed 18 May 2023].
- 38 Northern Ireland Statistics and Research Agency (NISRA). Northern Ireland multiple deprivation measure 2017. 2017. Available: https://www.nisra.gov.uk/sites/nisra.gov.uk/files/publications/NIMDM17_SOAreults.xls [Accessed 18 May 2023].