

The performance, health and development of youth women's footballers: A systematic scoping review

International Journal of Sports Science

& Coaching

1–32

© The Author(s) 2026



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/17479541251411068

journals.sagepub.com/home/spo



Thomas Adams¹, Sally Waterworth¹ , Tracy Lewis¹ , Naomi Datson² ,
Carla Harkness-Armstrong³ , Ruth Lowry⁴, Paul Freeman¹,
and Alice Harkness-Armstrong¹

Abstract

The primary aim of this scoping review was to summarise the current scientific literature on the performance, health and development of youth women's footballers. The review provides a summary of the research topics, including methodological approaches adopted and key findings, and identifies gaps in the literature. A systematic search of electronic databases was conducted in December 2023 and June 2025, with keywords relating to the population, football, and performance, health and development. Studies which involved youth women's footballers playing at any competitive standard, and quantified at least one aspect of performance, health or development were included. Of the 16,473 studies identified in the database searches, 294 studies met the eligibility criteria to be included in the review. Of the eight research topics investigating the performance, health and development of youth women's footballers, physical qualities was the most investigated ($n = 119$; 40%), followed by injury ($n = 49$; 17%), biomechanics ($n = 40$; 14%), psychology ($n = 31$; 11%), match-play ($n = 20$; 7%), nutrition ($n = 14$; 5%), fatigue and recovery ($n = 13$; 4%) and training load ($n = 8$; 3%). Players competing in regional (42%) and national competitions (32%), and within an U17 age-group (23%) were the most investigated. Over half of all studies (56%) were published from 2020 onwards, demonstrating recent rapid growth in youth women's football literature. This comprehensive resource can be used to inform practices supporting the performance, health and development of youth women's footballers across various competitive standards. Furthermore, multiple research areas are highlighted as underdeveloped, and areas for future research concerning this population to explore.

Keywords

Adolescence, biomechanics, female, injury, psychology, soccer

Introduction

Women's football has seen rapid growth within recent years. To support the sustainable development across competitive standards, women's football strategies have been implemented by global (i.e., FIFA¹), continental (e.g., OFC,² UEFA³) and national governing bodies (e.g., The FA,⁴ Irish FA⁵). Consequentially, there has been increased investment, provision and support provided across playing standards, from grassroots; increasing the number of organised women's clubs to create new playing opportunities and consequentially increasing participation rates,⁶ to elite environments; professionalising senior domestic environments,⁷ an increase in the number of teams competing at senior international major tournaments (e.g., 2023 FIFA Women's World Cup; 2028 Olympics), and the establishment of new elite competitions

(e.g., FIFA Women's Champions Cup 2026; FIFA Women's Club World Cup 2028⁸). Beyond the increased

Reviewer: Nuno Nunes (Solent University, UK)

¹School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Colchester, UK

²Department of Sport & Exercise Sciences, Manchester Metropolitan University Institute of Sport, Manchester, UK

³Centre for Physical Activity in Health and Disease, Department of Sport, Health, and Exercise Sciences, Brunel University of London, London, UK

⁴School of Psychology, Ulster University, Coleraine, UK

Corresponding author:

Alice Harkness-Armstrong, School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Wivenhoe Park, Colchester, CO4 3SQ, UK.
Email: a.harkness-armstrong@essex.ac.uk

opportunities for players, the sport has seen increased and record-breaking attendances at international and domestic fixtures, increasing media coverage and viewership, and fan engagement.⁷ The development of the sport can also be observed beyond applied environments, with a recent scoping review highlighting the recent rapid growth in the volume of scientific literature investigating women's football.⁹

The growth and development of the sport has also extended into youth environments, with women's football strategies inclusive of youth populations,¹⁻⁵ greater numbers of girls participating,⁶ and global (e.g., FIFA Talent Development Scheme^{10,11}), continental (e.g., UEFA Women's Football Development Programme¹²; CAF IMPACT Programme¹³) and national initiatives (e.g., The FA Emerging Talent Centre and Pro-Game Academies within England¹⁴; FAW National Academi within Wales¹⁵) which have increased investment, provision and support provided for youth players within talent development programmes (e.g., national talent pathways, domestic club youth academies).¹⁶⁻¹⁸ Further, FIFA has emphasised the importance of women's clubs establishing clear talent pathways for the sustained growth and development of women's football.⁷ Whilst the growth and development of youth women's football is apparent in applied environments, the same cannot be said for the body of scientific literature. Research investigating women's football has predominantly focussed upon senior populations,⁹ and where research has attempted to systematically evaluate and summarise the scientific literature within women's football, reviews have either focussed upon isolated research areas and reported limited research conducted with youth players (e.g., match-play¹⁹; menstrual cycle²⁰; coaching practice²¹), or have intentionally excluded youth players within eligibility criteria.²²⁻²⁴ Therefore, it is difficult to gain an overall perspective on the current state and extent of the scientific literature across youth women's football.

The lack of understanding of the current evidence-base in youth women's football is problematic for practitioners attempting to implement evidence-informed applied practices which support players regarding performance, health and development. Practitioners may inappropriately rely on informing practices based on a more developed evidence-base from different populations (e.g., senior women's, youth men's²⁵). However, generalisations from one population to another (e.g., senior to youth women's football) are inappropriate due to both population (e.g., biological maturation,^{26,27} physiological, biomechanical,^{28,29} training age) and contextual differences (e.g., training and competition structures, provision and support relating to sport science and medicine, governance and regulations for applied environments, rules and regulations within competitions, employment/dual-career/education). Therefore, there is a need to understand the current state and extent of scientific literature within youth women's football, to

both provide a critical and comprehensive resource for applied practitioners, but also to understand current gaps in understanding, and identify areas of future research.

Therefore, the aims of this scoping review are to: 1) systematically scope and review the scientific literature on the performance, health and development of youth women's footballers, 2) determine the methodological approaches adopted, 3) summarise the findings of research areas, and 4) identify gaps within the literature. In turn, being able to guide the focus of future research, to further benefit the performance, health and development of youth women's footballers.

Methods

Design and search strategy

This scoping review was conducted in accordance with the extension for Scoping Reviews (PRISMA-ScR; S1 Appendix).³⁰ A systematic search of electronic databases (CINAHL, EMBASE, IndMed, LILACS, MEDLINE, PubMed, Scopus, SPORTDiscus, Web of Science) was conducted on 17th December 2023, with no date or language restrictions applied. An updated search was conducted on 11th June 2025, to identify any additional studies published since the date of the original search. The search strategy included the terms for the population ('female' OR 'women's' OR 'girls' AND 'youth' OR 'adolescent') AND sport ('football' OR 'soccer' OR 'futbol') AND performance, health, or development ('perform' OR 'competit*' OR 'match' OR 'game' OR 'training' OR 'demands' OR 'activity' OR 'tactical' OR 'technical' OR 'physiolog*' OR 'physical*' OR 'testing' OR 'qualities' OR 'power' OR 'speed' OR 'fitness' OR 'change of direction' OR 'agility' OR 'skill' OR 'health' OR 'menstrual' OR 'energy' OR 'nutrition' OR 'fuel*' OR 'diet*' OR 'composition' OR 'mental' OR 'psycholog*' OR 'well*' OR 'fatigue' OR 'recover*' OR 'load' OR 'lifestyle' OR 'injur*' OR 'rehabilit*' OR 'risk' OR 'incidence' OR 'exposure' OR 'development' OR 'athletic*' OR 'hormon*' OR 'maturation*' OR 'growth' OR 'anthropomet*'). Additionally, the search strategy included AND NOT ('futsal' OR 'American football' OR 'rugby' OR 'Australian rules football' or 'Gaelic football'). The search strategy was informed by recent reviews within sports science and medicine.^{9,19,23,24,31} No review protocol was registered for this scoping review.

Study selection

After removing duplicate studies, a two-stage screening process was used. Initial screening consisted of two researchers (original search: TA, AHA; updated search: AHA, CHA) independently screening title, abstract, and keywords of studies against the eligibility criteria. Before

moving to the next stage, both researchers met to discuss and resolve any disagreements. No third reviewer was required to facilitate this process. Full-text screening was then conducted independently by the same researchers, using the same eligibility criteria, again followed by the same process where both researchers discussed and resolved any disagreements. Selected studies following this two-stage screening process were included within this review.

The eligibility criteria were guided by the elements of the Population, Concept, and Context (PCC) framework.³² Studies were included if they involved youth women's football players (PPC Population), playing at any competitive standard (PPC Context, i.e., organised competitions; local, regional, national, or international). For the purposes of this review, youth was defined as players aged 19yrs or younger (as per the National Institute of Health's definition for adolescent).³³ Studies must have quantified at least one aspect of performance, health or development (PPC Concept). Performance includes the technical, tactical, physiological and psychological characteristics of a player or team.³⁴ Health relates to the physical, mental and social well-being of the player.³⁵ Whilst development refers to the chronological and biological development of the player.³⁶ Only original research published in peer-reviewed journals were included, with conference proceedings/abstracts, book chapters, case-studies/reports, review articles, or student theses excluded. Studies which only included youth women's football players in non-competitive environments (PPC Context, i.e., recreational, physical activity, school settings), senior women's (i.e., >19yrs), youth age-groups including players older than 19yrs (i.e., U20, U23), men's, mixed-gender, or other football codes (i.e., American football, Australian rules, futsal, Gaelic football or rugby) or sports were excluded. Additionally, where studies only presented data which were not distinguishable or unique to youth women's football players (i.e., youth women's data grouped with men's,³⁷ non-competitive,³⁸ or senior women's³⁹ players' data) these studies were also excluded. Lastly, studies which used youth women's football players as participants, but did not investigate the performance, health and/or development of youth women's football players were excluded.⁴⁰

Data extraction

One researcher extracted the data (original search: TA; updated search: AHA), which was checked by another member of the research team who had expertise in the respective research topic (biomechanics: CHA; fatigue and recovery: SW, ND; injuries: TL; match-play: AHA, ND; nutrition: SW; physical qualities: ND, SW, AHA; psychology: RL, PF; training load: AHA, ND). These topics aligned to a previous systematic scoping review of the sport science and medicine literature in another sport.³¹ Studies were categorised into topics based on their primary

aim and outcome variables; this process was completed independently by two researchers (original search: TA, AHA; updated search: AHA, CHA), with any disagreements resolved via discussion with the respective researcher with expertise in the topic (e.g., biomechanics; CHA). Data relating to study characteristics (e.g., authors, year of publication, geographical location), participant characteristics (e.g., age, height, body mass, playing standard, age-group), study aim(s), methodological approach (e.g., number of participants, number of teams/clubs, data collection tools, outcome variables, comparison groups), and key findings were extracted from the full-text. Where studies included irrelevant data (e.g., playing position data which grouped senior and youth players),⁴¹ only the relevant data was extracted.

Data synthesis

Due to the nature of a scoping review in mapping, identifying and summarising the evidence within a topic, no statistical analysis was conducted.³⁰ To determine the methodological approaches adopted, findings and gaps within the literature, data relating to study and participant characteristics, and methodological approach were summarised across all studies and per topic using descriptive statistics (i.e., count, percentages, mean $\pm$ SD, or range), whilst data relating to key findings were qualitatively summarised according to topic.

Results and discussion

Search and selection of studies

A total of 16,473 studies were identified through the database searches. After removing duplicates ($n = 6,436$) and conducting the initial and full-text screening, 294 studies were deemed eligible for inclusion in this review (Figure 1).

Study characteristics

Study characteristics: research topics. Of the 294 studies included in this review, the performance, health and development of youth women's footballers were covered across eight research topics: biomechanics ($n = 40$; 14%), fatigue and recovery ($n = 13$; 4%), injury ($n = 49$; 17%), match-play ($n = 20$; 7%), nutrition ($n = 14$; 5%), physical qualities ($n = 119$; 40%), psychology ($n = 31$; 11%) and training load ($n = 8$; 3%) (Figure 2).

Study characteristics: publication year. There has been a steady increase in the number of studies investigating the performance, health and development of youth women's footballers (Figure 3), with 90% of studies ($n = 264$) published since 2010. Further, in the last five years, there has been a period of particularly rapid growth, with 56% ($n = 165$) of studies published between 2020 and 2025.




Figure 1. Flow of the selection process to determine eligible studies.




Figure 2. Number of studies within topics investigating the health, performance and development of youth women's footballers.

Similar to trends observed within senior women's football research,⁹ there appears to be an increase in the volume of studies published in the year following a major global

senior international competition (i.e., 2011, 2015, 2019, and 2023 FIFA Women's World Cups).

The earliest studies focussed on psychology (1979), injury (1985) and physical qualities (1992; Figure 4), which is reflective of the topics investigated in the early evidence-base within senior women's football research (sociology = 1939; psychology = 1975; injury = 1975).⁹ In contrast, fatigue and recovery, training load and match-play are the least established topics within youth women's football research, which may be due to the increasing accessibility of technology (e.g., global positioning systems; GPS) within recent years. Lastly, the volume of research within recent years has predominantly investigated physical qualities, with the number of publications within the research topic progressively increasing since 2017.

Study characteristics: geography of studies. Studies were predominantly from European ($n=162$; 55%) and North American ($n=81$; 28%) countries, with South American ($n=15$; 5%), Asian ($n=13$; 4%), Oceania ($n=10$; 3%) and African ($n=7$; 2%) countries contributing a smaller proportion of the evidence-base. A total of 42 different countries were represented within this review (Figure 5). The majority of studies came from: USA ($n=53$; 18%), England ($n=31$; 11%), Spain ($n=28$; 10%), Canada ($n=27$; 9%), Sweden ($n=20$; 7%), Norway ($n=16$; 6%), Germany ($n=15$; 5%), Denmark ($n=10$; 3%), Australia ($n=9$; 3%) and Brazil ($n=9$; 3%). The large proportion of research published from these countries seem unsurprising when considered in the wider context of women's football. These countries are prominent women's football nations, with successful history in senior and youth global and continental international competitions (e.g., FIFA Women's World Cup, CONCACAF Women's Championship, UEFA European Championships), and are home to the most established professional senior women's football leagues (e.g., National Women's Super League in USA, Women's Super League in England, Liga F in Spain, Damallsvenskan in Sweden, Frauen Bundesliga in Germany).⁴² However, caution may be required for practitioners using the evidence-base predominantly derived from such countries, with consideration given to the potential contextual differences (e.g., talent pathways and structures, investment, provision and support) which may exist between comparable youth women's football populations (e.g., international, national or regional competitive standards).²⁵

The representation of countries according to research topic is presented in S1 Figure. As the largest topic, the physical qualities research topic also had the widest representation of countries ($n=35$; 83%), inclusive of all six continents. Publications from European countries contributed to all research topics, with studies from England providing the largest contribution for the fatigue and recovery ($n=6$; 46%) and match-play ($n=4$; 20%) topics, whilst Spain had the largest contribution to physical qualities ($n=18$; 15%). Within




Figure 3. Number of studies published per year investigating the health, performance and development of youth women's footballers.

European countries, the focus of studies from individual countries was typically the investigation of physical qualities (e.g., Spain: $n = 18$; 64%; England: $n = 14$; 45%). However, studies from Scandinavian countries tended to focus on injuries (Sweden: $n = 10$; 50%; Norway: $n = 8$; 50%; Denmark: $n = 6$; 60%). Similarly, publications from North American countries contributed to all research topics, with the USA having the largest contribution to the biomechanics ($n = 20$; 50%), injury ($n = 11$; 22%) and psychology ($n = 10$; 32%) topics, and Canada leading the contribution for nutrition studies ($n = 4$; 29%). Studies from countries in under-represented continents (Africa, Oceania, Asia and South America) predominantly investigated physical qualities.

Participant characteristics

Participant characteristics: competitive standard. The competitive standard of players was reported in most studies ($n = 256$; 87%). Players participating in regional ($n = 124$; 42%) and national ($n = 95$; 32%) competitions were the most investigated, followed by international ($n = 32$; 11%) and local ($n = 30$; 10%) players. When considering competitive standards investigated within research topics (Figure 6), national players were the most investigated in physical qualities ($n = 49$; 41%), nutrition ($n = 7$; 50%) and training load ($n = 4$; 50%) studies, regional players were the most investigated in fatigue and recovery ($n = 11$; 85%), injury ($n = 24$; 49%), match-play ($n = 9$; 45%), and psychology ($n = 15$; 48%) studies, whilst local players were the most investigated in biomechanics studies ($n = 15$; 38%). The over-representation of local players in

biomechanics studies (half of all studies involving local players) may be due to convenience sampling (e.g., ease of accessibility to participants), or logistical considerations for time-intensive, labour-intensive, or inconvenient data collection procedures (e.g., laboratory-based testing as opposed to field-based observational studies).⁴³ International players were predominantly involved in physical qualities studies ($n = 21$; 66%), notably anthropometrics and physical characteristics ($n = 12$; 38%) and studies investigating relative age effect (RAE) within international competitions ($n = 8$; 25%; e.g., FIFA Women's U17 World Cup; UEFA Women's U19 European Championships). The lack of international representation within the literature may be due to the challenge and difficulties in accessing these elite populations, exemplified by all anthropometric and physical characteristics studies involving international players consisting of a single nation/team sample (where stated) likely consequential of convenience sampling (e.g., researchers with applied roles or networks/connections within respective organisations). Furthermore, it is worth noting that no study investigated youth women's para-football formats of any competitive standard. Consequently, there is a clear gap in literature investigating disability formats of youth women's football, which is reflective of the gaps in senior women's football literature.⁹ Future research should focus attention on disability formats of both youth and senior women's football, to start to develop an evidence-base for practitioners working with these players.

Participant characteristics: age-group. Age-group was not stated or indistinguishable for over a third of studies ($n = 108$;




Figure 4. Number of studies published per year investigating the health, performance and development of youth women's footballers according to topic; A) biomechanics, B) fatigue and recovery, C) injury, D) match-play, E) nutrition, F) physical qualities, G) psychology, and H) training load.

38%). The U17 age-group was most investigated ($n = 67$; 23%), followed by U15 ($n = 58$; 20%), U14 ($n = 57$; 19%), and U16 ($n = 55$; 19%) age-groups (Fig. 7). Whilst research focussed upon older age-groups, there was a lower proportion of research at U18 ($n = 41$; 14%) and U19 ($n =$

32; 11%) age-groups which may be due to the exclusion of studies which did not differentiate youth players' data from senior players (e.g., senior, college/university samples). There is a clear lack of research investigating younger youth women's football players (U10: $n = 6$; 2%; U11: $n =$




Figure 5. Geography of studies investigating the health, performance and development of youth women's footballers (grouped by continent; colour).

4; 1%), which may be due to these younger age-groups competing in participation or non-competitive environments (therefore not eligible for inclusion in this review), talent development environments (e.g., youth talent pathway structures; regional and national environments) starting for older age-groups,^{44,45} or researchers' opting (consciously or subconsciously) to focus investigations on older age-groups instead.

Participant characteristics: number of participants. The number of players and teams/clubs participating in studies

within each research topic is presented in Table 1. Most studies ($n = 284$; 97%) reported the number of participating players. Match-play had the highest proportion of studies not reporting the number of participating players ($n = 5$; 25%), which were predominantly studies quantifying heading frequency on a team-level, and therefore, instead reported the number of teams and matches observed.^{46–49} Injury studies typically involved the largest number of participants, which may be unsurprising due to the large-scale epidemiology studies conducted at a league- or tournament-wide level.^{50–52} This was followed by psychology studies,




Figure 6. Competitive standard of players involved in studies investigating the health, performance and development of youth women's footballers (A), according to topic; B) biomechanics, C) fatigue and recovery, D) injury, E) match-play, F) nutrition, G) physical qualities, H) psychology, and I) training load.

where methodological approaches (e.g., questionnaires) facilitated larger sample sizes than those typically reported in observational or intervention-based research in other research topics. Lastly, when removing the studies investigating relative age effect (median = 537; range = 40–12,257), the median and range of participants involved in

physical qualities reduced (median = 34; range = 12–673) to align more closely with the remaining research topics.

Of the 177 studies (60%) which reported the number of teams/clubs involved, 90 (51%) studies investigated single team/club samples. Further, for all research topics except injury, single team/club samples were




Figure 7. Age-groups involved in studies investigating the health, performance and development of youth women's footballers (A), and according to topic; B) biomechanics, C) fatigue and recovery, D) injury, E) match-play, F) nutrition, G) physical qualities, H) psychology, and I) training load.

the most investigated. Given the contextual differences that may exist between teams/clubs (e.g., training and competition structures, provision of/access to sports science and medicine support), results from single team/club samples may not be generalisable to the wider population. Therefore, where possible, future youth

women's football research should adopt a multi-club approach.

Participant characteristics: comparison groups. Common comparisons were between; age-groups ($n = 94$; 32%), including comparisons between youth and senior populations (n

Table 1. Number of players and teams/clubs participating in studies within research topics.

	Number of Players		Number of Teams/Clubs	
	Median (Range)	Not Stated (n; %)	Median (Range)	Not Stated (n; %)
Biomechanics	21 (6–351)	0 (0%)	2 (1–33)	16 (40%)
Fatigue and Recovery	32 (15–93)	0 (0%)	1 (1–1)	9 (70%)
Injury	363 (24–19,147)	3 (6%)	27 (1–341)	14 (29%)
Match-Play	58 (12–828)	5 (25%)	3 (1–74)	3 (15%)
Nutrition	21 (7–205)	0 (0%)	1 (1–3)	8 (57%)
Physical Qualities	40 (12–12,527)	2 (2%)	1 (1–73)	49 (41%)
Psychology	175 (3–1430)	0 (0%)	4 (1–113)	16 (52%)
Training Load	18 (10–259)	0 (0%)	1 (1–1)	2 (25%)

= 28; 10%), genders ($n = 76$; 26%), playing positions ($n = 27$; 9%), and competitive standards ($n = 21$; 7%). Excluding intervention comparisons (e.g., pre- vs post-intervention), a variety of time periods were investigated ($n = 34$; 12%), predominantly between and within training sessions and/or matches ($n = 19$; 6%), and between and within seasons ($n = 15$; 5%). Less frequent comparisons included chronological age ($n = 12$; 4%), maturity status ($n = 5$; 2%), and stage of menstrual cycle ($n = 5$; 2%).

Biomechanics

Forty studies investigated the biomechanics of youth women's football (S1 Table) assessing jumping, landing and/or cutting ($n = 22$; 55%),^{53–74} heading ($n = 14$; 35%),^{75–88} kicking ($n = 2$; 5%),^{89,90} postural control ($n = 1$; 3%)⁹¹ and methodological aspects ($n = 1$; 3%).⁹²

Biomechanics: jumping, cutting and landing. Regional ($n = 7$; 32%),^{55,57,65,66,69,70,72} and local ($n = 6$; 27%)^{53,54,56,62,73,74} players were most commonly investigated across the sub-topic, with only one study (5%) involving national players.⁶⁸ Eight studies (36%) did not state playing standard.^{58–61,63,64,67,71} Common outcome variables included knee kinematics ($n = 16$; 73%)^{55–57,59,60,62,63,65,67–74} and kinetics ($n = 8$; 36%),^{56,57,61,65,69,71,73,74} ground reaction force ($n = 4$; 18%)^{63,64,68,69} and jump performance ($n = 6$; 27%).^{58,62–64,67,72}

Most studies quantified jumping, cutting and/or landing biomechanics using force platforms ($n = 14$; 64%)^{55–58,61,63–66,68,69,71,73,74} and 3D motion analysis ($n = 13$; 59%),^{55–59,61,63,65,68,69,71,73,74} whereas four studies (18%) used 2D video analysis^{62,67,70,72} and four (12%) used IMUs.^{53–55,60} One study used a combination of a force plate and Swift speed mat to determine age-related changes in jump performance.⁶⁴ Studies found sex-specific movement strategies in jumping, cutting and landing movements^{60,65,66} as well as age-related changes in landing asymmetry,⁶⁸ suggesting landing asymmetry reduces with increased age. Intervention studies typically found improvements in lower limb kinematics and kinetics

following implementation of an injury-prevention training programme.^{56,61,67,69,73,74} Younger players showed greater improvements in knee valgus moment during a double leg jump task,⁷⁴ however no other age-related effects following intervention have been reported. Additionally, improved knee kinematics and kinetics were found during an unanticipated side cutting and side-hopping task, after immediate verbal instruction.⁵⁶ While multiple studies showing improvements in kinematics and kinetics following intervention is promising, all studies to date have been conducted in controlled laboratory environments. Therefore, it is unclear whether improvements can translate to training and match scenarios.

Biomechanics: heading. Players of local standard were most commonly investigated across this sub-topic ($n = 8$; 57%),^{78,80,81,84–88} followed by regional ($n = 3$; 21%)^{75,76,79} and national standards ($n = 2$; 14%),^{82,83} with only one study (7%) not reporting playing standard.⁷⁷ All studies bar one quantified head impact kinematics using inertial measurement units, in the form of trackers ($n = 8$; 57%)^{75,76,79,80,82,83,86,87} or mouthpieces ($n = 5$; 36%).^{77,78,81,84,85} Common outcome variables also included total number of headers or head impacts ($n = 9$; 64%),^{75–77,79,81,84,85,87,88} and neck muscle strength ($n = 3$; 21%).^{80,82,87} Two studies (14%) assessed balance and neurocognitive function.^{86,88} Finally, variables were most commonly compared across ball-delivery method or game scenario ($n = 6$; 43%)^{75,77–79,81,84} or age-group ($n = 4$; 29%).^{75–77,85}

Better heading technique (i.e., impact location, technique score) was associated with improved heading kinematics.^{78,79} Head impact magnitudes differed depending on match scenario (e.g., goal kick, corner),^{78,79,81,84,85} and ball characteristics.⁸³ Older players experienced a higher frequency of head impacts (>15 g) than younger players (U14 vs. U12).⁷⁶ Four studies assessed the effects of an intervention on heading kinematics.^{80,82,86,87} These studies predominantly ($n = 3$) improved neck muscle strength, which subsequently led to improvements in heading kinematics.^{80,82,87} Therefore, with guidance and restrictions

being introduced by national governing bodies (UEFA Heading Guidelines for youth players)⁹³ to reduce exposure to repeated head impacts, particularly for younger age-groups, it is essential to ensure that appropriate training (e.g., heading technique, neck strength etc.) is implemented to optimally prepare players for purposeful heading.

Fatigue and recovery

Thirteen studies^{94–106} investigated the fatigue and recovery of youth women's football players (S1 Table), assessing muscular function ($n = 6$; 46%),^{96–99,101,106} nervous system function ($n = 2$; 15%),^{95,100} physical qualities ($n = 2$; 15%),^{102,103} well-being ($n = 2$; 15%),^{94,105} and vascular function ($n = 1$; 8%).¹⁰⁴

A variety of outcome variables were used to investigate fatigue and recovery. Variables assessing neuromuscular function (e.g., muscular delay,⁹⁶ stiffness,⁹⁷ hamstring to quadriceps ratio,⁹⁸ strength and delayed onset of muscle soreness (DOMS),¹⁰¹ knee flexion and valgus angle¹⁰⁶) were quantified using a variety of equipment (e.g., dynamometer,^{96,98,106} RPE,^{101,106} electromyography (EMG),⁹⁶ contact mat,⁹⁷ force platform^{99,106}). Variables assessing nervous system function (e.g., heart rate, heart rate delay,^{95,100}) were quantified using a heart rate monitor¹⁰⁰ or electrocardiogram (ECG) electrodes.⁹⁵

Studies investigated differences between age-groups ($n = 4$; 31%),^{96–98,101} exposure to heading tasks ($n = 2$; 15%),^{95,100} and training load ($n = 1$; 8%),¹⁰⁵ but the majority explored fatigue and recovery across various time periods ($n = 10$; 77%).^{96–99,101–106} Ten studies (77%) investigated fatigue and recovery within an acute period, including; pre- vs post-match,^{99,101–104} pre- vs post-field-based soccer protocol,^{96–98,106} and stage of tournament.^{103,104} Two studies (22%) investigated fatigue and recovery over a longitudinal period, including a 3-month period,¹⁰⁵ and, competitive season⁹⁵).

Following a football-simulation protocol, older age groups had less electromechanical delay,⁹⁶ and greater leg stiffness⁹⁷ compared to younger age-groups. Further, under fatigue, negative effects on hamstring-to-quadriceps ratio were observed for U15 players but beneficial effects for U17 players, particularly at joint angles near full extension where injury risk is higher.⁹⁸ These age-group differences may be due to differences in relative workload, as U15s were found to cover similar total distance in the football-simulation protocol to U17s but in less time (80 min for U15 players compared to 90 min for U17 players). Additionally, U17 players are generally more physically mature, having passed peak height velocity. Maturation is associated with increased neuromuscular control and stability, improved muscle coordination, and enhanced strength.⁹⁸ As a result, more mature players, who are typically better prepared for higher physical demands given an increased playing exposure, are less susceptible to fatigue.

Furthermore, following competitive match-play, U17 players were found to fully recover within a 168-h testing period, while U13 and U15 players continued to exhibit elevated creatine kinase levels and delayed onset muscle soreness beyond this period.¹⁰¹ This indicates greater muscle damage and delayed recovery in younger players, which may suggest that younger age-groups are more susceptible to residual fatigue and potential injury risk if they don't have sufficient recovery time between matches. This may have important implications for fixture scheduling, load management, recovery strategies, and substitution strategies.

The fatigue and recovery of youth women's footballers has had limited investigation, and future research is required to address our gaps in understanding. Future research could look to investigate how relative workload during training and matches affects fatigue-related injury risks and recovery across different age groups, as well as the effectiveness of conditioning programs tailored to specific maturation stages. Due to current research often not accounting for total relative workload completed prior to and during data collection, consideration should be given to how training and fixture schedules of different age groups effects their fatigue and recovery, furthermore, quantifying this (e.g., via GPS) would allow for a more comprehensive analysis. From a practitioner's perspective, consideration needs to be given to the fact that, with more mature players being less susceptible to the effects of fatigue due to more advanced physical qualities, there needs to be a focus on developing the physical qualities of players from earlier age-groups. At the same time, attention could be given to adjusting the volume of training or fixtures of younger age-groups to accommodate for decreased recovery abilities.

Injury

A total of 49 studies investigated injuries within youth women's football (S1 Table), predominantly exploring epidemiology ($n = 22$; 45%)^{50–52,107–125} and risk factors ($n = 17$; 35%),^{126–142} with the remaining studies ($n = 10$; 20%) exploring the efficacy of injury interventions.^{143–152}

Injury: epidemiology. Of the 22 studies which explored epidemiology in youth women's football, there was a range of playing standards investigated (local: $n = 2$, 9%^{108,124}; regional: $n = 6$, 27%^{111,118,120–122,125}; national: $n = 6$, 27%^{107,109,112–114,117}; international: $n = 4$, 18%^{51,52,110,119}; not stated: $n = 4$, 18%^{50,115,116,123}). The most common outcome variables included injury type and location ($n = 18$; 82%),^{50,51,107,109,110,112–124} injury frequency ($n = 17$; 77%),^{50–52,107,109–117,119–121,123,124} match and training exposure or load ($n = 8$; 36%)^{50,107,110,112,115–117,125} and injury severity and duration ($n = 6$; 27%).^{50–52,107,110,114} Practitioners, including coaches and medical professionals,

collected data in 41% ($n=9$) of studies.^{52,107,110,113,114,116,117,119,120} Questionnaires or surveys were used in 41% ($n=9$) of studies^{52,108,111,120–125} and self-report techniques (e.g., text message,¹⁰⁹ self-referral to a medical tent^{50,51,115}) were employed in 18% ($n=4$) of studies. One study used the International Classification of Diseases (ICD-10) alongside X-ray to diagnose injuries.¹¹²

Comparisons of age-group were most common across this sub-topic (60%; $n=13$),^{50–52,107,108,111–114,116,118,123,124} followed by game type (training and matches) ($n=8$; 36%),^{52,108,113–115,121,122,125} injury status (injured and non-injured players) ($n=7$; 32%),^{114,115,118–121,124} playing standard ($n=5$; 23%),^{109,114,118,120,124} and time period ($n=4$; 18%) (year,^{110,115} and (season))^{52,114}). Further comparison groups such as playing position ($n=1$; 6%),¹²² injury type ($n=1$; 5%),¹²³ and injury surveillance type (e.g., parental internet based or certified athletic trainer) ($n=1$; 5%)¹²⁰ featured just once across the sub-topic, therefore warranting further investigation.

Younger age-groups typically experienced highest injury frequency, and thus greater injury risk.^{50,52,111–114,116,118,123} These findings align with existing literature on youth men's players, suggesting that higher injury rates relate to skeletal maturity factors (e.g., proximity to peak height velocity,¹⁵³ skeletal age falling behind chronological age,¹⁵⁴ immature skeleton¹⁵⁵). However, two studies observed injury incidence increased with age.^{51,107}

Studies which compared differences between matches and training presented inconsistent findings, with two reporting injury incidence was highest in training,^{113,122} one observing more injuries in matches⁵² and one reporting no difference.¹²¹ Consideration should be given to the differing methodological approaches adopted within these studies (e.g., collected via survey or practitioners), and the contexts in which these studies occurred. For example, Sprouse et al.⁵² investigated injury incidence during international camps, which is inclusive of competitive matches and training within a congested period (e.g., 7-day period), therefore training and match demands may not reflect those experienced in league-based or domestic environments. Further, as the observation period varied across studies (e.g., whole season,^{107,109,114} tournaments^{50,115,116}), findings from studies observing a shorter time-span cannot be generalised to longer footballing period formats, and vice-versa.

Injury: risk factors. Players within regional competitions were most frequently studied across this sub-topic ($n=11$; 65%)^{127,129–133,138–142} followed by those from national competitions ($n=5$; 29%).^{126,128,134–136} One study did not report playing standard.¹⁴⁰

Common outcome variables included injury frequency ($n=6$; 36%),^{128,135–138,141} injury location ($n=8$; 47%),^{126,129,132,136,138,139,141,142} injury severity (time loss due to injury) ($n=6$; 36%),^{128,129,135,136,138,139} injury

history ($n=4$; 24%),^{126,127,130,131} training load ($n=6$; 36%)^{128,130,131,133,139,140} and physical performance outcomes (e.g., knee strength, jump height, hip and trunk strength,^{127,129,130,134} $\dot{V}O_2$ max^{129,141}). Methods of determining injury characteristics (e.g., injury location, severity, history etc.) and training load of players varied, including practitioner-reported injuries ($n=4$; 24%^{131,132,136,138}), questionnaires ($n=9$; 53%^{126,128,130,132,133,137,139,140,142}), and self-reported injuries ($n=2$; 12%^{135,141}). Questionnaires were also used to assess variables not directly related to injury, such as well-being (Brief-COPE questionnaire),¹⁴⁰ General Health Questionnaire (GHQ-12),^{130,133,140} technical-tactical attributes¹³⁷ and perceptions of early football specialisation.¹⁴² However, self-reporting techniques have limitations, particularly due to reliance on participants' accurate recall of injury detail which may affect accuracy of results.

Injury status was most commonly compared in this sub-topic (53%; $n=9$).^{126,128,130–133,139–141} Typically, injury status reflected current injuries, but four studies specifically examined previous injury as a risk factor for future injury,^{128,130,131,139} concluding that players with prior knee,^{128,131} groin, or ankle¹³⁹ injury had a significantly increased risk of a new injury occurring to the same region. One study reported an increased risk of injuries resulting in time-loss,¹²⁸ although lacked detail regarding type, severity, side and number of previous injuries. Future research should include detailed injury histories to better understand the risk of previous injury on injury re-occurrence. Lastly, due to low ACL and acute knee injury rates (3.5 events), there was low statistical power in risk factor modelling (Cox regression analyses),¹³¹ therefore results from the ACL risk factor model in this study should be interpreted cautiously.

Other comparison groups included playing standard,^{127,137,142} playing surface (grass, artificial turf),^{132,136,138} age-group,¹³⁶ type of sport,¹³⁵ and playing position.¹³² Notably, age-group and playing-position (common comparison groups in other topics) were less utilised across the sub-topic. Future research should explore these further to better understand how maturation and playing position affect risk of injury occurrence and characteristics. Of the three studies exploring playing surface, two found no significant differences in injury frequency between grass and artificial surfaces,^{136,138} while one reported significantly higher training injuries on grass, but no significant differences in matches.¹³² This is in contrast with previous research on senior players which suggests the opposite.^{156–158} This may potentially be due to grass pitches for youth players not being of (or maintained to) as high a standard as senior pitches.

Two of three studies exploring playing standard found that more skilled players had an increased risk of injury,^{137,142} which aligns with previous research in youth men's football,¹⁵⁹ where greater exposure to contact events such as tackles and ariel duels is associated with higher

injury rates. Factors such as greater training volume, intensity, and competitive demands in elite settings may also contribute to increased injury risk and should be considered when interpreting these findings. Future research should account for training experience and load to better isolate the effect of playing standard on injury risk.

Injury: interventions. Ten studies investigated the effects of interventions on injury outcomes in youth women's football (regional: $n = 7$, 70%^{143,144,146,148,149,151,152}; national: $n = 2$, 20%^{145,147} not stated: $n = 1$; 10%).¹⁵⁰ Injury incidence was the most common outcome variable, included in 90% ($n = 9$) of studies.^{144–152} Other outcome variables were more sporadically quantified, for example; compliance or adherence to intervention ($n = 4$; 40%^{144,148–150}), training and match load ($n = 3$; 30%^{144,149,152}), injury location ($n = 3$; 30%^{147,148,152}), injury type ($n = 3$; 30%^{147,148,151}), physical qualities ($n = 3$; 30%^{143,145,149}), and injury severity ($n = 2$; 20%^{147,148}). Data collection of injury characteristics (frequency, type, severity, etc.) was predominantly conducted via practitioners such as coaches, therapists and other medical staff ($n = 5$; 50%),^{146–149,151} followed by questionnaires and surveys ($n = 4$; 40%).^{145,146,150,152}

Interventions varied, including warm-up programs ($n = 4$; 40%),^{148–151} training programs ($n = 2$; 20%),^{143,144} injury prevention strategies ($n = 1$; 10%)¹⁵⁰ and a change in ball ($n = 1$; 10%).¹⁵² The longest intervention was two competitive seasons,¹⁴⁶ however, just over half of the studies had a season-long intervention ($n = 6$; 60%),^{144,147,148,150–152} with others lasting four months ($n = 2$; 20%).^{143,149} Of the two remaining studies, one excluded the intervention group,¹⁴⁵ the other compared the effects of the intervention on trained and un-trained players.¹⁴⁶ Five studies reported significant reductions in injury rates for players in intervention groups,^{143,147,150–152} while two studies found no significant differences in injury rate between the intervention and control groups^{144,148}; however, in one study, low compliance in one specific tertile of the intervention group partially explained the lack of effect,¹⁴⁴ with the other tertile (with a high-compliance rate) demonstrating a significantly lower injury rate than the control group. Comparison of injury intervention effects across comparative groups such as playing position, playing level, game type, fitness level and trained and untrained players were each examined in singular studies within youth women's football. Future research should explore these variables more consistently, particularly by comparing variables such as injury frequency between these comparative groups, to better understand the effectiveness of injury interventions. Furthermore, future research should also explore which variables affect adherence and compliance to injury interventions to inform the design and delivery of interventions (e.g., what is delivered, when and how is it delivered, and by whom?).

Match-Play

Twenty studies quantified the match-play characteristics of youth women's football players (S1 Table). Of those studies, 40% ($n = 8$) investigated physical characteristics,^{41,160–166} 15% ($n = 3$) quantified both physical and technical characteristics,^{167–169} 10% ($n = 2$) quantified only technical characteristics,^{170,171} and the remaining 35% ($n = 7$) examined heading incidence.^{46–49,172–174} No study attempted to quantify the tactical characteristics of youth women's football match-play and therefore future research is warranted in this area.

Match-Play: physical characteristics. Players in regional competitions were the most investigated playing standard in studies across this sub-topic ($n = 4$; 36%),^{162,163,167,168} with two studies (11%) investigating international match-play.^{41,161} Studies examining physical characteristics used a range of variables, such as, heart rate,^{160,169} total distance,^{41,160–169} distance covered in different speed zones,^{41,160–169} and number of accelerations.^{162,163,166} Studies predominantly used GPS devices, albeit of differing frequencies (i.e., 1 Hz,¹⁶⁰ 5 Hz,^{166,169} 10 Hz,^{41,161–164,168} 15 Hz¹⁶⁵), to quantify distance- and speed-based variables, whilst one study used 2D video-based motion analysis.¹⁶⁷ Consistent with research quantifying match-play physical characteristics of senior women's players,¹⁹ there were discrepancies between studies in the speed thresholds and qualitative descriptors for speed zones. Further inconsistencies in the methodological approach were observed for the inclusion criteria for observations (e.g., whole match,^{41,161,163,164,166,169} positional observations,^{162,168} >70 min¹⁶⁷). Therefore, caution is advised when interpreting and comparing results of studies, particularly the distances covered in speed zones.

Studies which involved multiple age-groups ($n = 6$; 55%),^{41,162–164,166,168} quantified age-group specific physical characteristics and consistently observed increasing physical characteristics with increasing age. However, when differing match durations between age-groups were accounted for, age-group differences were less pronounced. Therefore, future research should include both absolute and relative data when investigating match-play physical characteristics across youth-age-groups.¹⁶² Five studies (45%) investigated time-based differences (e.g., match-half/match-periods,^{160,163,166} segmental periods^{164,167}), typically observing reductions in physical performance as the match progresses. All three studies (27%) which quantified position-specific physical characteristics^{41,162,163} observed differences between playing positions. Given that physical characteristics of youth women's football appear to be position-specific, future research should quantify physical match-play characteristics according to playing position and ensure appropriate differentiation of positional categorisation to truly capture position-specific characteristics (e.g., CD and WD^{41,162,163} instead of DEF¹⁶⁶). One study

found different ball characteristics did not influence physical characteristics,¹⁶⁷ whilst one study observed minimal differences in physical characteristics when manipulating the number of players (7v7 vs 8v8), and another reported physical characteristics were dependent upon the possession (i.e., in-possession, out-of-possession, ball-out-of-play) and match status (i.e., drawing, losing or winning).¹⁶⁸

Match-play: technical characteristics. Of the four studies which reported playing standard, three involved players competing in regional competitions and one involved players within national competitions. The frequency of possession-based, offensive, and defensive technical actions during match-play were analysed in studies examining technical characteristics.^{167–171} All five studies used video recordings and notational analysis to collect technical data, likely consequential of the inaccessibility of more advanced data collection systems within youth environments (e.g., optical tracking systems).

Two studies found age-group differences in technical characteristics,^{168,171} one of which also observed position-specific differences within and between age-groups.¹⁷¹ For example, U14 central players performed more technical actions in-possession compared to wide players, whilst technical actions were more evenly distributed across playing positions at U16. The other study found technical actions differed depending upon match status.¹⁶⁸ For example, U14 and U16 age-groups technical data suggested a higher turnover of possession when losing compared to drawing or winning which may be consequential of differing tactical strategies or engaging in high-risk turnover situations (e.g., dribbles, crosses). The findings suggest that age-group-specific coaching practices focussed on technical-tactical aspects of performance may be required (e.g., match strategies, training design, talent development). Additionally, future research should explore the influence of wider contextual factors which have been shown to influence technical characteristics in other populations. For example, contextual factors such as match status and outcome,¹⁷⁵ team and opposition quality,¹⁷⁶ and environmental factors,¹⁷⁷ have all been explored in senior women's football. Lastly, one study found more technical actions were performed by U11 players during a 7v7 match compared to an 8v8.¹⁶⁹ Further research is warranted to understand how manipulating task constraints (e.g., pitch size, number of players, match duration) may influence technical characteristics and how this may vary between age-groups, to assist with informing training design and talent development practices within youth women's football environments.

Match-play: heading. Of the six studies which reported playing standard, four involved regional players whilst two involved national players. All seven studies reported heading frequency during match-play,^{46–49,172–174} with four further quantifying heading characteristics (e.g., impact

location,^{47,48,173} match situation^{47,48,173}). All but one study⁴⁹ conducted post-match notational analysis, instead utilising a live notational analysis approach. The most common finding amongst studies which compared by age group was that heading frequency increased with age.^{47–49,174} Midfielders tended to perform more headers than defenders and attackers regardless of age-group.^{48,172,173} One study also quantified heading frequency during training, and observed significantly more headers were performed during training compared to match-play.¹⁷³ It is important that coaches are made aware of this, so that training can reflect the heading exposure of players in matches, resulting in players only receiving what is deemed to be the necessary level of exposure to improve heading technique. Lastly, none of the studies explored how heading incidence or characteristics may vary across a match, which may have implications for informing policy (e.g., heading rules and regulations), or practice (e.g., substitution strategies).

Nutrition

Fourteen studies^{178–191} explored nutrition surrounding youth women's footballers (S1 Table). Topics varied, with studies assessing energy intake and expenditure,^{178,180,186,187,189} hydration,^{182,184} bone health,^{181,185} food consumption,^{188,190} nutritional status¹⁸³ and disordered eating behaviours.¹⁹¹ Across studies, the most common outcome variables were; body mass measurements (e.g., fat mass, fat free mass, muscle mass (kg, %)).^{178,181–190} Macronutrient measurements were commonly presented in studies exploring energy intake, food consumption and nutritional status.^{178,180,183,185,187,188,190} Outcome variables were compared between age-group,^{180,184,187} playing position,¹⁸⁷ hydration status^{179,182} and dietary condition^{185,188} or between different game days and training sessions.^{179,184,186,187}

Five studies used self-reported food diaries,^{178,186} dietary recall methods^{185,189} or remote food photography¹⁸⁷ along with physical and biochemical assessments, to investigate players' nutritional habits and their impact on health and performance. Results showed that a significant number of players did not meet energy,^{178,180,186,187} carbohydrate,^{178,186,187} protein,^{178,183} or micronutrient requirements, specifically vitamin D, E, A and B12 as well as folate and calcium.^{178,183} These findings, however, should be interpreted with caution as self-reported dietary intake is often inaccurate. Indeed, after adjusting for known under-reporting, one study reported that prevalence of low energy availability dropped from 34% to 5%.¹⁸⁶ Furthermore, prevalence of players with low energy availability significantly reduced (15% vs 5%) when using objective doubly labelled water derived measures compared to self-reported dietary intake.¹⁸⁷ Additionally, all studies had limited sample sizes, either a small sample or one age-group.^{178,186} Despite the associations of energy availability with bone

health, women's football players were reported to have significantly higher bone mineral density compared to non-athletes¹⁸¹ and short-term consumption of Greek yoghurt did not affect bone metabolism,¹⁸⁵ but had a possible effect on the acute anti-inflammatory response during periods of intense training.¹⁸⁸

Two studies investigated hydration using urine specific gravity and found a significant proportion of players were hypohydrated before and during training sessions. Chapelle et al.¹⁷⁹ reported that between 44% and 78% of players were at least minimally hypohydrated during a tournament, with fluid intake insufficient to offset losses, even after an educational intervention. Similarly, Gibson et al.¹⁸⁴ reported that 45% of players arrived at training hypohydrated, and most consumed less than 250 mL of water during sessions, leading to mild dehydration despite moderate sweat losses. These studies were limited by small sample sizes, reliance on urine specific gravity for short-term hydration assessment, and self-reported fluid intake. Additionally, the studies were conducted over brief periods, limiting the understanding of long-term hydration behaviours, and did not account for varying environmental conditions that could impact hydration needs.

One study¹⁹¹ assessed disordered eating behaviours and found that approximately one in six players demonstrated risky eating behaviours. The prevalence was higher in older players (U19s) compared to younger players (U15s). While this study used a larger sample across multiple academies, it was limited by the self-report nature of the EAT-26 and absence of diagnostic follow-up.

These studies highlight significant nutritional gaps among youth women's football players, particularly in energy availability, carbohydrate intake, and micronutrient adequacy. Deficiencies in key nutrients such as iron and vitamin D were prevalent across studies, potentially impacting performance and recovery. Additionally, hydration status was suboptimal, with a high percentage of players starting training sessions in a hypohydrated state. The reliance on self-reported food diaries and dietary recall methods introduces potential biases, particularly under-reporting, which was addressed in some studies by adjusting intake estimates and using objective measures underscoring the importance of accurate data collection. However, the small sample sizes and the short duration of some studies limit the ability to generalize the findings. Further research with larger cohorts and more objective tracking methods is needed to better assess the nutritional practices and their impact on performance and health among youth women's football players.

Physical qualities

Of the 119 studies investigating the physical qualities of youth women's footballers (S1 Table), the majority quantified anthropometrics and physical characteristics ($n = 70$; 59%^{37,192–260}), followed by training interventions ($n = 22$;

18%^{261–282}), relative age effect ($n = 20$; 17%^{283–302} and validity and reliability studies ($n = 7$; 6%^{38303–308}).

Physical qualities: anthropometrics and physical characteristics. Of the 70 studies in this sub-topic, the majority involved regional ($n = 28$; 40%)^{37,193,197,199,202,203,209,212,214,215,219,222,224,225,227,230,233,236,237,242,247,249,250,255,256,258,259} or national ($n = 28$; 40%)^{194,196,198,200,201,204,205,207,213,216–218,221,223,226,227,231,232,234,236,238–240,243,247,253,255,256} standard players, whilst only 12 (17%) and two (3%) studies investigated international^{192,208,220,228,235,236,241,243,244,253,257,260} and local players^{246,252}, respectively. A number of different outcome variables were used, including; jump height or distance ($n = 39$; 56%)^{37,192,195–201,203,204,206,208,210–212,218,221,223,225,227,228,231,233,234,236,239,240,243–245,247,249–251,256,257,260,303} sprint time or distance ($n = 32$; 46%)^{192,195–197,199,200,203–206,208,210–212,226,227,231,232,234,236,239,240,243,244,247,249–251,256–258,260} force ($n = 15$; 21%) (e.g., peak force, peak torque)^{192,194,209–213,220,228,233,235,240,248,255,257} bone health ($n = 8$; 11%) (e.g., bone mineral density or content, bone length bone age)^{214,216,217,224,225,229,230,254} cardiovascular variables ($n = 7$; 10%) (e.g., blood pressure, ventricular wall thickness)^{193,218,237,238,240,241,254} range of motion ($n = 1$; 1%)²²⁰ and menstrual cycle characteristics ($n = 1$; 1%)²⁰⁷. A variety of data collection techniques were deployed across the sub-topic, with 33% ($n = 23$) of studies using a stadiometer (for anthropometric measurements)^{192,194,197,203,208–211,218,221,222,224–226,229,232,234,240,241,251–254} 27% ($n = 19$) using timing gates^{192,196,199,203–205,208,210–212,227,232,236,240,247,250,251,256,258} 17% ($n = 12$) using a dynamometer^{194,200,213,215,219–221,228,233,240,248,255} and 13% ($n = 9$) using skinfold callipers.^{193,202,203,230,231,233,237,244,251} Studies measuring jump height and/or distance used optical measurement systems (e.g., photocells) ($n = 16$; 23%)^{37,192,201,210–212,221,225,227,231,234,239,245,247,249,250} jump, contact or timing mats ($n = 7$; 10%)^{196,203,204,236,243,244,249,256} and force plates ($n = 4$; 6%)^{208,223,260}.

The most common comparative group was age-group with 44% ($n = 31$) of studies comparing variables between at least two age-groups.^{194,201,202,204–206,208–211,213,215,226–229,231,232,235,238,239,241–244,246,252,253,255,256,259} Of these, the U17 age category was the most common amongst studies ($n = 23$), with U10 being the least ($n = 4$).^{209–212,235,253} Two studies worth noting compared variables between seven²³² and eight²⁵⁶ different age-groups respectively. Findings of these studies have general consistency with youth male literature, where jump height and jump distance has been demonstrated to increase across age-groups.^{309–311} Further comparison groups include asymmetry (right and left, or dominant and non-dominant limb)^{37,198,199,206,213,215,239,242,245,248,255} which was used in 16% ($n = 11$) of studies, maturation stage^{193,196,212,234,254} and playing

level^{197,203,236,244,246} were used in five different studies (7%) each. Seven studies (10%) compared variables over a time period, such as pre- and post-season and training period,^{211,217,227,237,240,258,259} all of these studies found significant changes in variables (increased performance) across their compared time period. Only four studies accounted for playing position,^{192,236,246,260} with significant differences only observed between outfield playing positions and goalkeepers. A key limitation was that biological maturation of players over the period of data collection was often not considered, primarily due to cross-sectional research design. Therefore, it is important for future research to compare variables over time (e.g., pre- and post-season). Additionally, it is clear that more research on positional differences in anthropometric and physical characteristics is needed, as the only two studies which did take this approach had a low sample size^{246,260} and so differences between outfield positions were difficult to determine. This may have implications for talent development and talent identification and recruitment practices.

Physical qualities: training interventions. Of the 22 studies investigating the efficacy of training interventions, participants were predominantly players competing in national competitions ($n = 11$; 50%),^{261,262,266,268,269,272,273,275,277,278,281} followed by regional competitions ($n = 7$; 32%),^{264,267,271,276,278,280,282} local competitions ($n = 2$; 9%),^{274,279} and international competitions ($n = 1$; 5%).²⁶³ Common outcome variables included, jump height and distance ($n = 16$; 73%),^{261,263–270,272–275,277,279,280} sprint time and distance ($n = 13$; 59%),^{261,263,265–267,269–273,275,279,282} balance variables generated from the Y-Balance test (e.g., left/right posterolateral movement) ($n = 5$; 23%),^{268–270,272,277} force or strength ($n = 4$; 18%),^{261–263,278} and power (W) ($n = 3$; 14%).^{265,266,276} These variables were commonly quantified using timing gates ($n = 10$; 45%)^{261,263,266,269–272,275,279,282} and optoelectric cell systems ($n = 9$; 41%).^{261,263,265,267,271–273,280,305} Sport-specific items (i.e., footballs) were used in two studies (9%)^{274,275} to quantify kicking distance and kicking velocity respectively.

Different intervention programs were used, including warm up and activation programs ($n = 4$; 184%),^{268,270,277,281} neuromuscular or endurance programmes ($n = 2$; 9%),^{261,262} injury prevention programs ($n = 2$; 9%),^{272,279} transcranial direct current stimulations ($n = 1$; 5%),²⁷⁸ and sprint ($n = 5$; 23%),^{265,271,274,276,282} strength ($n = 5$; 23%),^{263,266,269,272,276} jumping ($n = 2$; 9%)^{264,275} and HIIT/football ($n = 2$; 9%)^{267,280} training programs. The intervention period for the majority of studies ($n = 17$; 77%) lasted between 4–12 weeks.^{261–266,268,270–273,275,276,279,280,282} Three studies (14%) had a season-long intervention duration^{267,269,277} and one (5%) lasted only 2-weeks.²⁷⁸ Most studies ($n = 14$; 64%) compared an intervention group against a control group,^{263,267,268,270–277,279–}

²⁸¹ some compared pre- vs post-intervention ($n = 10$; 45%)^{261–264,271,273,274,278–280} and others the effects of different types of training ($n = 6$; 27%).^{261,262,265,266,269,280} Sixteen studies (73%) observed significant improvements in physical characteristics following an intervention,^{261–263,265–269,271–277,280} two reported improvements in only the dominant limb,^{264,278} whilst four either saw decrements or no significant changes in physical characteristics as a result of an intervention.^{270,279,281,282}

Studies which introduced an intervention alongside players current training made it difficult to determine the effects of the intervention in isolation ($n = 14$; 64%).^{264,265,267–275,277,279,281} Furthermore, studies which did not account for baseline fitness levels saw vast differences in individual player physical characteristics ($n = 4$; 18%).^{264,265,276,278} Therefore, future research should look to introduce interventions outside of players' usual playing seasons. This, alongside establishing baseline fitness levels of participants, may allow for more isolated differences to be established.

Physical qualities: relative age effects. Twenty studies investigated RAE in national ($n = 10$, 50%^{284,286–292,301,302}), international ($n = 7$; 33%^{285,288,293–297}), regional ($n = 5$; 25%^{283,284,298,300,302}) and local ($n = 3$; 15%^{291,299,300}) competitions. Studies predominantly conducted secondary analysis ($n = 18$; 90%) using data from regional ($n = 2$ ^{291,300}), national ($n = 12$ ^{284–286,286,288–290,292,294,294,298,299,301,302}), continental (e.g., UEFA; $n = 1$ ²⁹³) or global governing bodies (i.e., FIFA; $n = 3$ ^{295–297}). The remaining two studies conducted primary data analysis, utilising a questionnaire (no specific details provided), and anthropometric testing.^{283,287} The majority of studies ($n = 18$; 90%) explored birth distribution using birth quartiles^{283–285,287,288,290,292,293,295–302} to investigate RAE, with half year birth distribution ($n = 6$; 33%),^{290–292,298–300} selection probability ($n = 2$; 11%)^{284,285} and predicted adult height ($n = 1$; 6%)²⁸³ used less frequently.

Age-group was the most popular comparative group amongst studies ($n = 14$; 70%),^{283,285,287,289–293,295,296,299,301} followed by competitive level (e.g., regional, national, international; $n = 8$; 40%)^{284,286–288,294,298,300,302} and playing position ($n = 6$; 30%),^{285,286,294,296–298} whilst two studies compared data between geographical location and continental governing bodies (i.e., AFC, CAF, CONCACAF, CONMEBOL, OFC and UEFA^{296,297}). Further comparison groups included: selection status ($n = 2$; 10%),^{283,291} maturity status ($n = 2$; 10%),^{286,287} year of competition ($n = 1$; 5%),²⁹⁵ success of team ($n = 1$; 5%),²⁹⁹ skill rating ($n = 1$; 5%),²⁸⁶ and chronological age ($n = 1$; 5%).³⁰⁰ RAE within different age-groups (U12 to U19) were explored, with the most common being U17 ($n = 12$; 60%)^{285,286,289,290,293–298,301,302} and U19 ($n = 9$; 45%)^{285,286,288–290,293,298,299,302} players. However, no significant

differences in RAE were determined between age-groups in seven studies.^{283,285,287,289,294,299,301} Twelve (60%) of the twenty studies concluded there was a significant over-representation of players born in the first half of the year (birth quartiles Q1 and Q2)^{285,286,290–293,295–298,300,302} compared to the second half (Q3, Q4). However, six studies found no significant RAE.^{283,288,289,294,299,301} The lack of significance in findings could be due to these studies choosing to only assess RAE in older age groups. Females tend to mature earlier compared to males,³¹² and so age-related differences which may directly affect selection rate may be less pronounced in older age groups. Albeit, the two studies which assessed younger age groups (U12^{283,287}; U13²⁸³), observed no RAE. However, these only investigated national players within two isolated countries (Canada, Spain). Future research should focus on investigating RAE across age-groups, considering whether patterns exist in geographical location (e.g., regional, country, continental). Future insights into RAE can help practitioners to mitigate biases in selection processes, which often occur with youth male players,³¹³ therefore developing a more equitable system for supporting the talent development of youth women's football players.

Physical qualities: testing. Testing studies ($n = 7$) assessed the reliability and validity of fitness tests, equipment and data collection techniques. Five studies (71%) assessed performance tests (e.g., multi-stage fitness test,³⁰³ side-hop test,³⁰⁴ anaerobic sprint test³⁸ jumping, sprinting and agility tests^{306,307}). One study assessed the validity of an inertial measurement system (Gyko, Microgate, Italy) and force plate (1000 Hz, Kistler, Switzerland) to determine performance changes across a season,³⁰⁵ and another study determined differential RPE scales were reliable for measurement of training and match load.³⁰⁸ Despite a variety of testing aims being presented, three studies compared age-group,^{304,306,307} whilst two studies compared data collection technique.^{303,305} Playing status³⁸ and training type³⁰⁸ were stand-alone comparison groups in the other two studies. None of the seven studies (where it was applicable) used playing position as a comparative group, this with four studies reporting limited sample size by four studies.^{38,106,303,305} Multiple studies only included one age-group,^{38,106,303,305} limiting the generalisability of findings to the wider youth women's football population. Further limitations of studies included assumptions made on limb dominance,³⁰⁴ an inability to determine maximum effort from participants in fitness tests.³⁰⁷ For the study assessing equipment validity, the inertial measurement device was not intended for stand-alone use and so findings may be inaccurate.³⁰⁵ Future research should aim to adopt a larger sample size and include competitive standard, age-group, and playing position as comparative groups to determine potential validity and reliability differences across the youth women's football population.

Psychology

Thirty-one studies investigated psychology surrounding youth women's footballers (S1 Table).^{314–344} Studies most frequently examined motivational climate ($n = 6$; 21%)^{329–333,336} and motor development and cognitive function ($n = 9$; 29%).^{314,315,321–324,327,337,341} As a result, most outcome variables were psychometric scales representing anxiety,^{316,326,333,340,342} moral functioning,^{344,330,343} leadership,^{319,338} self-confidence,^{316,333} motivation,^{321,328,335,338} coach-athlete relationship,³²⁸ and self-talk.^{316,317} These latent constructs were operationalised using several different questionnaires and scales such as: The Psychological Skills Inventory motivation subscale,^{321,345} Perceived Motivational Climate in Sport Questionnaire,^{329–331,333,336,342,346} the Sport Motivation Scale³²⁸ and the Task and Ego Orientation in Sport Questionnaire^{316,317,344,343,346} to measure motivational constructs. The Go/No-Go Task^{314,315,347} was used to measure variables representing motor development (e.g., reaction time to determine impulsiveness). Outcome variables were most commonly compared across age-group ($n = 5$; 18%),^{315,320,326,327,343} time period (e.g., pre- and post-game) ($n = 4$, 14%)^{314,318,320,334} and game or session stage ($n = 3$; 11%).^{322,339,340}

Studies exploring cognitive function and motor development assessed: cognitive function over time,³¹⁴ the validation of soccer competence scale³²¹ and whether motor performance is a predictor for footballing success in adulthood.³²⁷ Therefore, comparisons cannot be made as they deal with distinct topics. Contrastingly, studies exploring motivational climate are more comparable, with most establishing relationships between motivational climate and other characteristics and traits such as cognitive anxiety,³³³ peer acceptance,³³⁶ and sporting behaviour,³²⁹ determining that higher levels of intrinsic or self-determined motivation are positively related to biopsychosocial outcomes of football involvement. Age-group specific results were presented in five (18%) studies,^{315,320,326,327,343} however direct comparisons between them were only made in two.^{315,343}

None of the 31 studies used a control group of a non-athletic comparison group to establish psychological changes due to engaging in playing football or a football related training program or intervention. Therefore, future research could consider comparative designs to establish the differences or similarities between women's footballers and their non-sporting or other sporting peers. To a similar end, a more consistent approach to examining age related differences would establish developmental changes in the psychological profile of women footballers. Lastly, future research could explore performance-related psychological variables, such as visual exploratory activity (VEA) of players during training or matches³⁴⁸ and explore whether VEA of youth women's football players may differ

according to age-group, playing position or the influence of contextual factors (e.g., pitch location). This may have implications for talent identification and development practices and informing training design and delivery.

Training load

Eight studies quantified the training load of youth women's footballers (S1 Table).^{349–356} These studies involved participants competing in international competitions,^{349,351} national competitions^{353,354}; Rumpf et al.³⁵⁵ also included local and regional players, whilst Williams et al.³⁵⁶ and Idarraga and Valencia-Sánchez³⁵² only included regional players. The eight studies had distinct aims for investigating training load within youth women's football, including; describing seasonal variations in training, anthropometry, body composition and physical fitness,³⁵³ comparing physical exposure, wellness and psychological variables during domestic and international playing period,³⁴⁹ quantifying differences in internal and external load between starters and substitutes,³⁵¹ investigating training profiles and motivation across different age-groups and competitive standards,³⁵⁵ determining the prevalence of non-functional overreaching and overtraining in elite youth players,³⁵⁶ comparing the influence of obstacles in SSGs on tactical, physical and emotional responses of players,³⁵⁴ evaluating a passing-skill training intervention,³⁵⁰ and evaluating the effectiveness of psychokinetic games on tactical creativity.³⁵² Consequently, methodological approaches for data collection and outcome variables varied across the eight studies. Given the size of the current training load evidence base, and the range of aims, methodological techniques and findings, it is difficult to interpret and summarise this area of literature on youth women's football. As such, the focus for future research is broad. Further research needs to develop our understanding of the training demands in different age-groups and competitive standards within youth women's football environments.³⁰⁸ Further, it is important to understand how representative these training environments are to competition or whether they are providing sufficient stimulus for physiological adaptations and athletic development, and therefore similar metrics should be investigated to those quantified for understanding the demands of match-play.^{162,164,41} Further, understanding whether training demands are appropriately preparing players for transitioning between youth age-groups (e.g., given the age-related increases in match demands within youth female football^{162,164}) or into senior female football environments.^{357–360} Lastly, investigating how training demands may differ according to drill type (as opposed to simply quantifying whole-session demands), could help coaches inform the design and delivery of activities within training sessions (e.g., age-group specific benchmarks).³⁵⁹

Limitations

This scoping review adhered to the PRISMA-ScR Checklist,³⁰ assessing several electronic databases, scoping studies with no date nor language restrictions, and utilising a broad range of search terms inclusive of performance, health and development. Further, two authors independently screened studies at both initial and full-text stage, with all extracted data checked independently by another member of the research team. As such, this study has comprehensively reviewed and summarised the scientific literature investigating the performance, health and development of youth women's football players. However, there are limitations which should be acknowledged. Firstly, whilst no language restrictions were imposed, the search terms were predominantly written in English, and therefore relevant studies written in non-English may have been excluded from this review. Secondly, the search strategy did not include grey literature, and therefore, this review does not consider unpublished (or non-peer-reviewed) evidence relating to the performance, health and development of youth women's football. Whilst population search terms related to youth and adolescents, some studies may have used U (e.g., U12, or U16) as their descriptor, and thus, these studies may not have been captured. Thirdly, due to the nature of this review and the quantity of eligible studies, it was not practically feasible to manually screen the reference list of included studies within the review. Therefore, it may be possible that additional eligible studies could have been identified through this process. Fourthly, studies were systematically categorised into topics based on their primary research aim. However, there may have been studies where secondary or tertiary aims could have been categorised into alternative topics. Lastly, there were a notable proportion of studies which either poorly reported or did not state key information (e.g., number of participants, number of teams/clubs represented within the sample, age-group(s) or competitive standard). Therefore, when conducting data extraction, this resulted in notable data reported as not stated (NS). Therefore, the distribution of studies within the literature may be misrepresented within the results of this scoping review (e.g., U17 may not have been the most investigated age-group). Future research across all topics should ensure methods are clearly detailed to facilitate understanding of the approach adopted, and the individuals participating within studies.

Implications for future research & applied practice

Throughout this scoping review, gaps in literature have been identified within respective topics and recommendations made for future research, both from the perspective of developing knowledge and understanding, and with regards to methodological approaches adopted.

Overall, there is a lack of research investigating international youth women's football players, and future research is warranted to understand the performance, health and development of players competing at the international standard. This insight would be useful for coaches and practitioners supporting and preparing players for involvement across competitive standards (e.g., domestic to international), and for understanding age-related differences which may exist (e.g., youth international to senior international environments) to aid transition and progression between environments. Overall, there is a lack of research which has consistently investigated multiple age-groups across youth women's football, instead primarily focussing on isolated age-groups, and particularly investigating U14–U17 age-groups. There is a need for research to investigate across age-groups, to understand potential age-related differences or considerations for applied practice, particularly due to the differences which can be placed on these age-groups from a talent development perspective (e.g., long-term talent development: learning to train, training to train, training to compete, training to win). There is no research which has investigated youth women's para-football formats, and therefore, a warranted area for future research. Importantly, the evidence-base was predominantly from Western European and North American countries, and thus translation and application for youth women's populations in other countries may be inappropriate (e.g., differing talent development pathways, training and competition structures, playing styles, provision and support). Therefore, greater diversity and representation within future research would facilitate a more global understanding of youth women's football, and aid translation and application within populations in currently under-represented countries (i.e., South American, Asian, African, Oceanic).³⁶¹ Further, global, continental, and national governing bodies should prioritise supporting performance, health and development research within youth women's populations (i.e., in addition to senior women's populations, and participation-focussed research in youth populations), whilst also attempting to address key challenges in facilitating inclusive and diverse research (e.g., funding, resources, access to expertise, socio-cultural expectations of youth women's football).

Studies across all topics (except injury) predominantly involved single team/club samples, and future research should consider collaborative research (e.g., with national governing bodies) who can facilitate recruitment and participation for enabling multi-club approaches (e.g., league-wide) to understand potential team/club variation and facilitate more generalisable insights into youth women's football. There are discrepancies in the number of studies within respective topics, and regardless of methodological inconsistencies or considerations that exist and limit the transfer of knowledge or application findings within topics, the least investigated areas of research were training load,

fatigue and recovery and nutrition. However, the body of research has predominantly investigated the performance, health and development of youth women's football in isolated topics. Therefore, to gain a more holistic understanding, future research should aim to adopt a more interdisciplinary approach to investigating the performance, health and development of the youth women's footballer.

This scoping review has provided a comprehensive summary of the current evidence-base within youth women's football, identifying gaps and recommending areas for future research. However, it is important that research attempts to bridge the gap with practitioners, setting an agenda for research investigating the performance, health and development within youth women's football which accounts for the needs of those working with this population. Therefore, the next steps are to engage with wider experts (academic, research, and applied practitioners) by conducting a Delphi study to gain consensus on the required areas for future research, using this scoping review as the basis for current understanding.

Conclusion

This study has scoped the peer-reviewed scientific literature investigating the performance, health and development of youth women's footballers. A total of 294 studies were included, and categorised into eight research topics: biomechanics, fatigue and recovery, injury, match-play, nutrition, physical qualities, psychology and training load. The most researched area was physical qualities, whilst the most investigated players were those competing in regional competitions and those competing in an U17 age-group. The study provides a comprehensive, critical and objective resource of the scientific research involving youth women's soccer players, which can be used to inform applied practices within this population. By determining the methodological approaches adopted, summarising the key findings and identifying the gaps within scientific literature, recommendations have been provided for future research. Further, the current needs, questions or challenges facing applied practitioners working within youth women's football in relation to our current understanding from the scientific literature is required, to establish consensus on priorities for future research.

ORCID iDs

Sally Waterworth  <https://orcid.org/0000-0002-9955-4497>

Tracy Lewis  <https://orcid.org/0000-0002-6493-2072>

Naomi Datson  <https://orcid.org/0000-0002-5507-9540>

Carla Harkness-Armstrong  <https://orcid.org/0000-0002-6093-365X>

Alice Harkness-Armstrong  <https://orcid.org/0000-0002-7258-4469>

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Data availability statement

The authors confirm that the data supporting the findings of this study are available within its supplementary materials.

Supplemental material

Supplemental material for this article is available online.

References

1. Fédération Internationale de Football Association (FIFA). Women's football strategy: 2024 – 2027. In: *Women's football strategy: 2024 – 2027 [Internet]*. 2024 [cited 13 May 2025]. Available: <https://inside.fifa.com/womens-football/strategy>
2. Oceania Football Confederation (OFC). All in: OFC women's football strategy 2027. 2021. Available: <https://www.oceaniafootball.com/cms/wp-content/uploads/2021/07/All-IN-OFC-Womens-Football-Strategy-compressed.pdf>
3. Union of European Football Associations (UEFA). UEFA women's football strategy 2024–2030: Unstoppable. 2024. Available: https://editorial.uefa.com/resources/0292-1c3532529b72-3306578495fb-1000/unstoppable_uefa_women_s_football_strategy.pdf
4. The Football Association (FA). Reaching higher: Our 2024–2028 women's and girls' strategy. 2024. Available: <https://www.thefa.com/-/media/thefacom-new/files/about-the-fa/2024/fa-womens-girls-football-strategy-202428.ashx?la=en>
5. Irish Football Association (Irish FA). Together we thrive: elevating the game for everyone. Strategic plan for women's & girls' football 2025–2029. 2024. Available: <https://www.irishfa.com/media/57554/together-we-thrive.pdf>
6. Fédération Internationale de Football Association (FIFA). Women's football member associations survey report 2023. 2023. Available: <https://digitalhub.fifa.com/m/28ed34bd888832a8/original/FIFA-Women-s-Football-MA-Survey-Report-2023.pdf>
7. Fédération Internationale de Football Association (FIFA). FIFA benchmarking report women's football: Setting the pace 2024 edition. 2024. Available: https://digitalhub.fifa.com/m/4e63f61e61a63a93/original/FIFA-Womens-Football-Benchmarking-Report_EN.pdf
8. Fédération Internationale de Football Association (FIFA). FIFA women's champions Cup™ and FIFA women's club world Cup™ announced. 2025 [cited 13 May 2025]. Available: <https://www.fifa.com/en/news/articles/womens-champions-cup-club-world-cup-timelines-formats>
9. Kryger K, Wang A, Mehta R, et al. Research on women's football: a scoping review. *Sci Med Footb* 2022; 6: 549–558.
10. Fédération Internationale de Football Association (FIFA). Introduction to the FIFA talent development scheme. In: *FIFA Training Centre [Internet]*. 2022 [cited 1 Dec 2025]. Available: <https://www.fifatrainingcentre.com/en/environment/talent-development/talent-development/introduction-to-the-fifa-talent-development-scheme.php>
11. Fédération Internationale de Football Association (FIFA). One region, one goal: FIFA talent academies strengthen football in Concacaf. In: *FIFA Training Centre [Internet]*. 2025 [cited 1 Dec 2025]. Available: <https://www.fifatrainingcentre.com/en/environment/news/high-performance/one-region-one-goal.php>
12. Union of European Football Associations (UEFA). UEFA women's football development programme: National association projects. 2020. Available: https://editorial.uefa.com/resources/025f0fff0dce7b19-5db19c89f686-1000/uefa_wfdp_national_association_projects_2020.pdf
13. Confédération Africaine de Football (CAF). CAF launches CAF IMPACT Programme to promote sound financial Governance, Youth Competitions and Community Outreach with Member Associations. [cited 1 Dec 2025]. Available: <https://www.cafonline.com/inside-caf/news/caf-launches-caf-impact-programme-to-promote-sound-financial-governance-youth-competitions-and-community-outreach-with-member-associations/>
14. The Football Association (FA). Women's and girls' football sees record growth in a historic four years for the game. In: *www.thefa.com [Internet]*. 2024 [cited 1 Dec 2025]. Available: <http://www.thefa.com/news/2024/sep/27/women-and-girls-stratgy-progress-inspiring-positive-change-20242709>
15. Football Association Wales (FAW). National AcademiFAW National Academi: Unearthing talent and connecting football [Internet] 2025 [cited 1 Dec 2025]. Available: <https://faw.cymru/national-academi/>
16. Ghana Football Association. Ghana advances Women's Football Development with expanded national team structure. 2024 [cited 13 May 2025]. Available: <https://www.ghanafa.org/ghana-advances-womens-football-development-with-expanded-national-team-structure>
17. Scottish Football Association (Scottish FA). New girls' talent centres to help identify future Scotland stars. 2024 [cited 13 May 2025]. Available: <https://www.scottishfa.co.uk/news/new-girls-talent-centres-to-help-identify-future-scotland-stars/>
18. In numbers: Premier League's support for wider football. 2025 [cited 13 May 2025]. Available: <https://www.premierleague.com/news/4237126>
19. Harkness-Armstrong A, Till K, Datson N, et al. A systematic review of match-play characteristics in women's soccer. *PLoS One* 2022; 17: e0268334.
20. Hamed D, Gonzalez-Muñoz A, Cuevas Cervera M, et al. Effects of the menstrual cycle on the performance of female

- football players. A systematic review. *Front Physiol* 2024; 15: 1359953.
21. Mulvenna C, Evans N, Brady A, et al. A scoping review of the research supporting coaching practice in women's football – as the game grows the research strives to keep up. *Sci Med Footb* 2025; 1–10. doi:10.1080/24733938.2025.2488525
 22. Costa JA, Rago V, Brito P, et al. Training in women soccer players: a systematic review on training load monitoring. *Front Psychol* 2022; 13: 943857.
 23. Lopez-Valenciano A, Raya-González J, Garcia-Gomez J, et al. Injury profile in women's football: a systematic review and meta-analysis. *Sports Med* 2021; 51: 423–442.
 24. Pettersen SD, Adolfsen F and Martinussen M. Psychological factors and performance in Women's football: A systematic review. *Scand J Med Sci Sports* 2021; 32: 161–175.
 25. Emmonds S, Heyward O and Jones B. The challenge of applying and undertaking research in female sport. *Sports Med – Open* 2019; 5: 51.
 26. Bonito FP, Emmonds S, Teles J, et al. Growth and maturation in youth female football players and its association with body composition and physical performance: a scoping review. *Int J Sports Sci Coach* 2025; 17479541251380019. doi:10.1177/17479541251380019
 27. Malina RM, Rogol AD, Cumming SP, et al. Biological maturation of youth athletes: assessment and implications. *Br J Sports Med* 2015; 49: 852–859.
 28. Ramachandran AK, Pedley JS, Moeskops S, et al. Changes in lower limb biomechanics across various stages of maturation and implications for ACL injury risk in female athletes: a systematic review. *Sports Med Auckl NZ* 2024; 54: 1851–1876.
 29. Butcher AJ, Ward S, Clissold T, et al. Maturation and biomechanical risk factors associated with anterior cruciate ligament injury: is there a link? A systematic review. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2024; 68: 31–50.
 30. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018; 169: 467–473.
 31. Whitehead S, Weakley J, Cormack S, et al. The applied sports science and medicine of netball: a systematic scoping review. *Sports Med* 2021; 51: 1715–1731.
 32. Peters MDJ, Godfrey CM, Khalil H, et al. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc* 2015; 13: 141–146.
 33. Sacks D, Canadian Paediatric Society, Adolescent Health Committee. Age limits and adolescents. *Paediatr Child Health*. 2003;8: 577.
 34. Bangsbo J. Performance in sports—with specific emphasis on the effect of intensified training. *Scand J Med Sci Sports* 2015; 25: 88–99.
 35. Cueto M. The world health organization. In: MCB Raviglione, F Tediosi, S Villa, N Casamitjana and A Plasència, editors. *Global health essentials*. Cham: Springer International Publishing; 2023, pp. 421–424. doi:10.1007/978-3-031-33851-9_63
 36. The Editors of Encyclopaedia Britannica. Biological development summary | Britannica. [cited 12 Feb 2025]. Available: <https://www.britannica.com/summary/biological-development>
 37. Mollinedo-Gómez L, Osmani F and Lago-Fuentes C. Asimetrías en jugadores alevines de fútbol de una cantera de élite: diferencias entre sexos. *Sport Sci J Sch Sport Phys Educ Psychomot* 2023; 9: 545–562.
 38. Jastrzebska A. Comparison of usefulness of two tests measuring anaerobic performance of untrained and soccer-training girls U12. *Sci Rep* 2023; 13: 19498.
 39. Nayiroğlu S, Yılmaz AK, Silva AF, et al. Effects of small-sided games and running-based high-intensity interval training on body composition and physical fitness in under-19 female soccer players. *BMC Sports Sci Med Rehabil* 2022; 14: 119.
 40. Faustino-da-Silva YV, Martinho DV, Coelho-E-Silva MJ, et al. Reproducibility and inter-observer agreement of greulich-Pyle protocol to estimate skeletal age among female adolescent soccer players. *BMC Pediatr* 2020; 20: 494.
 41. Ramos GP, Nakamura FY, Penna EM, et al. Activity profiles in U17, U20, and senior women's Brazilian national soccer teams during international competitions: are there meaningful differences? *J Strength Cond Res* 2019; 33: 3414–3422.
 42. CIES. Worldwide demographic analysis of women's football – CIES football observatory *Football Observatory – Monthly Report [Internet]*. 2024 [cited 12 Feb 2025]. Available: <https://football-observatory.com/MonthlyReport96>.
 43. Glazier PS and Mehdizadeh S. Challenging conventional paradigms in applied sports biomechanics research. *Sports Med Auckl NZ* 2019; 49: 171–176.
 44. Andrew M, Ford PR, Alder SE, et al. Talent development in female soccer: developmental activities of professional players in England. *J Sports Sci* 2024; 42: 853–864.
 45. McEwan GP, Unnithan VB, Carter M, et al. Talent identification and development strategies in elite women's soccer: a pan-European perspective. *Sci Med Footb* 2024; 1–9. doi:10.1080/24733938.2024.2404920
 46. Beaudouin F, Gioftsidou A, Larsen MN, et al. The UEFA heading study: heading incidence in children's and youth' football (soccer) in eight European countries. *Scand J Med Sci Sports* 2020; 30: 1506–1517.
 47. Harriss A, Johnson AM, Walton DM, et al. The number of purposeful headers female youth soccer players experience during games depends on player age but not player position. *Sci Med Footb* 2019; 3: 109–114.
 48. Peek K, Vella T, Meyer T, et al. The incidence and characteristics of purposeful heading in male and female youth football (soccer) within Australia. *J Sci Med Sport* 2021; 24: 603–608.
 49. Sandmo SB, Andersen TE, Koerte IK, et al. Head impact exposure in youth football—are current interventions hitting the target? *Scand J Med Sci Sports* 2020; 30: 193–198.

50. Andreassen I, Faunø P, Lund B, et al. Soccer injuries among youth. *Scand J Med Sci Sports* 1993; 3: 62–66.
51. Schmidt-Olsen S, Bünnemann LK, Lade V, et al. Soccer injuries of youth. *Br J Sports Med* 1985; 19: 161–164.
52. Sprouse B, Alty J, Kemp S, et al. The football association injury and illness surveillance study: the incidence, burden and severity of injuries and illness in men's and women's international football. *Sports Med* 2020; 54: 213–232.
53. Alanen A-M, Barrons Z, Jordan MJ, et al. Between-day reliability of inertial measurement unit parameters during soccer-specific change of direction test. *Int J Sports Sci Coach* 2023; 19: 1050–1057.
54. Alanen A-M, Benson LC, Jordan MJ, et al. Effects of game-specific demands on accelerations during change of direction movements: analysis of youth female soccer. *Biomechanics* 2023; 3: 250–257.
55. Benjaminse A, Nijmeijer EM, Gokeler A, et al. Application of machine learning methods to investigate joint load in agility on the football field: creating the model, part I. *Sensors* 2024; 24: 3652.
56. Celebrini R, Eng J, Miller W, et al. The effect of a novel movement strategy in decreasing ACL risk factors in female adolescent soccer players. *J Strength Cond Res Natl Strength Cond Assoc* 2012; 26: 3406–3417.
57. Celebrini RG, Eng JJ, Miller WC, et al. Effect of a novel movement strategy in decreasing ACL risk factors in female adolescent soccer players: a randomized controlled trial. *Clin J Sport Med Off J Can Acad Sport Med* 2014; 24: 134–141.
58. DeLang MD, Hannon JP, Goto S, et al. Female adolescent soccer players utilize different neuromuscular strategies between limbs during the propulsion phase of a lateral vertical jump. *Int J Sports Phys Ther* 2021; 16: 695–703.
59. DiCesare CA, Kiefer AW, Bonnette S, et al. High-Risk lower-extremity biomechanics evaluated in simulated soccer-specific virtual environments. *J Sport Rehabil* 2020; 29: 294–300.
60. Di Paolo S, Gokeler A, Benjaminse A, et al. On-field kinematics of cut maneuvers in football players: are wearable sensors reliable for assessing anterior cruciate ligament injury risk? *J Sports Sci* 2025; 43: 1256–1266.
61. Ford KR, Taylor JB, Westbrook AE, et al. Outcomes of a randomized controlled trial of neuromuscular training with real-time biofeedback in young female athletes. *BMC Musculoskelet Disord* 2025; 26: 366.
62. Grandstrand SL, Pfeiffer RP, Sabick MB, et al. The effects of a commercially available warm-up program on landing mechanics in female youth soccer players. *J Strength Cond Res* 2006; 20: 331–335.
63. Grooten W, Karlefur O and Conradsson D. Effects of verbal knee alignment instructions on knee kinematics, kinetics and the performance of a single-leg jump in female adolescent soccer players. *Eur J Physiother* 2020; 22: 106–114.
64. Jeras N, Bovend'Eerd T and McCrum C. Biomechanical mechanisms of jumping performance in youth elite female soccer players. *J Sports Sci* 2020; 38: 1335–1341.
65. Landry SC, McKean KA, Hubley-Kozey CL, et al. Neuromuscular and lower limb biomechanical differences exist between male and female elite adolescent soccer players during an unanticipated side-cut maneuver. *Am J Sports Med* 2007; 35: 1888–1900.
66. Landry S, McKean K, Hubley-Kozey C, et al. Gender differences exist in neuromuscular control patterns during the pre-contact and early stance phase of an unanticipated side-cut and cross-cut maneuver in 15-18 years old adolescent soccer players. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 2009; 19: e370–e379.
67. Lindblom H, Waldén M, Carlffjord S, et al. Limited positive effects on jump-landing technique in girls but not in boys after 8 weeks of injury prevention exercise training in youth football. *Knee Surg Sports Traumatol Arthrosc* 2020; 28: 528–537.
68. Lucarno S, Zago M, Rossi E, et al. Influence of age and sex on drop jump landing strategies in elite youth soccer players. *Int J Sports Sci Coach* 2021; 16: 344–351.
69. Nijmeijer EM, Kempe M, Elferink-Gemser MT, et al. Practice, and improve? Enhancing sidestep cutting execution in talented female soccer players: a four-week intervention program with video instruction. *J Strength Cond Res* 2024; 38: e430–e439.
70. O'Kane JW, Tencer A, Neradilek M, et al. Is knee separation during a drop jump associated with lower extremity injury in adolescent female soccer players? *Am J Sports Med* 2016; 44: 318–323.
71. Ortiz A, Trudelle-Jackson E, McConnell K, et al. Effectiveness of a 6-week injury prevention program on kinematics and kinetic variables in adolescent female soccer players: a pilot study. *P R Health Sci J* 2010; 29: 40–48.
72. Richardson MC, Evans W, Chesterton P, et al. The effects of a 6-week sand- vs. land-based jump training programme on frontal plane knee angle and jump performance in adolescent female football players. *J Sports Sci* 2025; 43: 523–535.
73. Thompson JA, Tran AA, Gatewood CT, et al. Biomechanical effects of an injury prevention program in preadolescent female soccer athletes. *Am J Sports Med* 2017; 45: 294–301.
74. Thompson-Kolesar JA, Gatewood CT, Tran AA, et al. Age influences biomechanical changes after participation in an anterior cruciate ligament injury prevention program. *Am J Sports Med* 2018; 46: 598–606.
75. Brooks JS, Allison W, Harriss A, et al. Purposeful heading performed by female youth soccer players leads to strain development in deep brain structures. *Neurotrauma Rep* 2021; 2: 354–362.
76. Chrisman SPD, Ebel BE, Stein E, et al. Head impact exposure in youth soccer and variation by age and sex. *Clin J Sport Med Off J Can Acad Sport Med* 2019; 29: 3–10.
77. Filben TM, Pritchard NS, Miller LE, et al. Header biomechanics in youth and collegiate female soccer. *J Biomech* 2021; 128: 110782.

78. Filben TM, Tomblin BT, Pritchard NS, et al. Assessing the association between on-field heading technique and head impact kinematics in a cohort of female youth soccer players. *Sci Med Footb* 2023; 8: 341–350.
79. Harriss A, Johnson AM, Walton DM, et al. Head impact magnitudes that occur from purposeful soccer heading depend on the game scenario and head impact location. *Musculoskelet Sci Pract* 2019; 40: 53–57.
80. Le Flao E, Pichardo AW, Ganpatt S, et al. An accessible, 16-week neck strength training program improves head kinematics following chest perturbation in young soccer athletes. *J Sport Rehabil* 2021; 30: 1158–1165.
81. Miller LE, Pinkerton EK, Fabian KC, et al. Characterizing head impact exposure in youth female soccer with a custom-instrumented mouthpiece. *Res Sports Med Print* 2020; 28: 55–71.
82. Peek K, Andersen J, McKay MJ, et al. The effect of the FIFA 11 + with added neck exercises on maximal isometric neck strength and peak head impact magnitude during heading: a pilot study. *Sports Med Auckland NZ* 2022; 52: 655–668.
83. Peek K, McKay M, Fu A, et al. The effect of ball characteristics on head acceleration during purposeful heading in male and female youth football players. *Sci Med Footb* 2021; 5: 195–203.
84. Pritchard S, Filben T, Haja S, et al. Head impact exposure in youth soccer: comparing across activity types. *Proc Inst Mech Eng Part P J Sports Eng Technol* 2025; 239: 533–544.
85. Tomblin BT, Pritchard NS, Filben TM, et al. Characterization of on-field head impact exposure in youth soccer. *J Appl Biomech* 2021; 37: 36–42.
86. Wahlquist VE and Kaminski TW. Analysis of head impact biomechanics in youth female soccer players following the get aHEAD safely in Soccer™ heading intervention. *Sensors*. 2021;21: 3859.
87. Wahlquist VE, Glutting JJ and Kaminski TW. Examining the influence of the get aHEAD safely in Soccer™ program on head impact kinematics and neck strength in female youth soccer players. *Res Sports Med Print* 2022; 32: 17–27.
88. Wahlquist VE, Buckley TA, Caccese JB, et al. Youth soccer heading exposure and its effects on clinical outcome measures. *Sports* 2024; 12: 342.
89. Khatun M, Boby FA, Sanjaykumar S, et al. Gender disparities in long passing performance among senior football players: implications for inclusive training strategies. *Retos* 2024; 60: 886–893.
90. Lyle MA, Sigward SM, Tsai L-C, et al. Influence of maturation on instep kick biomechanics in female soccer athletes. *Med Sci Sports Exerc* 2011; 43: 1948–1954.
91. Przysucha E, Zerpá C, Vidotto E, et al. Postural sway characteristics in static balance control of youth female soccer players: NA. *Acta Kinesiol* 2024; 18: 61–68.
92. Di Paolo S, Nijmeijer E, Bragonzoni L, et al. Comparing lab and field agility kinematics in young talented female football players: implications for ACL injury prevention. *Eur J Sport Sci* 2023; 23: 859–868.
93. The Football Association. Heading in football. <https://www.Englandfootball.com> [Internet]. [cited 12 Feb 2025]. Available: <https://www.Englandfootball.com/participate/learn/brain-health/heading-in-football>
94. Ahmedi M, Sitko S and Mayolas-Pi C. The influence of competitive level on sleep quality and quality of life in adolescent soccer players. *Cureus* 2024; 16: e71395.
95. Bonn MM, Harriss AB, Thompson JWG, et al. Performing more than 20 purposeful gameplay headers in a soccer season may alter autonomic function in female youth soccer players. *Res Sports Med Print* 2021; 29: 440–448.
96. De Ste Croix MBA, Priestley AM, Lloyd RS, et al. ACL Injury risk in elite female youth soccer: changes in neuromuscular control of the knee following soccer-specific fatigue. *Scand J Med Sci Sports* 2015; 25: e531–e538.
97. De Ste Croix MBA, Hughes JD, Lloyd RS, et al. Leg stiffness in female soccer players: intersession reliability and the fatiguing effects of soccer-specific exercise. *J Strength Cond Res* 2017; 31: 3052–3058.
98. De Ste Croix M, Priestley A, Lloyd R, et al. Age-related differences in functional hamstring/Quadriceps ratio following soccer exercise in female youth players: an injury risk factor. *Pediatr Exerc Sci* 2018; 30: 376–382.
99. Fahey JT, Comfort P and Ripley NJ. Changes in single leg countermovement jump force-time characteristics pre, post and 2 days postmatch in elite Girls' youth soccer. *J Strength Cond Res* 2025; 39: e788–e797.
100. Harriss AB, Abbott K, Kimpinski K, et al. An evaluation of heart rate variability in female youth soccer players following soccer heading: a pilot study. *Sports* 2019; 7: 229.
101. Hughes JD, Denton K, Lloyd SR, et al. The impact of soccer match play on the muscle damage response in youth female athletes. *Int J Sports Med* 2018; 39: 343–348.
102. Sedano S and Maroto-Izquierdo S. Effectiveness of different neuromuscular recovery strategies in elite youth female football players. *Sports* 2025; 13: 36.
103. Spalding J. Changes in vertical jump height and sprint time during a congested match schedule in youth female soccer players. *Sport J* 2022; 24: 1–15.
104. Sugawara J, Ogoh N, Watanabe H, et al. Impact of a brief series of soccer matches on vascular conditions in youth women. *Front Physiol* 2024; 15: 1476627.
105. Watson A and Brickson S. Relationships between sport specialization, sleep, and subjective well-being in female adolescent athletes. *Clin J Sport Med Off J Can Acad Sport Med* 2019; 29: 384–390.
106. Wright M, Chesterton P, Wijnbergen M, et al. The effect of a simulated soccer match on anterior cruciate ligament injury risk factors. *Int J Sports Med* 2017; 38: 620–626.
107. Beech J, Jones B, King R, et al. The incidence and burden of injuries in elite English youth female soccer players. *Sci Med Footb* 2022; 6: 605–615.

108. Butera CL, Clark MJ, Georges KJ, et al. Skin cancer risk perception and sunscreen use in adolescent female soccer athletes. *J Dermat Nurses Asso* 2014; 7: 89–96.
109. Clausen MB, Zebis MK, Møller M, et al. High injury incidence in adolescent female soccer. *Am J Sports Med* 2014; 42: 2487–2494.
110. Häggglund M, Waldén M and Ekstrand J. UEFA Injury study—an injury audit of European championships 2006 to 2008. *Br J Sports Med* 2009; 43: 483–489.
111. Inoue Y, Tamura A and Misu S. Age of adolescent female soccer players at first ankle sprain and menarche. *Prog Rehabil Med* 2023; 8: 20230042.
112. Kolstrup LA, Koopmann KU, Nygaard UH, et al. Injuries during football tournaments in 45,000 children and adolescents. *Eur J Sport Sci* 2016; 16: 1167–1175.
113. Le Gall F, Carling C and Reilly T. Injuries in young elite female soccer players: an 8-season prospective study. *Am J Sports Med* 2008; 36: 276–284.
114. Larruskain J, Lekue JA, Angulo P, et al. An injury burden heat map of all men's and women's teams of a professional football club over a decade. *Res Sports Med Print* 2024; 32: 740–750.
115. Lindman I, Högne W, Johansson G, et al. High injury incidence among youth in the World's largest football tournament. *Scand J Med Sci Sports* 2025; 35: e70072.
116. Lislevand M, Andersen TE, Junge A, et al. Injury surveillance during a 2-day national female youth football tournament in Kenya. *Br J Sports Med* 2014; 48: 924–928.
117. Maehlum S, Dahl E and Daljord OA. Frequency of injuries in a youth soccer tournament. *Phys Sportsmed* 1986; 14: 73–79.
118. O'Kane JW, Neradilek M, Polissar N, et al. Risk factors for lower extremity overuse injuries in female youth soccer players. *Orthop J Sports Med* 2017; 5: 2325967117733963.
119. Rosenbaum DA. Issues encountered by physicians during international travel with youth national soccer teams. *Sports Health* 2011; 3: 230–234.
120. Schiff MA, Mack CD, Polissar NL, et al. Soccer injuries in female youth players: comparison of injury surveillance by certified athletic trainers and internet. *J Athl Train* 2010; 45: 238–242.
121. Schneider A, Mayer M, Geißler U, et al. Injuries in male and female adolescent soccer players. *Sportverletz Sportschaden Organ Ges Orthopadisch-Traumatol Sportmed* 2013; 27: 34–38.
122. Schneider AS, Schneider C, Rumpf MC, et al. Knieverletzungen bei männlichen und weiblichen adolescenten fußballspielern. *Sport-Orthop - Sport-Traumatol – Sports Orthop Traumatol* 2013; 29: 40–44.
123. Söderman K, Adolphson J, Lorentzon R, et al. Injuries in adolescent female players in European football: a prospective study over one outdoor soccer season. *Scand J Med Sci Sports* 2001; 11: 299–304.
124. Solstad BE, Sersland A, Torstveit MK, et al. The association between psychosocial factors and reported injuries among youth grassroots soccer players. *Eur J Sport Sci* 2025; 25: e12314.
125. Wezenberg D, Lindblom H, Sonesson S, et al. Prevalence and intensity of pain in male and female amateur football players: a prospective cohort study. *J Sci Med Sport* 2024; 27: 222–227.
126. Cheng AL, Merlo JA, Hunt D, et al. Are hip physical examination findings predictive of future lower-body injury rates in elite adolescent female soccer athletes at minimum 5-year follow-up? *J Sport Rehabil* 2020; 29: 476–482.
127. Chrisman SP, O'Kane JW, Polissar NL, et al. Strength and jump biomechanics of elite and recreational female youth soccer players. *J Athl Train* 2012; 47: 609–615.
128. Clausen MB, Tang L, Zebis MK, et al. Self-reported previous knee injury and low knee function increase knee injury risk in adolescent female football. *Scand J Med Sci Sports* 2016; 26: 919–926.
129. Emery CA. Risk factors for injury in indoor compared with outdoor adolescent soccer. *Am J Sports Med* 2006; 34: 1636–1642.
130. Fältström A, Asker M, Weiss N, et al. Poor knee strength is associated with higher incidence of knee injury in adolescent female football players: the Karolinska football injury cohort. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 2025; 33: 3179–3193.
131. Häggglund M. Risk factors for acute knee injury in female youth football. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 2016; 24: 737–746.
132. O'Kane JW, Gray KE, Levy MR, et al. Shoe and field surface risk factors for acute lower extremity injuries among female youth soccer players. *Clin J Sport Med Off J Can Acad Sport Med* 2016; 26: 245–250.
133. Orzali L, Asker M, Weiss N, et al. Psychological distress, skipped meals, and insufficient sleep, and the occurrence of back pain in adolescent female soccer players: the Karolinska football injury cohort study. *Sports Health* 2025; 19417381251326498. doi:10.1177/19417381251326498
134. Parpa K and Michaelides M. Knee mechanics, strength and flexibility: assessing injury risk in female adolescent soccer players. *J Funct Morphol Kinesiol* 2025; 10: 77.
135. Sedeaud A, De Larochelambert Q, Moussa I, et al. Does an optimal relationship between injury risk and workload represented by the “Sweet spot” really exist? an example from elite French soccer players and pentathletes. *Front Physiol* 2020; 11: 1034.
136. Soligard T, Bahr R and Andersen TE. Injury risk on artificial turf and grass in youth tournament football. *Scand J Med Sci Sports* 2012; 22: 356–361.
137. Soligard T, Grindem H, Bahr R, et al. Are skilled players at greater risk of injury in female youth football? *Br J Sports Med* 2010; 44: 1118–1123.
138. Steffen K, Andersen TE and Bahr R. Risk of injury on artificial turf and natural grass in young female football players. *Br J Sports Med* 2007; 41: i33–i37.
139. Steffen K, Myklebust G, Andersen TE, et al. Self-reported injury history and lower limb function as risk factors for

- injuries in female youth soccer. *Am J Sports Med* 2008; 36: 700–708.
140. Tranaeus U, Ivarsson A, Johnson U, et al. The role of the results of functional tests and psychological factors on prediction of injuries in adolescent female football players. *Int J Environ Res Public Health* 2022; 19: 143.
 141. Watson A, Brickson S, Brooks M, et al. Preseason aerobic fitness predicts in-season injury and illness in female youth athletes. *Orthop J Sports Med* 2017; 5: 232596711772697.
 142. Xiao M, Lemos JL, Hwang CE, et al. High specialization among female youth soccer players is associated with an increased likelihood of serious injury. *Med Sci Sports Exerc* 2021; 53: 2086–2092.
 143. De Ste Croix M, Hughes J, Ayala F, et al. Efficacy of injury prevention training is greater for high-risk vs low-risk elite female youth soccer players. *Am J Sports Med* 2018; 46: 3271–3280.
 144. Häggglund M, Atroshi I, Wagner P, et al. Superior compliance with a neuromuscular training programme is associated with fewer ACL injuries and fewer acute knee injuries in female adolescent football players: secondary analysis of an RCT. *Br J Sports Med* 2013; 47: 974–979.
 145. Leppänen M, Uotila A, Tokola K, et al. Players with high physical fitness are at greater risk of injury in youth football. *Scand J Med Sci Sports* 2022; 32: 1625–1638.
 146. Mandelbaum B, Silvers-Granelli H, Watanabe D, et al. Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes 2-year follow-up. *Am J Sports Med* 2005; 33: 1003–1010.
 147. Soligard T, Myklebust G, Steffen K, et al. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *Br Med J* 2008; 337: a2469.
 148. Steffen K, Myklebust G, Olsen OE, et al. Preventing injuries in female youth football – A cluster-randomized controlled trial. *Scand J Med Sci Sports* 2008; 18: 605–614.
 149. Steffen K, Meeuwisse WH, Romiti M, et al. Evaluation of how different implementation strategies of an injury prevention programme (FIFA 11+) impact team adherence and injury risk in Canadian female youth football players: a cluster-randomised trial. *Br J Sports Med* 2013; 47: 480–487.
 150. Suits W, Darmofal M, Roe O, et al. Comparison of two implementation strategies for anterior cruciate ligament injury prevention in amateur girls' youth soccer: a prospective cohort study. *BMJ Open Sport Exerc Med* 2024; 10: e002031.
 151. Waldén M, Atroshi I, Hedevik H, et al. Prevention of acute knee injuries in adolescent female football players: cluster randomised controlled trial. *Br Med J* 2012; 344: e3042.
 152. Zebis MK, Thorborg K, Andersen LL, et al. Effects of a lighter, smaller football on acute match injuries in adolescent female football: a pilot cluster-randomized controlled trial. *J Sports Med Phys Fitness* 2018; 58: 644–650.
 153. van der Sluis A, Elferink-Gemser MT, Brink MS, et al. Importance of peak height velocity timing in terms of injuries in talented soccer players. *Int J Sports Med* 2015; 36: 327–332.
 154. Malina RM. Maturity status and injury risk in youth soccer players. *Clin J Sport Med Off J Can Acad Sport Med* 2010; 20: 132.
 155. Le Gall F, Carling C, Reilly T, et al. Incidence of injuries in elite French youth soccer players: a 10-season study. *Am J Sports Med* 2006; 34: 928–938.
 156. Ekstrand J, Häggglund M and Fuller CW. Comparison of injuries sustained on artificial turf and grass by male and female elite football players. *Scand J Med Sci Sports* 2011; 21: 824–832.
 157. Ekstrand J, Timpka T and Häggglund M. Risk of injury in elite football played on artificial turf versus natural grass: a prospective two-cohort study. *Br J Sports Med* 2006; 40: 975–980.
 158. Bjørneboe J, Bahr R and Andersen TE. Risk of injury on third-generation artificial turf in Norwegian professional football. *Br J Sports Med* 2010; 44: 794–798.
 159. Schwebel DC, Banaszek MM and McDaniel M. Brief report: behavioral risk factors for youth soccer (football) injury. *J Pediatr Psychol* 2007; 32: 411–416.
 160. Barbero-Álvarez J, Gómez López M, Barbero-Álvarez V, et al. Heart rate and activity profile for young female soccer players. *J Hum Sport Exerc* 2008; 3: 1–11.
 161. Gómez-Villaseca M, Rehbein C, Earp J, et al. Match-related activity profiles of youth female and male football players during world cup classification tournaments and their relationship to body composition. *J Sports Med Phys Fitness* 2024; 64: 784–792.
 162. Harkness-Armstrong A, Till K, Datson N, et al. Whole and peak physical characteristics of elite youth female soccer match-play. *J Sports Sci* 2021; 39: 1320–1329.
 163. Harkness-Armstrong A, Till K, Datson N, et al. Determining age-specific velocity thresholds for elite youth female soccer players. *Sci Med Footb* 2022; 6: 581–588.
 164. Mordillo E and Zago M. Motion characteristics of under-15, under-17 and under-19 Italian youth women soccer matches. *J Sports Med Phys Fitness* 2023; 63: 223–229.
 165. Sydney MG, Wollin M, Chapman DW, et al. Impaired pre-competition wellbeing measures can negatively impact running performance in developmental youth female soccer players. *Biol Sport* 2024; 41: 145–152.
 166. Vescovi JD. Motion characteristics of youth women soccer matches: female athletes in motion (FAiM) study. *Int J Sports Med* 2014; 35: 110–117.
 167. Andersen TB, Bendiksen M, Pedersen JM, et al. Kicking velocity and physical, technical, tactical match performance for U18 female football players—effect of a new ball. *Hum Mov Sci* 2012; 31: 1624–1638.
 168. Harkness-Armstrong A, Till K, Datson N, et al. Influence of match status and possession status on the physical and

- technical characteristics of elite youth female soccer match-play. *J Sports Sci* 2023; 41: 1437–1449.
169. Ørntoft C, Larsen MN, Andersen TB, et al. Technical actions, heart rate, and locomotor activity in 7v7 and 8v8 games for female youth soccer players. *J Strength Cond Res* 2016; 30: 3298–3303.
 170. Arroyo del Bosque R, Amatria Jiménez M and Maneiro Dios R. Análisis técnico del fútbol 7 femenino en etapas formativas, categoría Sub12. *Cuad Psicol Deporte* 2025; 25: 183–202.
 171. Harkness-Armstrong A, Till K, Datson N, et al. Technical characteristics of elite youth female soccer match-play: position and age group comparisons between under 14 and under 16 age groups. *Int J Perform Anal Sport* 2020; 20: 942–959.
 172. Harriss A, Walton DM and Dickey JP. Direct player observation is needed to accurately quantify heading frequency in youth soccer. *Res Sports Med Print* 2018; 26: 191–198.
 173. Reeschke R, Haase FK, Dautzenberg L, et al. Training matters: heading incidence and characteristics in children's and youth football (soccer) players. *Scand J Med Sci Sports* 2023; 33: 1821–1830.
 174. Wahlquist VE, Buckley TA, Caccese JB, et al. Quantitative analysis of ball-head impact exposure in youth soccer players. *J Sports Sci Med* 2023; 22: 591–596.
 175. Maneiro R. Tactical differences between winning and losing teams in elite women's football – INEFC. *Educ Fisica Esports* 2022; 147: 45–54.
 176. Villaseca-Vicuña R, Pérez-Contreras J, Merino-Muñoz P, et al. Influence of the quality of the opponent on physical, technical, and tactical performance of elite female soccer players in international competitions. *Sport Sci Health* 2024; 20: 863–872.
 177. Trewin J, Meylan C, Varley MC, et al. Effect of match factors on the running performance of elite female soccer players. *J Strength Cond Res* 2018; 32: 2002–2009.
 178. Braun H, von Andrian-Werburg J, Schänzer W, et al. Nutrition status of young elite female German football players. *Pediatr Exerc Sci* 2018; 30: 157–167.
 179. Chapelle L, Tassignon B, Aerenhouts D, et al. The hydration status of young female elite soccer players during an official tournament. *J Sports Med Phys Fitness* 2017; 57: 1186–1194.
 180. Cherian KS, Sainoji A, Nagalla B, et al. Energy balance coexists with disproportionate macronutrient consumption across pretraining, during training, and posttraining among Indian junior soccer players. *Pediatr Exerc Sci* 2018; 30: 506–515.
 181. Düppe H, Gärdsell P, Johnell O, et al. Bone mineral density in female junior, senior and former football players. *Osteoporos Int J Establ Result Coop Eur Found Osteoporos Natl Osteoporos Found USA* 1996; 6: 437–441.
 182. Francescato MP, Venuto I, Buoite Stella A, et al. Sex differences in hydration status among adolescent elite soccer players. *J Hum Sport Exerc* 2019; 14: 265–280.
 183. Gibson JC, Stuart-Hill L, Martin S, et al. Nutrition status of junior elite Canadian female soccer athletes. *Int J Sport Nutr Exerc Metab* 2011; 21: 507–514.
 184. Gibson JC, Stuart-Hill LA, Pethick W, et al. Hydration status and fluid and sodium balance in elite Canadian junior women's soccer players in a cool environment. *Appl Physiol Nutr Metab Physiol Appl Nutr Metab* 2012; 37: 931–937.
 185. Klentrou P, McKee K, McKinlay BJ, et al. Circulating levels of bone markers after short-term intense training with increased dairy consumption in adolescent female athletes. *Child Basel Switz* 2021; 8: 961.
 186. McHaffie SJ, Langan-Evans C, Strauss JA, et al. Under-fuelling for the work required? Assessment of dietary practices and physical loading of adolescent female soccer players during an intensive international training and game schedule. *Nutrients* 2023; 15: 4508.
 187. McHaffie SJ, Langan-Evans C, Strauss JA, et al. Energy expenditure, intake and availability in female soccer players via doubly labelled water: are we misrepresenting low energy availability. *Exp Physiol* 2025; 110: 1705–1720.
 188. McKinlay BJ, Wallace PJ, Olansky S, et al. Intensified training in adolescent female athletes: a crossover study of Greek yogurt effects on indices of recovery. *J Int Soc Sports Nutr* 2022; 19: 17–33.
 189. Paludo AC, Gimunová M, Michaelides M, et al. Description of the menstrual cycle status, energy availability, eating behavior and physical performance in a youth female soccer team. *Sci Rep* 2023; 13: 11194.
 190. Soto M, Sánchez-Díaz S, Castillo D, et al. Food intake, body composition and physical assessments in young male and female soccer players. *Rev Espanola Nutr Humana Diet* 2021; 25: e1398.
 191. Vázquez-Díaz F, Aguayo-Muela ADC, Radesca K, et al. Prevalence of disordered eating risk attitudes in youth elite male and female football players. *J Clin Med* 2024; 13: 6178.
 192. Alexe DI, Čaušević D, Čović N, et al. The relationship between functional movement quality and speed, agility, and jump performance in elite female youth football players. *Sports* 2024; 12: 214.
 193. Nicolao A, Pedrinelli A, Zogaib P, et al. Influence of sexual maturation in lactate threshold in female soccer players. *Rev Bras Med Esporte* 2010; 16: 335–338.
 194. Andrade MS, Junqueira MS, Andre Barbosa De Lira C, et al. Age-related differences in torque in angle-specific and peak torque hamstring to quadriceps ratios in female soccer players from 11 to 18 years old: α cross-sectional study. *Res Sports Med Print* 2021; 29: 77–89.
 195. Ávila-Martínez JD, Castro-Malaver MA, Becerra-Patiño BA, et al. Gender differences for the relative age effect on physical skills and emotional intelligence in child volleyball and soccer athletes. *J Funct Morphol Kinesiol* 2024; 9: 244.
 196. Beavan A, Spielmann J, Altmann S, et al. Longitudinal development of physical, perceptual-cognitive and skill

- predictors of talent in academy and professional female football players. *J Sports Sci* 2023; 41: 1952–1959.
197. Becerra-Patiño BA, Paucar Uribe JD, Ávila Martínez JD, et al. Analysis of physical variables as an indicator of performance in a sample of Colombian women's soccer players: influence of being a starter and a non-starter. *J Phys Educ Sport* 2023; 23: 1481–1487.
 198. Bishop C, Pereira LA, Reis VP, et al. Comparing the magnitude and direction of asymmetry during the squat, countermovement and drop jump tests in elite youth female soccer players. *J Sports Sci* 2020; 38: 1296–1303.
 199. Bishop C, Read P, McCubbine J, et al. Vertical and horizontal asymmetries are related to slower sprinting and jump performance in elite youth female soccer players. *J Strength Cond Res* 2021; 35: 56–63.
 200. Brännström A, Yu J-G, Jonsson P, et al. Vitamin D in relation to bone health and muscle function in young female soccer players. *Eur J Sport Sci* 2017; 17: 249–256.
 201. Castagna C. Vertical jump performance in Italian male and female national team soccer players. *J Strength Cond Res* 2013; 27: 1156–1161.
 202. Cherian KS, Shahkar F, Sainoji A, et al. Resting metabolic rate of Indian junior soccer players: testing agreement between measured versus selected predictive equations. *Am J Hum Biol Off J Hum Biol Counc* 2018; 30: e23066.
 203. Datson N, Weston M, Drust B, et al. High-intensity endurance capacity assessment as a tool for talent identification in elite youth female soccer. *J Sports Sci* 2020; 38: 1313–1319.
 204. Datson N, Weston M, Drust B, et al. Reference values for performance test outcomes relevant to English female soccer players. *Sci Med Footb* 2022; 6: 589–596.
 205. De Marco K, Lyons M, Joyce C, et al. The relationship between relative lower-body strength, sprint and change of direction ability in elite youth female soccer athletes. *Int J Sports Sci Coach* 2024; 19: 805–811.
 206. De Oliveira F, Corrêa Neto V, Barbosa R, et al. Power and anaerobic capacity in female soccer: A comparison between different age-Categories. *Apunts Sports Med* 2023; 58: 100421.
 207. Donnelly J, Valentin S, Easton C, et al. Perceptions and experiences of the menstrual cycle amongst elite adult and adolescent football players. *Sci Med Footb* 2025: 1–10. doi:10.1080/24733938.2025.2476485
 208. Doyle B, Browne D and Horan D. Differences in anthropometric and physical performance characteristics between U17, U19, and senior Irish female international football players. *Int J Sports Sci Coach* 2021; 16: 352–359.
 209. Emmonds S, Morris R, Murray E, et al. The influence of age and maturity status on the maximum and explosive strength characteristics of elite youth female soccer players. *Sci Med Footb* 2017; 1: 209–215.
 210. Emmonds S, Till K, Redgrave J, et al. Influence of age on the anthropometric and performance characteristics of high-level youth female soccer players. *Int J Sports Sci Coach* 2018; 13: 779–786.
 211. Emmonds S, Sawczuk T, Scantlebury S, et al. Seasonal changes in the physical performance of elite youth female soccer players. *J Strength Cond Res* 2020; 34: 2636–2643.
 212. Emmonds S, Scantlebury S, Murray E, et al. Physical characteristics of elite youth female soccer players characterized by maturity status. *J Strength Cond Res* 2020; 34: 2321–2328.
 213. Eustace SJ, Page RM and Greig M. Isokinetic strength differences between elite senior and youth female soccer players identifies training requirements. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2019; 39: 45–51.
 214. Fagundes U, Vancini RL, Seffrin A, et al. Adolescent female handball players present greater bone mass content than soccer players: a cross-sectional study. *Bone* 2022; 154: 116217.
 215. Fältström A, Skillgate E, Tranaeus U, et al. Normative values and changes in range of motion, strength, and functional performance over 1 year in adolescent female football players: data from 418 players in the Karolinska football injury cohort study. *Phys Ther Sport* 2022; 58: 106–116.
 216. Ferry B, Duclos M, Burt L, et al. Bone geometry and strength adaptations to physical constraints inherent in different sports: comparison between elite female soccer players and swimmers. *J Bone Miner Metab* 2011; 29: 342–351.
 217. Ferry B, Lespessailles E, Rochcongar P, et al. Bone health during late adolescence: effects of an 8-month training program on bone geometry in female athletes. *Joint Bone Spine* 2013; 80: 57–63.
 218. Gradidge PJ-L, Demitri. A comparative study on the cardiac morphology and vertical jump height of adolescent black South African male and female amateur competitive footballers. *Cardiovasc J Afr* 2018; 29: 32–35.
 219. Haag T, Schneider A, Beckmann C, et al. A test battery to investigate back pain in female soccer players. *Sport Sci Health* 2016; 12: 361–367.
 220. Heijboer WMP, Thijs KM, Weir A, et al. Normal hip strength and range of motion values in youth and adult female national football teams: data from 504 assessments. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2024; 69: 51–58.
 221. Kobus M, Żądzińska E, Michaelides M, et al. Primary headaches and physical performance: a professional youth female soccer team study. *Brain Sci* 2022; 12: 1702.
 222. Kottaras S, Stoikos J, McKinlay BJ, et al. Bone turnover markers and osteokines in adolescent female athletes of high- and low-impact sports compared with nonathletic controls. *Pediatr Exerc Sci* 2023; 35: 41–47.
 223. Lilić A, Petkovic E, Stankovic M, et al. Influence of body composition parameters on explosive power performance in female adolescent football players. *Facta Univ Ser Phys Educ Sport* 2022; 20: 175–185.
 224. Lozano-Berges G, Matute-Llorente Á, Gómez-Bruton A, et al. Bone geometry in young male and female football

- players: a peripheral quantitative computed tomography (pQCT) study. *Arch Osteoporos* 2018; 13: 57.
225. Lozano-Berges G, Matute-Llorente Á, Gómez-Bruton A, et al. Do serum 25-hydroxyvitamin D concentrations affect body composition, physical fitness, bone strength and bone biomarkers in female children and adolescent football players? A one-season study. *Int J Environ Res Public Health* 2022; 19: 15394.
 226. Mainer-Pardos E, Gonzalo-Skok O, Nobari H, et al. Age-related differences in linear sprint in adolescent female soccer players. *BMC Sports Sci Med Rehabil* 2021; 13: 97.
 227. Mainer-Pardos E, Bishop C and Gonzalo-Skok O. Seasonal variations in performance and asymmetry data for jump and change of direction abilities in female soccer players. *J Strength Cond Res* 2024; 38: 1768–1777.
 228. Manson SA, Brughelli M and Harris NK. Physiological characteristics of international female soccer players. *J Strength Cond Res* 2014; 28: 308–318.
 229. Martinho DV, Coelho-E-Silva MJ, Gonçalves Santos J, et al. Body size, fatness and skeletal age in female youth soccer players. *Int J Sports Med* 2023; 44: 711–719.
 230. McCulloch RG, Bailey DA, Whalen RL, et al. Bone density and bone mineral content of adolescent soccer athletes and competitive swimmers. *Pediatr Exerc Sci* 1992; 4: 319–330.
 231. Mujika I, Santisteban J, Impellizzeri F, et al. Fitness determinants of success in men's and women's football. *J Sports Sci* 2009; 27: 107–114.
 232. Norikazu H. Age differences in change-of-direction performance and its subelements in female football players. *Int J Sports Physiol Perform* 2015; 10: 440–445.
 233. Nyland JA, Caborn DNM, Brosky JA, et al. Anthropometric, muscular fitness, and injury history comparisons by gender of youth soccer teams. *J Strength Cond Res* 1997; 11: 92–97.
 234. O'Brien-Smith J, Bennett K, Fransen J, et al. Same or different? A comparison of anthropometry, physical fitness and perceptual motor characteristics in male and female youth soccer players. *Sci Med Footb* 2020; 4: 37–44.
 235. O'Connor C, Chrystal R, Mc Intyre M, et al. Hip adduction and abduction strength values in elite-level male and female youth soccer players: a comparison between sexes, and across age-groups. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2024; 70: 7–14.
 236. Oddsson HR, Friðgeirsdóttir KÝ, Hafliðadóttir L, et al. Differences in anthropometric parameters, physical fitness, and kicking speed in young football players according to performance level, playing position, and relative age effect: a population-based study. *J Strength Cond Res* 2025; 39: e561–e568.
 237. Oyón P, Rubio FJ and Valero A. Young women soccer players. Anthropometric and physiological characteristics. Evolution in a sports season. *Arch Med Deporte* 2016; 33: 24–28.
 238. Pambo P, Adu-Adadey M, Ankrah PT, et al. Electrocardiographic and echocardiographic findings in Ghanaian female soccer players. *Clin J Sport Med Off J Can Acad Sport Med* 2021; 31: e367–e372.
 239. Mainer Pardos E, Bishop C, Gonzalo-Skok O, et al. Associations between inter-Limb asymmetries in jump and change of direction speed tests and physical performance in adolescent female soccer players. *Int J Environ Res Public Health* 2021; 18: 3474.
 240. Parpa K, Katanic B and Michaelides M. Seasonal variation and the effect of the transition period on physical fitness parameters in youth female soccer players. *Sports Basel Switz* 2024; 12: 84.
 241. Petrovic T, Zdravkovic M, Djelic M, et al. Influence of ACE and ACTN3 genes polymorphisms on cardiovascular adaptation in female football players. *Genetika* 2022; 54: 1035–1047.
 242. Philp F, Telford C, Reid D, et al. Normative performance values of modified star excursion balance test and limb symmetry in female adolescent footballers. *Transl Sports Med* 2020; 3: 328–336.
 243. Poehling RA, Tsai M-C, Manson SA, et al. Physical performance development in a female national team soccer program. *J Sci Med Sport* 2021; 24: 597–602.
 244. Ramos GP, Nakamura FY, Penna EM, et al. Comparison of physical fitness and anthropometrical profiles among Brazilian female soccer national teams from U15 to senior categories. *J Strength Cond Res* 2021; 35: 2302–2308.
 245. Raya-González J, Clemente FM and Castillo D. Analyzing the magnitude of interlimb asymmetries in young female soccer players: a preliminary study. *Int J Environ Res Public Health* 2021; 18: 475.
 246. Romero-Caballero A, Varela Olalla D and Loëns-Gutiérrez C. Fitness evaluation in young and amateur soccer players: reference values for vertical jump and aerobic fitness in men and women. *Sci Sports* 2021; 36: 141–e1.
 247. Roso-Moliner A, Gonzalo-Skok O, Villavicencio Álvarez VE, et al. Analysing the influence of speed and jumping performance metrics on the percentage change of direction deficit in adolescent female soccer players. *Life* 2024; 14: 466.
 248. Rusu L, Dobre A, Chivaran A, et al. Relation between muscle force of knee extensors and flexor muscles and sleep quality of women soccer players: a pilot study. *Appl Sci* 2023; 13: 2289.
 249. Sánchez García M, Rodríguez-Fernandez A, Villa-delBosque M, et al. Effects of the menstrual phase on the performance and well-being of female young soccer players. *Cult Cienc Deporte* 2022; 17: 113.
 250. Sašek M, Smajla D and Spudić D. Specificity of curvilinear sprint performance in youth female soccer players: comparison with linear sprint and relationship with vertical jumps. *Int J Perform Anal Sport* 2025; 1–10. doi:10.1080/24748668.2025.2507414
 251. Svytnos G, Michailidis Y, Kotsakis P, et al. Effects of vertical and horizontal jumping asymmetries on linear and change-of-direction speed performance of female soccer players. *Appl Sci* 2024; 14: 3901.

252. Swanson G, Haraldsdottir K, Anderson S, et al. The associations between acute and chronic sleep and aerobic capacity in adolescent female soccer players. *Sport Sci Health* 2023; 20: 435–441.
253. Sweeney L, Lundberg TR, Sweeney C, et al. Biological maturity but not relative age biases exist in female international youth soccer players relative to the general population. *Biol Sport* 2025; 42: 249–256.
254. Martinho VD, Valente-Dos-Santos J, Coelho-E-Silva MJ, et al. Scaling left ventricular mass in adolescent female soccer players. *BMC Pediatr* 2020; 20: 157.
255. Vargas VZ, Motta C, Peres B, et al. Knee isokinetic muscle strength and balance ratio in female soccer players of different age groups: a cross-sectional study. *Phys Sportsmed* 2020; 48: 105–109.
256. Vescovi JD, Rupf R, Brown TD, et al. Physical performance characteristics of high-level female soccer players 12–21 years of age. *Scand J Med Sci Sports* 2011; 21: 670–678.
257. Villaseca-Vicuña R, Morales-Acuña F, Rojas-Trelles L, et al. Comparison of neuromuscular performance, perceived effort, and well-being across menstrual cycle phases in female youth soccer players. *J Phys Educ Sport* 2024; 24: 1814–1822.
258. Wright MD. Changes in sprint-related outcomes during a period of systematic training in a Girls' soccer academy. *J Strength Cond Res* 2019; 33: 793–800.
259. Wright M. Application and interpretation of the yo-yo intermittent recovery test to the long-term physical development of girls association football players. *Sci Med Footb* 2019; 3: 297–306.
260. Zhang J, Lin J, Wei H, et al. Relationships between functional movement quality and sprint and jump performance in female youth soccer athletes of team China. *Child Basel Switz* 2022; 9: 1312.
261. Belamjahad A, Tourny C, Jebabli N, et al. Effects of a pre-season neuromuscular training program vs. an endurance-dominated program on physical fitness and injury prevention in female soccer players. *Sports Med – Open* 2024; 10: 76.
262. Belamjahad A, Tourny C, Hackney AC, et al. Preseason neuromuscular versus endurance training leads to greater improvements in isokinetic strength, muscle damage, and inflammation parameters in highly trained female soccer players. *BMC Sports Sci Med Rehabil* 2025; 17: 107.
263. Darragi M, Zouhal H, Bousselmi M, et al. Effects of in-season strength training on physical fitness and injury prevention in north African elite young female soccer players. *Sports Med – Open* 2024; 10: 94.
264. Fahey JT, Comfort P, Jones P, et al. Effect of 6-week single leg countermovement jump training on force time metrics in elite female youth footballers. *J Sports Sci* 2025; 43: 1196–1206.
265. Fernandez Ortega JA, Los Reyes YGD and Garavito Peña FR. Effects of strength training based on velocity versus traditional training on muscle mass, neuromuscular activation, and indicators of maximal power and strength in girls soccer players. *Apunts Sports Med* 2020; 55: 53–61.
266. González García J, Morencos E, Balsalobre-Fernández C, et al. Effects of 7-Week hip thrust versus back squat resistance training on performance in adolescent female soccer players. *Sports* 2020; 55: 53–61.
267. Hammami MA, Ben Ayed K, Ali A, et al. The effects of a soccer season on anthropometric characteristics, physical fitness, and soccer skills in north African elite female youth soccer players. *Sci Sports* 2023; 38: 401–410.
268. Isla E, Romero-Moraleda B, Moya JM, et al. Effects of a neuromuscular warm-up program in youth female soccer players. *J Hum Kinet* 2021; 79: 29–40.
269. Lesinski M, Prieske O, Chaabene H, et al. Seasonal effects of strength endurance vs. power training in young female soccer athletes. *J Strength Cond Res* 2021; 35: S90–S96.
270. Lindblom H, Waldén M and Häggglund M. No effect on performance tests from a neuromuscular warm-up programme in youth female football: a randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 2012; 20: 2116–2123.
271. Mathisen G and Svein AP. The effect of speed training on sprint and agility performance in female youth soccer players. *J Phys Educ Sport* 2015; 15: 395–399.
272. Pardos-Mainer E, Casajús JA and Gonzalo-Skok O. Adolescent female soccer players' soccer-specific warm-up effects on performance and inter-limb asymmetries. *Biol Sport* 2019; 36: 199–207.
273. Pardos-Mainer E, Casajús JA, Bishop C, et al. Effects of combined strength and power training on physical performance and interlimb asymmetries in adolescent female soccer players. *Int J Sports Physiol Perform* 2020; 15: 1147–1155.
274. Rubley MD, Haase AC, Holcomb WR, et al. The effect of plyometric training on power and kicking distance in female adolescent soccer players. *J Strength Cond Res* 2011; 25: 129–134.
275. Sanchez M, Sanchez-Sanchez J, Bermejo-Martín L, et al. Effects of jump training on youth female soccer Player's physical fitness. *Open Sports Sci J* 2022; 22: e1875399X2208310.
276. Singha P. A study on the effects of 6 weeks of training on body composition, physical fitness and physiological variables of female football players. *J Hum Sport Exerc* 2023; 18: 879–892.
277. Steffen K, Emery CA, Romiti M, et al. High adherence to a neuromuscular injury prevention programme (FIFA 11+) improves functional balance and reduces injury risk in Canadian youth female football players: a cluster randomised trial. *Br J Sports Med* 2013; 47: 794–802.
278. Vargas VZ, Baptista AF, Pereira GOC, et al. Modulation of isometric quadriceps strength in soccer players with transcranial direct current stimulation: a crossover study. *J Strength Cond Res* 2018; 32: 1336–1341.
279. Vescovi JD and VanHeest JL. Effects of an anterior cruciate ligament injury prevention program on performance in adolescent female soccer players. *Scand J Med Sci Sports* 2010; 20: 394–402.

280. Wen X, Song F, Yang L, et al. Small-Sided soccer games promote greater adaptations on vertical jump and change-of-direction deficit and similar adaptations in aerobic capacity than high-intensity interval training in females. *J Sports Sci Med* 2024; 23: 445–454.
281. Whittaker JL. Impact of the FIFA 11 + on the structure of select muscles in adolescent female soccer players. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2015; 16: 228–235.
282. Wright M, Hurst C and Taylor J. Contrasting effects of a mixed-methods high-intensity interval training intervention in girl football players. *J Sports Sci* 2016; 34: 1808–1815.
283. Baxter-Jones ADG, Barbour-Tuck EN, Dale D, et al. The role of growth and maturation during adolescence on team-selection and short-term sports participation. *Ann Hum Biol* 2020; 47: 316–323.
284. Bennett KJM, Novak AR, Fransen J, et al. The prevalence of relative age effects and the influence of the talent pool size on Australian male and female youth football. *J Sports Sci* 2023; 41: 172–180.
285. Brustio PR, Modena R, Boccia G, et al. Youth-to-senior transition in women's and girls' football: towards a better understanding of relative age effects and gender-specific considerations. *PLoS One* 2023; 18: e0283781.
286. Finnegan L, van Rijbroek M, Oliva-Lozano JM, et al. Relative age effect across the talent identification process of youth female soccer players in the United States: influence of birth year, position, biological maturation, and skill level. *Biol Sport* 2024; 41: 241–251.
287. Ginés HJ, Huertas F, García Calvo T, et al. Age and maturation matter in youth elite soccer, but depending on competitive level and gender. *Int J Environ Res Public Health* 2023; 20: 2015.
288. Götze M. Relative age effect in elite German soccer: influence of gender and competition level. *Front Psychol* 2021; 11: 587023.
289. Grossmann B. Relative age effect (RAE) in football talents – the role of youth academies in transition to professional status in Germany. *Int J Perform Anal Sport* 2013; 13: 120–134.
290. Korgaokar AD, Farley RS, Fuller DK, et al. Relative age effect among elite youth female soccer players across the United States. *Sport Mont.* 2018; 16: 37–41.
291. Løgestad P, Steen I and Dalen T. Inevitable relative age effects in different stages of the selection process among male and female youth soccer players. *Sports Basel Switz* 2018; 6: 29.
292. Li Z, Mao L, Steingröver C, et al. Relative age effects in elite Chinese soccer players: implications of the “one-child” policy. *PLoS One* 2020; 15: e0228611.
293. Matušica M. The influence of relative age on the selection of players for elite events in junior football. *J Phys Educ Sport* 2023; 23: 790–799.
294. Morgans R, Radnor J, Nisbet D, et al. The relative age effect across an international soccer programme in male and female players aged 12 years old to seniors. *J Hum Kinet* 2024; 94: 157–169.
295. Pedersen AV, Aune TK, Dalen T, et al. Variations in the relative age effect with age and sex, and over time-elite-level data from international soccer world cups. *PLoS One* 2022; 17: e0264813.
296. Ribeiro E, Barreira J, Carraco D, et al. The relative age effect in under-17, under-20, and adult elite female soccer players. *Sci Med Footb* 2023; 8: 153–160.
297. Romann M. Influences of player nationality, playing position, and height on relative age effects at women's under-17 FIFA world cup. *J Sports Sci* 2013; 31: 32–40.
298. Sedano S, Vaeyens R and Redondo JC. The relative age effect in spanish female soccer players. Influence of the competitive level and a playing position. *J Hum Kinet* 2015; 46: 129–137.
299. Simon C, Carson F, Faber IR, et al. Low prevalence of relative age effects in Luxembourg's male and female youth football. *PLoS One* 2022; 17: e0273019.
300. Smith KL. An examination of relative age and athlete drop-out in female developmental soccer. *Sports Basel Switz* 2022; 10: 79.
301. van den Honert R. Evidence of the relative age effect in football in Australia. *J Sports Sci* 2012; 30: 1365–1374.
302. Vincent J. Gender differences in the relative age effect among US Olympic development program youth soccer players. *J Sports Sci* 2006; 24: 405–413.
303. Castagna C, Impellizzeri FM, Manzi V, et al. The assessment of maximal aerobic power with the multistage fitness test in young women soccer players. *J Strength Cond Res* 2010; 24: 1488–1494.
304. Fältström A, Häggglund M, Hedevik H, et al. The side hop test: validity, reliability, and quality aspects in relation to sex, age and anterior cruciate ligament reconstruction, in soccer players. *Phys Ther Sport Off J Assoc Chart Physiother Sports Med* 2023; 62: 39–45.
305. Lesinski M, Muehlbauer T and Granacher U. Concurrent validity of the gyro inertial sensor system for the assessment of vertical jump height in female sub-elite youth soccer players. *BMC Sports Sci Med Rehabil* 2016; 8: 35.
306. Pardos-Mainer E. Reliability and sensitivity of jumping, linear sprinting and change of direction ability tests in adolescent female football players. *Sci Med Footb* 2019; 3: 183–190.
307. Sonesson S, Lindblom H and Häggglund M. Performance on sprint, agility and jump tests have moderate to strong correlations in youth football players but performance tests are weakly correlated to neuromuscular control tests. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 2021; 29: 1659–1669.
308. Wright MD, Songane F, Emmonds S, et al. Differential ratings of perceived match and training exertion in Girls' soccer. *Int J Sports Physiol Perform* 2020; 15: 1315–1323.
309. Patel R, Nevill A, Cloak R, et al. Relative age, maturation, anthropometry and physical performance characteristics of players within an elite youth football academy. *Int J Sports Sci Coach* 2019; 14: 714–725.

310. Williams CA, Oliver JL and Faulkner J. Seasonal monitoring of sprint and jump performance in a soccer youth academy. *Int J Sports Physiol Perform* 2011; 6: 264–275.
311. Malina R, Eisenmann J, Cumming S, et al. Maturity-associated variation in the growth and functional capacities of youth football (soccer) players 13–15 years. *Eur J Appl Physiol* 2004; 91: 555–562.
312. Smoll FL and Smith RE. *Children and youth in sport: a biopsychosocial perspective*. 2nd ed. Iowa: USA: Kendall/Hunt Pub. Co., 1996, Available: <https://cir.nii.ac.jp/crid/1130282273185193984>.
313. Pérez-González B, León-Quismondo J, Bonal J, et al. The new generation of professional soccer talent is born under the bias of the RAE: relative age effect in international male youth soccer championships. *Child Basel Switz* 2021; 8: 1117.
314. Ando S, Ogoh N, Saito S, et al. Cognitive function among young women's football players in the summer heat. *Transl Sports Med* 2023; 2023: 5516439.
315. Beavan A, Spielmann J, Ehmann P, et al. The development of executive functions in high-level female soccer players. *Percept Mot Skills* 2022; 129: 1036–1052.
316. Burton D, Gillham A and Glenn S. Motivational styles: examining the impact of personality on the self-talk patterns of adolescent female soccer players. *J Appl Sport Psychol* 2011; 23: 413–428.
317. Burton D, Gillham A and Glenn S. The forgotten self-talk attribute: examining perceived effectiveness profiles and patterns of elite female junior soccer players. *Int J Sports Sci Coach* 2013; 8: 357–372.
318. Che X, Zhang Y, Lin J, et al. Two-dimensional and three-dimensional multiple object tracking learning performance in adolescent female soccer players: the role of flow experience reflected by heart rate variability. *Physiol Behav* 2023; 258: 114009.
319. Duguay AM, Loughhead TM and Cook JM. Athlete leadership as a shared process: using a social-network approach to examine athlete leadership in competitive female youth soccer teams. *Sport Psychol* 2020; 33: 189–202.
320. Fältström A, Skillgate E, Weiss N, et al. Lifestyle characteristics in adolescent female football players: data from the Karolinska football injury cohort. *BMC Sports Sci Med Rehabil* 2022; 14: 212.
321. Forsman H, Gråstén A, Blomqvist M, et al. Development and validation of the perceived game-specific soccer competence scale. *J Sports Sci* 2016; 34: 1319–1327.
322. Harris M, Casey LB, Meindl JN, et al. Using behavioral skills training with video feedback to prevent risk of injury in youth female soccer athletes. *Behav Anal Pract* 2020; 13: 811–819.
323. Hintermann M, Romann M, Schmid J, et al. The influence of scanning behaviour on performance during 4v4 small-sided games in youth female football. *J Sports Sci* 2024; 42: 1977–1985.
324. Höner O, Raabe J, Murr D, et al. Prognostic relevance of motor tests in elite girls' soccer: a five-year prospective cohort study within the German talent promotion program. *Sci Med Footb* 2019; 3: 287–296.
325. Kontos AP, Dolese A, Elbin RJ, et al. Relationship of soccer heading to computerized neurocognitive performance and symptoms among female and male youth soccer players. *Brain Inj* 2011; 25: 1234–1241.
326. Kuettel A, Durand-Bush N and Larsen CH. Mental health profiles of Danish youth soccer players: the influence of gender and career development. *J Clin Sport Psychol* 2022; 16: 276–293.
327. Leyhr D, Raabe J, Schultz F, et al. The adolescent motor performance development of elite female soccer players: a study of prognostic relevance for future success in adulthood using multilevel modelling. *J Sports Sci* 2020; 38: 1342–1351.
328. Lyons M, Conlon J, Nimphius S, et al. An athlete's perspective: comparing talent development environments for boys and girls in western Australia youth soccer. *Int J Sports Sci Coach* 2024; 19: 1995–2005.
329. Miller B, Roberts G and Ommundsen Y. Effect of motivational climate on sportpersonship among competitive youth male and female football players. *Scand J Med Sci Sports* 2004; 14: 193–202.
330. Miller B, Roberts G and Ommundsen Y. Effect of perceived motivational climate on moral functioning, team moral atmosphere perceptions, and the legitimacy of intentionally injurious acts among competitive youth football players. *Psychol Sport Exerc* 2005; 6: 461–477.
331. Møllerløkken NE, Lorås H and Pedersen AV. A comparison of Players' and Coaches' perceptions of the coach-created motivational climate within youth soccer teams. *Front Psychol* 2017; 8: 109.
332. Moore EW and Weiller-Abels K. Psychosocial climates differentially predict 12- to 14-Year-Old competitive soccer Players' goal orientations. *Women Sport Phys Act J* 2020; 28: 111–118.
333. Morales-Sánchez V, Caballero-Cerbán M, Postigo-Martín C, et al. Perceived motivational climate determines self-confidence and precompetitive anxiety in young soccer players: analysis by gender. *Sustainability* 2022; 14: 15673.
334. Munroe-Chandler K, Hall C, Fishburne G, et al. Using cognitive general imagery to improve soccer strategies. *Eur J Sport Sci* 2005; 5: 41–49.
335. Olmedilla A, Cánovas M, Olmedilla-Caballero B, et al. Characteristics relevant to sports performance: gender differences in youth soccer. *Cuad Psicol Deporte* 2021; 21: 127–137.
336. Ommundsen Y, Roberts GC, Lemyre P-N, et al. Peer relationships in adolescent competitive soccer: associations to perceived motivational climate, achievement goals and perfectionism. *J Sports Sci* 2005; 23: 977–989.
337. Perroni F, Gallotta M, Pisano S, et al. Gender differences in anthropometric parameters and technical performance of youth soccer players. *Sport Sci Health* 2018; 14: 399–405.

338. Price M and Weiss M. Peer leadership in sport: relationships among personal characteristics, leader behaviors, and team outcomes. *J Appl Sport Psychol* 2011; 23: 49–64.
339. Scanlan TK. Factors influencing the competitive performance expectancies of young female athletes. *J Sport Exerc Psychol* 1979; 1: 212–220.
340. Scanlan TK. Sources of competitive stress in young female athletes. *J Sport Exerc Psychol* 1979; 1: 151–159.
341. Shui Y, Che X, Zhang Y, et al. The role of offensive processes and age development for female soccer players' anticipation. *Sci Rep* 2024; 14: 6054.
342. Steffen K, Pensgaard AM and Bahr R. Self-reported psychological characteristics as risk factors for injuries in female youth football. *Scand J Med Sci Sports* 2009; 19: 442–451.
343. Stephens DE. Predictors of likelihood to aggress in youth soccer: an examination of coed and all-girls teams. *J Sport Behav* 2000; 23: 311–325.
344. Stephens DE. Moral atmosphere and judgments about aggression in girls' soccer: relationships among moral and motivational variables. *J Sport Exerc Psychol* 1996; 18: 158–173.
345. Mahoney MJ, Gabriel TJ and Perkins TS. Psychological skills and exceptional athletic performance. *Sport Psychol* 1987; 1: 181–199.
346. Duda JL. Task and ego orientation in sport questionnaire. *Contemp Educ Psychol* 1989. DOI: 10.1037/t21903-000
347. Gomez P, Ratcliff R and Perea M. A model of the go/no-go task. *J Exp Psychol Gen* 2007; 136: 389–413.
348. Feist J, Datson N, Runswick OR, et al. Visual exploratory activity in elite women's soccer: an analysis of the UEFA women's European championship 2022. *Int J Sport Exerc Psychol* 2025; 23: 281–303.
349. Andersen TR, Kästner B, Arvig M, et al. Monitoring load, wellness, and psychological variables in female and male youth national team football players during international and domestic playing periods. *Front Sports Act Living* 2023; 5: 1197766.
350. Carlsson T, Broman J, Isberg J, et al. Passing-skill training vs. small-sided games for improvement of side-foot kick accuracy among youth female soccer players. *Front Sports Act Living* 2025; 7: 1506563.
351. González-García J and Romero-Moraleda B. The impact of a congested competition schedule on load, recovery, and well-being in under-16 female soccer players: a comparison between starters and non-starters during a development tournament. *Appl Sci* 2024; 14: 8066.
352. Idarraga JV and Valencia-Sánchez W. Original article enhancing ball passing creativity and effectiveness in youth women's football: a single-case study. *J Phys Educ Sport* 2024; 24: 172–186.
353. Lesinski M, Prieske O, Helm N, et al. Effects of soccer training on anthropometry, body composition, and physical fitness during a soccer season in female elite young athletes: A prospective cohort study. *Front Physiol* 2017; 8: 1093.
354. Los Arcos A, Gonzalez Artetxe A, Bayer-Perez U, et al. Consequences of field obstacles on tactical, conditional, and emotional dimensions in young female soccer players during small-sided games. *Proc Inst Mech Eng Part P J Sports Eng Technol* 2023; 239: 70–77.
355. Rumpf M, Schneider A, Schneider C, et al. Training profiles and motivation of male and female youth soccer players. *Int J Sports Sci Coach* 2014; 9: 207–216.
356. Williams C, Winsley R, Rendeiro Pinho G, et al. Prevalence of non-functional overreaching in elite male and female youth academy football players. *Sci Med Footb* 2017; 1: 222–228.
357. Douchet T, Humbertclaude A, Cometti C, et al. Quantifying accelerations and decelerations in elite women soccer players during regular in-season training as an index of training load. *Sports Basel Switz* 2021; 9: 109.
358. Doyle B, Browne D and Horan D. Quantification of internal and external training load during a training camp in senior international female footballers. *Sci Med Footb* 2022; 6: 7–14.
359. Emmonds S, Dalton Barron N, Myhill N, et al. Locomotor and technical characteristics of female soccer players training: exploration of differences between competition standards. *Sci Med Footb* 2023; 7: 189–197.
360. Myhill N, Weaving D, Barrett S, et al. A multi-club analysis of the locomotor training characteristics of elite female soccer players. *Sci Med Footb* 2022; 6: 572–580.
361. Lovell R and Okholm Kryger K. Back to the future-Past learnings for prospective performance, medicine and health research recommendations in WOMEN's football. *Sports Med* 2025; 25: 1–9.