



Review



Cite this article: Cavanagh A, Matthews M. 2025

The heat is on: scaling improvements in photosynthetic thermal tolerance from the leaf to canopy to predict crop yields in a changing climate. *Phil. Trans. R. Soc. B* **380**: 20240235. <https://doi.org/10.1098/rstb.2024.0235>

Received: 27 November 2024

Accepted: 3 February 2025

One contribution of 21 to a theme issue 'Crops under stress: can we mitigate the impacts of climate change on agriculture and launch the 'Resilience Revolution'?'.

Subject Areas:

plant science

Keywords:

photosynthesis, temperature stress, photorespiration, climate change

Author for correspondence:

Amanda Cavanagh

e-mail: a.cavanagh@essex.ac.uk

The heat is on: scaling improvements in photosynthetic thermal tolerance from the leaf to canopy to predict crop yields in a changing climate

Amanda Cavanagh^{1,2} and Megan Matthews^{2,3}

¹School of Life Science, University of Essex - Colchester Campus, Colchester, UK

²Carl R Woese Institute for Genomic Biology, and ³Department of Civil and Environmental Engineering, Grainger College of Engineering, University of Illinois Urbana-Champaign, Urbana, IL, USA

AC, 0000-0001-5918-8093; MM, 0000-0002-5513-9320

Crop production must increase to sustain a growing global population, and this challenge is compounded by increased growing season temperatures and extreme heat events that are already causing significant yield losses in staple crops. Therefore, there is an urgent need to develop strategies to adapt crops to withstand the impacts of a warmer climate. Temperature-sensitive vegetative processes fundamentally related to yield, like photosynthesis, will be impacted by warming throughout the growing season, thus strategies to enhance their resilience hold promise to future-proof crops for a warmer world. Here, we summarize three major strategies to enhance C3 photosynthesis above the thermal optimum: enhanced rubisco activation, modified photorespiration and increased rates of ribulose biphosphate regeneration. We highlight recent experimental evidence demonstrating the efficacy of these strategies, and then use a mechanistic modelling approach to predict the benefit of these engineering strategies on leaf-level carbon assimilation and soybean yield at elevated temperatures. Our approach highlights that these three engineering targets, particularly when combined, can enhance photosynthetic rates and yield under both ambient and elevated temperatures. By targeting multiple aspects of photosynthetic metabolism, we can develop crops that are better equipped to withstand the challenges of a warming climate and contribute to future food security.

This article is part of the theme issue 'Crops under stress: can we mitigate the impacts of climate change on agriculture and launch the 'Resilience Revolution'?'.

1. Introduction

By 2050, global food demand is projected to increase up to an additional 50% of current demand than it is today [1]. However, current trends in agricultural productivity make achieving this target incredibly difficult [2,3]. This challenge is intensified by climate change, which poses significant threats to crop yields worldwide. Terrestrial mean global temperatures have increased approximately 1.6°C since the industrial revolution [4], and current global policies and actions will probably result in an increase from 3.3 to 5.7°C in mean global temperature by the end of the century [4,5]. These rising temperatures have already led to global yield losses in staple crops like maize and wheat [6,7] and further increases are projected to negatively impact the yields of these and other essential crops, such as rice and soybean [8]. Even if we are to achieve zero carbon emissions immediately, the warming caused by the accumulation of atmospheric CO₂ will persist owing to the

cumulative effects of past emissions. Therefore, there is an urgent need to develop strategies that not only improve crop yields but also adapt crops to withstand the adverse effects of a warmer climate.

In the face of both growing global food demands and the pressures of climate change, improving photosynthetic efficiency has emerged as a key strategy to improve crop yields. Photosynthesis is the fundamental process driving plant growth and the production of food, fibre and biofuels and photosynthetic efficiency directly influences crop yields [9–12]. In C3 plants, net photosynthetic carbon assimilation (A_N) is most often constrained by rubisco carboxylation or ribulose biphosphate (RuBP) regeneration capacity [13]. Under current atmospheric $[CO_2]$ and saturating light, the temperature dependence of A_N is co-limited by declines in both rubisco biochemistry and chloroplast electron transport rates [14–16]. Owing to the direct impacts increased chloroplast electron transport, RuBP regeneration and rubisco biochemistry have on the temperature response of photosynthesis, all have been an important target for transgenic modification (table 1).

Rubisco is regulated by a heat-sensitive chaperone protein known as rubisco activase (Rca), which plays a critical role in removing inhibitory sugar-phosphate compounds from rubisco's catalytic sites, thereby sustaining photosynthetic carboxylation efficiency [54–57]. Rubisco remains functional at temperatures exceeding 50°C; however, most Rca isoforms lose activity around 40°C [58,59]. Consequently, the *in vivo* deactivation of rubisco at elevated temperatures is linked to the thermal sensitivity of Rca. This, in turn, triggers photosynthetic inhibition, ultimately leading to yield reductions [56]. Many species have multiple isoforms of Rca with varying heat tolerance [60–62]. Transgenic overexpression of a thermostable Rca isoform confers improved photosynthetic performance and recovery after a short-term high-temperature exposure [17,63] and thermoprotective growth advantages at high growth temperatures [19].

Maintaining rubisco activation (RA) at higher temperatures may also promote rubisco oxygenation, and subsequent photorespiratory losses, although impacts of this remain relatively unexplored. Rubisco's substrate specificity for CO_2 also diminishes with rising temperatures, leading to an increase in RuBP oxygenation rather than carboxylation, and CO_2 release through photorespiration (PR) [64–67]. PR salvages one molecule of 3-phosphoglycerate (3-PGA) from two molecules of the oxygenation product 2-phosphoglycolate (2 PG) with the release of one molecule of CO_2 through a series of enzymatic conversions and transport steps spanning the chloroplast, peroxisome and mitochondria [68]. The photorespiratory pathway is energetically costly, requiring 3.5 ATP and 2 NADPH equivalents to recover RuBP from 2 PG and 3-PGA and releases NH_3 and 25% of the previously fixed CO_2 in the mitochondria. This can result in a dramatic yield drag in C3 crops, with modelled wheat yield losses of 20% across the USA annually and yield reductions of 50% or more observed in warmer growing regions [69]. Photorespiratory flux can be directly manipulated to enhance growth and photosynthesis in heated conditions through the introduction of synthetic 2 PG metabolic pathways ('bypasses') [37] and by the overexpression of key enzymes that may act to optimize photorespiratory flux [50].

As temperatures continue to rise, chloroplast electron transport and RuBP regeneration will become a limiting factor for A_N [15]. Additionally, elevated $[CO_2]$ levels are anticipated to shift the control of photosynthesis from rubisco to RuBP regeneration in C3 crops [70]. Transgenic rice plants with enhanced RuBP regeneration maintain higher rates of A_N after transient heat stress above 35°C [26]. In transgenic soybean, enhanced RuBP regeneration capacity conferred a thermoprotective benefit to in-field warming under elevated CO_2 conditions [31]. However, this strategy has not been fully explored to improve thermotolerance, and mechanistic measurements into the impact of enhanced RuBP regeneration on photosynthetic temperature responses remain unclear.

A key feature of the many strategies to improve photosynthetic carbon gain above the thermal optima is the modularity of the approaches. Many singular targets for manipulation have been identified to enhance RA, increase RuBP regeneration, or lower the cost of PR (table 1). Stacking of improvements made from several targets together will avoid co-limitation by other processes. Multigene constructs targeting the co-overexpression of RuBP regeneration and optimized photorespiratory flux demonstrate that additive gains in photosynthesis and biomass in controlled conditions are possible [20]. However, strategies remain untested over a broad range of temperatures. In this work, we take advantage of mechanistic modelling of photosynthesis at the leaf and canopy level to present a predictive approach to identify synergies between three important targets used to enhance photosynthesis above the thermal optima.

2. Modelling modified photosynthesis above the thermal optimum

The impact of targeted manipulations on net photosynthetic carbon assimilation (A_N) can be predicted using the mechanistic Farquhar von Caemmerer Berry (FvCB) model of photosynthesis [13,71]. The temperature response of light-saturated A_N can be modelled by the minimum of rubisco-limited photosynthesis (A_c) and RuBP-limited photosynthesis (A_j) as:

$$A_c = \frac{(C - \Gamma^*)V_{cmax}}{C + K_C(1 + O/K_O)} - R_l, \quad (2.1)$$

$$A_j = \frac{(C - \Gamma^*)J_{max}}{4C + 8\Gamma^*} - R_l, \quad (2.2)$$

where Γ^* represents the CO_2 compensation point, V_{cmax} is the maximum rate of rubisco carboxylation, J_{max} is the light-saturated electron transport rate, O and C are the oxygen and carbon dioxide concentrations, K_C and K_O are the Michaelis–Menten constants for CO_2 and O_2 , respectively, and R_l represents the carbon dioxide release in the light independent of PR. Although assimilation can also be limited by the rate of triose phosphate utilization, this is primarily observed at saturating CO_2 or low temperatures, and so we have ignored this limitation in our focus on high-temperature responses.

Table 1. Summary of validated targets demonstrating increased rubisco activation (RA), RuBP regeneration (RuBP) or lowered photorespiratory losses (PR) and their impact on photosynthesis and growth at ambient and warming conditions. (Rca = rubisco activase (EC 4.1.1.39); SBPase = sedoheptulose 1,7-bisphosphatase (EC 3.1.3.37); FBPaldolase = (EC 4.1.2.13); Cyt c_6 = algal cytochrome c_6 ; SBPase/FBPase = (EC3.1.3.37 + EC 3.1.3.11); GDC = glycine decarboxylase complex (EC 2.1.2.10, including subunits T, P, L and H); GlcDH = glycolate dehydrogenase (EC 1.1.99.14); GOX = glycolate oxidase (EC 1.1.3.15); WT = wild-type.)

strategy	target	species	impact on photosynthesis and growth
RA	enhanced thermotolerance of Rca	Arabidopsis	overexpression of thermostable Rca isoforms maintains increased rates of rubisco activation state, A_N and growth relative to WT under heat stress [17,18]
		rice	co-overexpression of thermostable Rca and rubisco increase V_{cmax} A_N and biomass relative to WT at 40°C [19]
RuBP	overexpression of SBPase	Arabidopsis, tobacco, tomato, wheat	increased V_{cmax} J_{max} A_N and biomass relative to WT under ambient conditions [20–23]; and increased seed and grain yield [20,24]. Increased J_{max} and growth relative to WT under elevated CO ₂ [25]
		rice	increased A_N > 35°C, and is associated with higher growth rates compared with WT under elevated temperature [26]
	overexpression of FBPaldolase	Arabidopsis, tobacco, tomato	increased V_{cmax} J_{max} A_N and biomass relative to WT under ambient conditions [20]. Increases A_N in tomato [27]. Further increases in A_N and growth evident at elevated CO ₂ [28]
	overexpression of bifunctional SBPase/FBPase	tobacco, lettuce	increases in A_N and biomass relative to controls [29,30]
		soybean	increased A_N V_{cmax} and J_{max} relative to WT. Protective against losses in seed yield loss only when grown under combined elevated temperature and CO ₂ field conditions [31]
	overexpression of algal Cyt c_6	Arabidopsis, tobacco	increased A_N and plant growth [32,33] and water-use [33]
	overexpression of Rieske FeS protein	Arabidopsis	increases in A_{max} J_{max} biomass and seed yield relative to WT controls [34]
	combined overexpression of algal Cyt c_6 with bifunctional SBPase/FBPase or SBPase	tobacco	transient increases in Cyt b_6f activity, but no increases in V_{cmax} J_{max} or growth [35]
PR	chloroplast expression of alternative photorespiration pathways	Arabidopsis, Camelina, tobacco, potato, rice	increases in J_{max} A_{max} biomass and water-use efficiency in-field conditions [36]
	overexpression of glycine decarboxylase (GDC) complex	Arabidopsis, tobacco	fully decarboxylating pathways demonstrate decreases in CO ₂ compensation point [37–43], increases in A_N at supra-ambient temperatures [37,44]. All show environmental-dependent increases in V_{cmax} and J_{max} [38,39,41–43,45]. Growth stimulation, particularly under high light, high temperature conditions [37–44,46]
	overexpression of 2 PG phosphatase (PGLP1)	Arabidopsis	increased A_N , particularly at high light [20,47,48]. Decreased CO ₂ compensation point [48,49]. Enhanced growth relative to WT controls, particularly under high light conditions
	overexpression of glycolate dehydrogenase (GlcDH)	Arabidopsis, potato, Camelina	increased A_{sat} photosynthetic and electron transport at elevated temperatures. Lower CO ₂ compensation point under prolonged heat stress or drought stress [50]
	overexpression of glycolate oxidase (GOX)	rice	decreases in CO ₂ compensation point [40,51], increases in A_N [45,51,52], V_{cmax} J_{max} [45] and photosynthetic electron transport [52]. Biomass and yield stimulation
			increase in A_N compared with WT at high temperature and high light conditions. Lines with 60–100% more GOX expression demonstrate increased plant height under ambient conditions [53]

The temperature dependencies of the six parameters (Γ^* , V_{cmax} J_{max} R_d K_C and K_O) are described by the Arrhenius equation:

$$\text{parameter} = \exp(c - \Delta H_a / RT_k), \quad (2.3)$$

where c is the scaling constant, ΔH_a is the activation energy, R is the molar gas constant (8.314 J mol⁻¹ K⁻¹) and T_k is the leaf temperature in Kelvin [72–74]. The mechanistic basis of enhanced or maintained A_N at increased temperature in plants with modified PR, increased RA or increased RuBP regeneration (RuBP) are likely to be related to changes in either the absolute value or the temperature responses of these six parameters. However, comparisons between improvement strategies are complicated because full temperature datasets on Γ^* , V_{cmax} and J_{max} are only available for tobacco plants expressing a synthetic glycolate metabolic pathway to divert photorespiratory flux [37]. In these plants, enhanced A_N above the thermal optima is probably driven by a change in the apparent ΔH_a for Γ^* such that the parameter is lower in transgenic plants at

Table 2. Parameters used to calculate the temperature response of A_N including the activation energy (ΔH_a) and scaling constant (c). Representative values for unmodified and modified PR plants are from [37]; altered parameters are based on experimental evidence for manipulated Rca [19] (RA) and RuBP regeneration [21,31] (RuBP) as described in text.

parameter	unmodified		RA		PR		RuBP	
	c	ΔH_a	c	ΔH_a	c	ΔH_a	c	ΔH_a
V_{cmax}	27.68	57.96	29.7	62.88				
J_{max}	21.6	41.76					21.78	41.76
Γ^* (Pa)	15.6	34.8			13.3	29.2		

temperatures above 35°C (table 2). Changes in V_{cmax} and J_{max} responses resulting from increased RA or enhanced electron transport can be mechanistically modelled from *in vitro* kinetic responses [16,75,76], but these may not always reflect the phenotype observed from *in planta* manipulations. In the absence of full temperature responses of these parameters for other manipulations, constants and ΔH_a of V_{cmax} and J_{max} can be extrapolated from experimental data to approximate the theoretical phenotype. Enhancing RA state through the overexpression of Rca has increased rice V_{cmax} in heated conditions but not controlled conditions [64], which can be modelled through altered ΔH_a (table 2). Enhanced RuBP regeneration/electron transport was modelled as a 15% increase in wild-type (WT) J_{max} at all temperatures (table 2), consistent with the upper range of reported increases in J_{max} when in plants expressing SBPase or Rieske FeS proteins [21,34,36], and experimental evidence finding no differential changes in the temperature response of J_{max} in transgenic plants with enhanced J_{max} from the overexpression of a bifunctional SBPase/FBPase [31]. The temperature response of A_N in unmodified plants was modelled using the minimum of equations (2.1) and (2.2), using the temperature responses of K_C and K_O from [74] and WT values of V_{cmax} and J_{max} from [37]. The temperature response of rubisco kinetic parameters (i.e. K_C and K_O) are available *in vivo* from *Nicotiana tabacum* (tobacco) and *Arabidopsis* [72,73,77], but not *Glycine max*. To minimize errors in estimating photosynthetic CO_2 assimilation without these key species-specific parameters [78,79] and enable stronger comparison with experimentally derived data in a modelled crop [21,36,37], leaf-level responses are modelled in tobacco. Parameters used to model temperature responses are given in table 2.

3. Photosynthesis above the thermal optimum

When modelled over a temperature range of 10–45°C, strategies that increase RA or modify PR have similar rates of A_N at temperatures leading up to peak photosynthetic rates (figure 1A–C). Maintaining higher rates of V_{cmax} through enhanced RA increases both the maximum photosynthetic rate (A_{max}) and the temperature associated with maximum A_N (T_{max}). Further increases in A_N of 5–12% are predicted at temperatures above T_{max} (figure 1A), in line with empirical work demonstrating a benefit to the expression of a thermostable variant of Rca [19,61,63]. The thermal optimum is unchanged by modified PR, but A_{max} is approximately 2% higher than WT levels, and at temperatures above approximately 30°C, where rubisco more strongly favours oxygenation over carboxylation, modified PR enhances A_N , maintaining rates that are approximately 7% greater than unmodified rates at 40°C (figure 1C). Unsurprisingly, modelled impacts of increased electron transport demonstrate increased A_N only below T_{max} , where rates of A_j are the predominant limitation (figure 1B).

The demonstrated impact that enhanced RA will have on A_N above the thermal optimum is aligned with recent species-independent models suggesting that the global decline in A_N can be accounted for by rubisco deactivation or declines in electron transport [16]. Our modelled responses of enhanced RA (figure 1A) support the position that PR does not limit photosynthetic carbon assimilation as much as rubisco deactivation or declines in electron transport rate, such that a more thermotolerant Rca would lead to an overall increase in photosynthesis at supra-optimal temperatures [16]. However, rubisco deactivation above the thermal optimum could act as a fine-tune control for flux into the photorespiratory pathway at elevated temperatures. Indeed, recent evidence suggests that PR enzymes downstream of rubisco do not acclimate to changes in growth temperature, such that the capacity of photorespiratory flux is scaled to rubisco activity [80]. This highlights a key consideration for improvement strategies, as enhanced RA at high temperature will increase both rubisco oxygenation and carboxylation rates, and increased metabolic flux through the photorespiratory pathway [69,81,82]. Therefore, we examined the impact of combined engineering strategies on the temperature response of photosynthesis (figure 2). When improvement strategies are combined, such that both rubisco deactivation and photorespiratory losses are minimized, modelling reveals an additive benefit to A_N at higher leaf temperatures (figure 2). At 35°C, predictions of increases relative to controls are approximately 4% when PR is modified, and approximately 8% when RA state is maintained, in line with empirical evidence from transgenic plants expressing these strategies [19,37]. Combining benefits from both strategies increases A_N by 12% at 35°C and by 19% at 40°C (figure 2). Owing to the improvements conferred by an increase in J_{max} below the thermal optimum, the largest benefit to assimilation across the breadth of the temperature response is realized when all three strategies are incorporated (figure 2).

4. Scaling from leaf-level photosynthesis to crop yield

To simulate the impact of enhanced leaf-level photosynthetic assimilation on crop yield, FvCB model parameters (table 2) were fed into the modular soybean growth simulator SOYBEAN-BioCRO [83–85] after being adjusted for the soybean V_{cmax} and




Figure 1. Predicted impacts of engineering strategies at and above the thermal optima. The temperature response of net photosynthetic CO_2 assimilation rate (A_N) predicted as the minimum of carboxylation-limited photosynthesis (A_C) or RuBP-regenerated photosynthesis (A_j) based on chloroplast electron transport. Modelled outputs were obtained at 21% O_2 and light-saturated conditions (i.e. $1500 \mu\text{mol m}^{-2}\text{s}^{-1}$) assuming an ambient CO_2 concentration of 410 ppm using parameters described in table 2. (A) The response of enhanced RA through rubisco activase (Rca) overexpression (RA; dashed green lines) compared with unmodified plants (WT; solid grey lines). (B) The response of enhanced electron transport or RuBP regeneration rates (RuBP; dashed blue lines) compared with unmodified plants (WT; solid grey lines). (C) The response of modified PR (dashed purple lines) compared with unmodified plants (WT; solid grey lines).




Figure 2. Combinatorial approaches enhance photosynthetic assimilation at and above the thermal optima. The temperature response of net photosynthetic CO_2 assimilation rate (A_N) predicted as the minimum of carboxylation-limited photosynthesis (A_C) or RuBP regenerated photosynthesis (A_j) based on chloroplast electron transport. Modelled outputs were obtained at 21% O_2 and light-saturated conditions (i.e. $1500 \mu\text{mol m}^{-2}\text{s}^{-1}$) assuming an ambient CO_2 concentration of 410 ppm using parameters described in table 2 for WT (black solid line) and combinations of PR+RA (dashed pink lines), PR+RuBP (dot-dash turquoise lines), RA+RuBP (two-dash pink lines), RA+RuBP (dotted yellow lines) and all manipulations combined (thick dashed blue line).

J_{max} values. SOYBEAN-BioCRO was simulated using 10 years of growing season weather data compiled as in [84] from the NOAA-ESRL SURFRAD site in Bondville, Illinois, USA (2006–2015; available from http://aftp.cmdl.noaa.gov/data/radiation/surf-rad/Bondville_IL/). These years represent a range of environmental fluctuations, including periods of drought, and give a good representation of average conditions in the region. The average air temperature over the 10 growing seasons was 21.28°C . The lowest average for a growing season was 19.9°C and the highest was 22.5°C . Yield projections were obtained for all modification combinations under ambient and elevated temperature (i.e. a constant 5°C increase above ambient), and under ambient (410 ppm) and elevated (610 ppm) atmospheric CO_2 concentrations. Overall, elevated temperature resulted in a 15% decrease in




Figure 3. Predicted impact of photosynthetic modifications on soybean yield. Average changes in end-of-season seed yield relative to unmodified controls calculated for the years 2006–2015 growing seasons (Amb) and modelled with changes to atmospheric CO₂ concentration (Elev; 610 ppm), and temperature (Amb + Heat; +5°C mean temperature increase), or combined climate changes (Elev + Heat; 610 ppm + 5°C mean temperature increase).

predicted yield at 410 ppm CO₂ concentrations (ambient: 5.5 ± 0.9 t ha⁻¹; elevated 4.67 ± 1.15 t ha⁻¹) and a 12% decrease at 610 ppm CO₂ concentrations (ambient: 7.1 ± 0.8 t ha⁻¹; elevated 6.3 ± 1.3 t ha⁻¹). On average, warming negates the predicted yield stimulation from elevated CO₂, as previously demonstrated for soybean using free air concentration enrichment studies [31,86].

Under non-heated conditions, modifications attributed to PR confer a slight advantage to yield compared with controls in both ambient and elevated CO₂ scenarios while those attributed to Rca offer little benefit, in line with predicted benefits to leaf-level photosynthesis at comparable temperatures (figures 2 and 3). While strategies with increased $J_{\max}$ (i.e. RuBP) confer the greatest predicted increase in leaf-level A_N at leaf temperatures <25°C (figure 3), this is not reflected in modelled yield increases under ambient conditions. As predicted from modelling and empirical studies [25,87], RuBP strategies confer a slight increase in yield stimulation relative to WT under elevated CO₂ conditions, with the largest changes relative to controls at elevated CO₂ seen in strategies combining enhanced $J_{\max}$ and lower photorespiratory losses. A 5°C increase in mean growing temperature lowers predicted yield overall, but the relative benefit of strategies to lower the cost of PR (PR+RA and PR+RuBP) is increased relative to ambient conditions (figure 3). However, changes attributed to RA, which demonstrate the largest benefit to light-saturated leaf-level assimilation do not confer a similarly strong predicted yield benefit under heat conditions in either ambient or elevated CO₂ levels (figure 3). Modelled heating scenarios demonstrate large variation in yield estimates for strategies driving mean increases >1 t ha⁻¹ (figure 3). This variation is largely driven by water stress, with the total rainfall for the simulated growing periods ranging from 264 to 657 mm. Strategies that increase A_N are often associated with increased stomatal conductance and subsequent water loss, which will have a more detrimental impact on yield in drier seasons. This is consistent with modelling demonstrating that increases in A_N offer limited benefit to C3 crop yield under water-limited growing conditions, despite driving yield increases under favourable conditions [88].

Scaling assimilation responses from a single leaf to a crop over a growing season adds several layers of complexity including microclimate effects throughout the canopy. Within a canopy, leaves in lower layers are shaded by overlying leaves, and typically receive only 10% of direct sunlight [89]. In crops like soybean that have dense canopies, this results in the average daytime canopy light environment being significantly lower than saturating conditions seen at the top of the canopy. To account for this, SOYBEAN-BIOCro uses a multilayer canopy model where average light is much less in bottom leaf layers compared with upper canopy layers [84,90]. In shaded conditions, the ATP:ADP ratio and redox potential in the chloroplast decline, downregulating Rca activity and leading to rubisco deactivation [54,75,91]. In these conditions, A_N is predominantly limited by electron transport and RuBP regeneration (i.e. A_j ; equation (2.2), which depends on the CO₂ compensation point Γ^*) equation (2.2). Changes in the temperature response of Γ^* account for enhanced thermotolerance of A_N in tobacco plants expressing modified PR [37] (table 2), which may explain the consistent yield benefit attributed to modified PR under all climate scenarios (figure 3). Although rates of rubisco oxygenation and photorespiratory flux will decrease under elevated CO₂ scenarios, prior modelling work has also identified persistent photorespiratory-driven yield losses under elevated CO₂ and increased temperature in wheat and soybean [69]. Therefore, lowering the cost of PR through alternative glycolate metabolism pathways [37–42] or the optimization of flux through targeted enzyme overexpression [47,50,66] remain a strategic target to improve crop yield and resilience to meet mid-century crop production goals.

Both leaf-level and crop model results assume photosynthesis is operating at a steady state. However, within the dynamic environment of the canopy, light is not constant and shaded leaves are exposed to momentary sun-flecks. These rapid periods of light induction feature changes in stromal pH, redox potential, [Mg²⁺] and [ATP], which lead to RA through Rca [92–94]. Transgenic plants overexpressing Rca exhibit faster rates of photosynthetic induction and improved growth in fluctuating light conditions [63,95,96]. Incorporating these faster induction rates would probably increase the estimated yield benefit of enhancing RA under heat stress, beyond what is captured by the crop model where only steady-state photosynthetic rates are considered (figure 3).

5. Conclusions: future directions and a few caveats

Models can be a powerful tool to explore photosynthetic improvement strategies under a variety of conditions in different crops before they are tested in the field. We have focused on sustained increased to atmospheric CO₂ and growing season temperature, but future work can explore timing of heat stress and the response of changes to relative humidity, to refine mechanistic models of predicted benefits [90]. Our modelled responses are informed by physiological data gathered from a range of species, and other changes in metabolic flux associated with photosynthetic manipulation such as changes in mesophyll conductance or light induction remain unaccounted for, despite their key role in regulating carbon assimilation in a dynamic environment. Thus, the true benefits of these strategies remain to be tested under field-relevant conditions in an agronomic crop. However, the recently observed thermoprotective benefit driven by lowering photorespiratory losses in tobacco [37] and potato [44], and the yield advantage for modified field-grown soybean with enhanced RuBP regeneration under combined increases in CO₂ and temperature [31] both suggest that our modelling outcomes are reasonable predictions. Our approach also highlights the importance of gathering mechanistic data on plants with altered photosynthetic metabolism to refine predictions of future performance. In particular, physiological temperature response data are lacking for plants overexpressing Rca or with increased RuBP regeneration capacity or electron transport rates and tends to be lacking even in non-altered metabolism plants from major crop species, complicating mechanistic modelling approaches.

Limitations to photosynthetic carbon assimilation above the thermal optimum, and strategies to mitigate these have been extensively reviewed [15,64,97–101]. Informed by empirical data from plants engineered for thermotolerance, our comparison reveals that predicted and observed benefits to leaf-level photosynthesis above the thermal optima can offer protection against yield losses owing to future warming (figure 3). Therefore, the optimization strategies discussed in this review represent strong candidates to future-proof crop yields in a changing environment.

Ethics. This work did not require ethical approval from a human subject or animal welfare committee.

Data accessibility. The weather data used to model crop responses is available at [102].

The code used to reproduce the results and figures in this manuscript are available in a public GitHub repository: [103].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. A.C.: conceptualization, visualization, writing—original draft, writing—review and editing; M.M.: software, visualization, writing—review and editing.

Both authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. No funding has been received for this article.

References

1. van Dijk M, Morley T, Rau ML, Saghai Y. 2021 A meta-analysis of projected global food demand and population at risk of hunger for the period 2010–2050. *Nat. Food* **2**, 494–501. (doi:10.1038/s43016-021-00322-9)
2. Ray DK, Mueller ND, West PC, Foley JA. 2013 Yield trends are insufficient to double global crop production by 2050. *PLoS ONE* **8**, e66428. (doi:10.1371/journal.pone.0066428)
3. Ray DK, Ramankutty N, Mueller ND, West PC, Foley JA. 2012 Recent patterns of crop yield growth and stagnation. *Nat. Commun.* **3**, 1293. (doi:10.1038/ncomms2296)
4. IPCC. 2023 *Contribution of working groups I, II and III to the sixth assessment report of the intergovernmental panel on climate change*, (eds CW Team, H Lee, J Romero), pp. 35–115. Geneva, Switzerland: IPCC.
5. Climate Action Tracker. 2024 2030 Emissions Gap: CAT projections and resulting emissions gap in meeting the 1.5°C Paris Agreement goal. See <https://climateactiontracker.org/global/cat-emissions-gaps/>.
6. Asseng S *et al.* 2015 Rising temperatures reduce global wheat production. *Nat. Clim. Chang.* **5**, 143–147. (doi:10.1038/nclimate2470)
7. Ortiz-Bobea A, Ault TR, Carrillo CM, Chambers RG, Lobell DB. 2021 Anthropogenic climate change has slowed global agricultural productivity growth. *Nat. Clim. Chang.* **11**, 306–312. (doi:10.1038/s41558-021-01000-1)
8. Zhao C *et al.* 2017 Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl Acad. Sci. USA* **114**, 9326–9331. (doi:10.1073/pnas.1701762114)
9. Kromdijk J, Long SP. 2016 One crop breeding cycle from starvation? How engineering crop photosynthesis for rising CO₂ and temperature could be one important route to alleviation. *Proc. R. Soc. B* **283**, 20152578. (doi:10.1098/rspb.2015.2578)
10. Long SP, Marshall-Colon A, Zhu XG. 2015 Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell* **161**, 56–66. (doi:10.1016/j.cell.2015.03.019)
11. Long SP, Zhu XG, Naidu SL, Ort DR. 2006 Can improvement in photosynthesis increase crop yields? *Plant Cell Environ.* **29**, 315–330. (doi:10.1111/j.1365-3040.2005.01493.x)
12. Simkin AJ, López-Calcano PE, Raines CA. 2019 Feeding the world: improving photosynthetic efficiency for sustainable crop production. *J. Exp. Bot.* **70**, 1119–1140. (doi:10.1093/jxb/ery445)
13. Farquhar GD, von Caemmerer S, Berry JA. 1980 A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta* **149**, 78–90. (doi:10.1007/bf00386231)
14. Busch FA, Sage RF. 2017 The sensitivity of photosynthesis to O₂ and CO₂ concentration identifies strong Rubisco control above the thermal optimum. *New Phytol.* **213**, 1036–1051. (doi:10.1111/nph.14258)
15. Sage RF, Kubien DS. 2007 The temperature response of C3 and C4 photosynthesis. *Plant Cell Environ.* **30**, 1086–1106. (doi:10.1111/j.1365-3040.2007.01682.x)
16. Scafaro AP, Posch BC, Evans JR, Farquhar GD, Atkin OK. 2023 Rubisco deactivation and chloroplast electron transport rates co-limit photosynthesis above optimal leaf temperature in terrestrial plants. *Nat. Commun.* **14**, 2820. (doi:10.1038/s41467-023-38496-4)
17. Kurek I, Chang TK, Bertain SM, Madrigal A, Liu L, Lassner MW, Zhu G. 2007 Enhanced thermostability of *Arabidopsis* rubisco activase improves photosynthesis and growth rates under moderate heat stress. *Plant Cell* **19**, 3230–3241. (doi:10.1105/tpc.107.054171)

18. Kumar A, Li C, Portis AR. 2009 *Arabidopsis thaliana* expressing a thermostable chimeric Rubisco activase exhibits enhanced growth and higher rates of photosynthesis at moderately high temperatures. *Photosynth. Res.* **100**, 143–153. (doi:10.1007/s11120-009-9438-y)
19. Qu Y, Sakoda K, Fukayama H, Kondo E, Suzuki Y, Makino A, Terashima I, Yamori W. 2021 Overexpression of both Rubisco and Rubisco activase rescues rice photosynthesis and biomass under heat stress. *Plant Cell Environ.* **44**, 2308–2320. (doi:10.1111/pce.14051)
20. Simkin AJ, Lopez-Calcano PE, Davey PA, Headland LR, Lawson T, Timm S, Bauwe H, Raines CA. 2017 Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphate aldolase and the photorespiratory glycine decarboxylase-H protein increases CO₂ assimilation, vegetative biomass and seed yield in *Arabidopsis*. *Plant Biotechnol. J.* **15**, 805–816. (doi:10.1111/pbi.12676)
21. Lefebvre S, Lawson T, Fryer M, Zakhleniuk OV, Lloyd JC, Raines CA. 2005 Increased sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis and growth from an early stage in development. *Plant Physiol.* **138**, 451–460. (doi:10.1104/pp.104.055046)
22. Simkin AJ, McAusland L, Headland LR, Lawson T, Raines CA. 2015 Multigene manipulation of photosynthetic carbon assimilation increases CO₂ fixation and biomass yield in tobacco. *J. Exp. Bot.* **66**, 4075–4090. (doi:10.1093/jxb/erv204)
23. Ding F, Wang M, Zhang S, Ai X. 2016 Changes in SBPase activity influence photosynthetic capacity, growth, and tolerance to chilling stress in transgenic tomato plants. *Sci. Rep.* **6**, 32741. (doi:10.1038/srep32741)
24. Driever SM *et al.* 2017 Increased SBPase activity improves photosynthesis and grain yield in wheat grown in greenhouse conditions. *Phil. Trans. R. Soc. B* **372**, 20160384. (doi:10.1098/rstb.2016.0384)
25. Rosenthal DM, Locke AM, Khozaei M, Raines CA, Long SP, Ort DR. 2011 Over-expressing the C3 photosynthesis cycle enzyme sedoheptulose-1-7 bisphosphatase improves photosynthetic carbon gain and yield under fully open air CO₂ fumigation (FACE). *BMC Plant Biol.* **11**, 123. (doi:10.1186/1471-2229-11-123)
26. Feng L, Wang K, Li Y, Tan Y, Kong J, Li H, Li Y, Zhu Y. 2007 Overexpression of SBPase enhances photosynthesis against high temperature stress in transgenic rice plants. *Plant Cell Rep.* **26**, 1635–1646. (doi:10.1007/s00299-006-0299-y)
27. Cai B, Ning Y, Li Q, Li X. 2022 Effects of the chloroplast fructose-1,6-bisphosphate aldolase gene on growth and low-temperature tolerance of tomato. *Int. J. Mol. Sci.* **23**, 728. (doi:10.3390/ijms23020728)
28. Uematsu K, Suzuki N, Iwamae T, Inui M, Yukawa H. 2012 Increased fructose 1,6-bisphosphate aldolase in plastids enhances growth and photosynthesis of tobacco plants. *J. Exp. Bot.* **63**, 3001–3009. (doi:10.1093/jxb/ers004)
29. Ichikawa Y, Tamoi M, Sakuyama H, Maruta T, Ashida H, Yokota A, Shigeoka S. 2010 Generation of transplastomic lettuce with enhanced growth and high yield. *GM Crop.* **1**, 322–326. (doi:10.4161/gmcr.1.5.14706)
30. Miyagawa Y, Tamoi M, Shigeoka S. 2001 Overexpression of a cyanobacterial fructose-1,6-/sedoheptulose-1, 7-bisphosphatase in tobacco enhances photosynthesis and growth. *Nat. Biotechnol.* **19**, 965–969. (doi:10.1038/nbt1001-965)
31. Köhler IH, Ruiz-Vera UM, VanLoocke A, Thomey ML, Clemente T, Long SP, Ort DR, Bernacchi CJ. 2016 Expression of cyanobacterial FBP/SBPase in soybean prevents yield depression under future climate conditions. *J. Exp. Bot.* **68**, 715–726. (doi:10.1093/jxb/erw435)
32. Chida H *et al.* 2007 Expression of the algal cytochrome c6 gene in *Arabidopsis* enhances photosynthesis and growth. *Plant Cell Physiol.* **48**, 948–957. (doi:10.1093/pcp/pcm064)
33. Yadav SK, Khatri K, Rathore MS, Jha B. 2018 Introgression of UfCyt c6, a thylakoid lumen protein from a green seaweed *Ulva fasciata* Delile enhanced photosynthesis and growth in tobacco. *Mol. Biol. Rep.* **45**, 1745–1758. (doi:10.1007/s11033-018-4318-1)
34. Simkin AJ, McAusland L, Lawson T, Raines CA. 2017 Overexpression of the RieskeFeS protein increases electron transport rates and biomass yield. *Plant Physiol.* **175**, 134–145. (doi:10.1104/pp.17.00622)
35. Heyno E, Ermakova M, Lopez-Calcano PE, Woodford R, Brown KL, Matthews JSA, Osmond B, Raines CA, von Caemmerer S. 2022 Rieske FeS overexpression in tobacco provides increased abundance and activity of cytochrome b6f. *Physiol. Plant* **174**, e13803. (doi:10.1111/ppl.13803)
36. López-Calcano PE, Brown KL, Simkin AJ, Fisk SJ, Vialet-Chabrand S, Lawson T, Raines CA. 2020 Stimulating photosynthetic processes increases productivity and water-use efficiency in the field. *Nat. Plants* **6**, 1054–1063. (doi:10.1038/s41477-020-0740-1)
37. Cavanagh AP, South PF, Bernacchi CJ, Ort DR. 2022 Alternative pathway to photorespiration protects growth and productivity at elevated temperatures in a model crop. *Plant Biotechnol. J.* **20**, 711–721. (doi:10.1111/pbi.13750)
38. South PF, Cavanagh AP, Liu HW, Ort DR. 2019 Synthetic glycolate metabolism pathways stimulate crop growth and productivity in the field. *Science* **363**, eaat9077. (doi:10.1126/science.aat9077)
39. Chen G *et al.* 2025 Synthetic photorespiratory bypass improves rice productivity by enhancing photosynthesis and nitrogen uptake. *Plant Cell* **37**, koaf015. (doi:10.1093/plcell/koaf015)
40. Kebeish R *et al.* 2007 Chloroplastic photorespiratory bypass increases photosynthesis and biomass production in *Arabidopsis thaliana*. *Nat. Biotechnol.* **25**, 593–599. (doi:10.1038/nbt1299)
41. Shen BR *et al.* 2019 Engineering a new chloroplastic photorespiratory bypass to increase photosynthetic efficiency and productivity in rice. *Mol. Plant* **12**, 199–214. (doi:10.1016/j.molp.2018.11.013)
42. Wang LM, Shen BR, Li BD, Zhang CL, Lin M, Tong PP, Cui LL, Zhang ZS, Peng XX. 2020 A synthetic photorespiratory shortcut enhances photosynthesis to boost biomass and grain yield in rice. *Mol. Plant* **13**, 1802–1815. (doi:10.1016/j.molp.2020.10.007)
43. Nayak L, Panda D, Dash GK, Lal MK, Swain P, Baig MJ, Kumar A. 2022 A chloroplast glycolate catabolic pathway bypassing the endogenous photorespiratory cycle enhances photosynthesis, biomass and yield in rice (*Oryza sativa* L.). *Plant Sci.* **314**, 111103. (doi:10.1016/j.plantsci.2021.111103)
44. Meacham-Hensold K *et al.* 2024 Shortcutting photorespiration protects potato photosynthesis and tuber yield against heatwave stress. *Glob. Chang. Biol.* **30**, e17595. (doi:10.1111/gcb.17595)
45. Dalal J *et al.* 2015 A photorespiratory bypass increases plant growth and seed yield in biofuel crop *Camelina sativa*. *Biotechnol. Biofuels* **8**, 175. (doi:10.1186/s13068-015-0357-1)
46. Maier A, Fahnstich H, von Caemmerer S, Engqvist MKM, Weber APM, Flüge UI, Maurino VG. 2012 Transgenic introduction of a glycolate oxidative cycle into *A. thaliana* chloroplasts leads to growth improvement. *Front. Plant Sci.* **3**, 38. (doi:10.3389/fpls.2012.00038)
47. López-Calcano PE, Fisk S, Brown KL, Bull SE, South PF, Raines CA. 2019 Overexpressing the H-protein of the glycine cleavage system increases biomass yield in glasshouse and field-grown transgenic tobacco plants. *Plant Biotechnol. J.* **17**, 141–151. (doi:10.1111/pbi.12953)
48. Timm S, Florian A, Arrivault S, Stitt M, Fernie AR, Bauwe H. 2012 Glycine decarboxylase controls photosynthesis and plant growth. *FEBS Lett.* **586**, 3692–3697. (doi:10.1016/j.febslet.2012.08.027)
49. Timm S *et al.* 2015 Mitochondrial dihydriolipoyl dehydrogenase activity shapes photosynthesis and photorespiration of *Arabidopsis thaliana*. *Plant Cell* **27**, 1968–1984. (doi:10.1105/tpc.15.00105)

50. Timm S, Woitschach F, Heise C, Hagemann M, Bauwe H. 2019 Faster removal of 2-phosphoglycolate through photorespiration improves abiotic stress tolerance of *Arabidopsis*. *Plants* **8**, 563. (doi:10.3390/plants8120563)
51. Nölke G, Houdelet M, Kreuzaler F, Peterhänsel C, Schillberg S. 2014 The expression of a recombinant glycolate dehydrogenase polypeptide in potato (*Solanum tuberosum*) plastids strongly enhances photosynthesis and tuber yield. *Plant Biotechnol. J.* **12**, 734–742. (doi:10.1111/pbi.12178)
52. Abbasi AZ *et al.* 2021 Expression of cyanobacterial genes enhanced CO₂ assimilation and biomass production in transgenic *Arabidopsis thaliana*. *PeerJ* **9**, e11860. (doi:10.7717/peerj.11860)
53. Cui LL, Lu YS, Li Y, Yang C, Peng XX. 2016 Overexpression of glycolate oxidase confers improved photosynthesis under high light and high temperature in rice. *Front. Plant Sci.* **7**, 1165. (doi:10.3389/fpls.2016.01165)
54. Amaral J, Lobo AKM, Carmo-Silva E. 2024 Regulation of Rubisco activity in crops. *New Phytol.* **241**, 35–51. (doi:10.1111/nph.19369)
55. Carmo-silva E, Scales JC, Madgwick PJ, Parry MAJ. 2015 Optimizing rubisco and its regulation for greater resource use efficiency. *Plant Cell Environ.* **38**, 1817–1832. (doi:10.1111/pce.12425)
56. Qu Y, Mueller-Cajar O, Yamori W. 2022 Improving plant heat tolerance through modification of Rubisco activase in C3 plants to secure crop yield and food security in a future warming world. *J. Exp. Bot.* **74**, 591–599. (doi:10.1093/jxb/erac340)
57. Salvucci ME, Portis AR, Ogren WL. 1985 A soluble chloroplast protein catalyzes ribulosebisphosphate carboxylase/oxygenase activation *in vivo*. *Photosynth. Res.* **7**, 193–201. (doi:10.1007/bf00037012)
58. Crafts-Brandner SJ, Salvucci ME. 2000 Rubisco activase constrains the photosynthetic potential of leaves at high temperature and CO₂. *Proc. Natl Acad. Sci. USA* **97**, 13430–13435. (doi:10.1073/pnas.230451497)
59. Yamori W, Masumoto C, Fukayama H, Makino A. 2012 Rubisco activase is a key regulator of non-steady-state photosynthesis at any leaf temperature and, to a lesser extent, of steady-state photosynthesis at high temperature. *Plant J.* **71**, 871–880. (doi:10.1111/j.1365-3113.2012.05041.x)
60. Degen GE, Worrall D, Carmo-Silva E. 2020 An isoleucine residue acts as a thermal and regulatory switch in wheat Rubisco activase. *Plant J.* **103**, 742–751. (doi:10.1111/tpj.14766)
61. Scafaro AP, Bautsoens N, den Boer B, Van Rie J, Gallé A. 2019 A conserved sequence from heat-adapted species improves rubisco activase thermostability in wheat. *Plant Physiol.* **181**, 43–54. (doi:10.1104/pp.19.00425)
62. Shivhare D, Mueller-Cajar O. 2017 *In vitro* characterization of thermostable CAM rubisco activase reveals a rubisco interacting surface loop. *Plant Physiol.* **174**, 1505–1516. (doi:10.1104/pp.17.00554)
63. Scafaro AP, Atwell BJ, Muylaert S, Reusel BV, Ruiz GA, Rie JV, Gallé A. 2018 A thermotolerant variant of rubisco activase from a wild relative improves growth and seed yield in rice under heat stress. *Front. Plant Sci.* **9**, 1663. (doi:10.3389/fpls.2018.01663)
64. Moore CE, Meacham-Hensold K, Lemonnier P, Slattery RA, Benjamin C, Bernacchi CJ, Lawson T, Cavanagh AP. 2021 The effect of increasing temperature on crop photosynthesis: from enzymes to ecosystems. *J. Exp. Bot.* **72**, 2822–2844. (doi:10.1093/jxb/erab090)
65. Ogren WL. 1984 Photorespiration: pathways, regulation, and modification. *Annu. Rev. Plant Physiol.* **35**, 415–442. (doi:10.1146/annurev.pp.35.060184.002215)
66. Timm S, Hagemann M. 2020 Photorespiration—how is it regulated and how does it regulate overall plant metabolism? *J. Exp. Bot.* **71**, 3955–3965. (doi:10.1093/jxb/eraa183)
67. Timm S. 2020 The impact of photorespiration on plant primary metabolism through metabolic and redox regulation. *Biochem. Soc. Trans.* **48**, 2495–2504. (doi:10.1042/bst20200055)
68. Peterhansel C, Horst I, Niessen M, Blume C, Kebeish R, Kürkcüoglu S, Kreuzaler F. 2010 Photorespiration. In *The Arabidopsis Book*, vol. **2010**. American Society of Plant Biologists. (doi:10.1199/tab.0130)
69. Walker BJ, VanLoocke A, Bernacchi CJ, Ort DR. 2016 The costs of photorespiration to food production now and in the future. *Annu. Rev. Plant Biol.* **67**, 107–129. (doi:10.1146/annurev-arplant-043015-111709)
70. Bernacchi CJ, Morgan PB, Ort SP. 2005 The growth of soybean under free air [CO₂] enrichment (FACE) stimulates photosynthesis while decreasing *in vivo* Rubisco capacity. *Planta* **220**, 434–446. (doi:10.1007/s00425-004-1320-8)
71. von Caemmerer S. 2000 *Biochemical models of leaf photosynthesis*. Collingwood, Australia: CSIRO. (doi:10.1071/9780643103405)
72. Bernacchi CJ, Portis AR, Nakano H, von Caemmerer S, Long SP. 2002 Temperature response of mesophyll conductance. Implications for the determination of Rubisco enzyme kinetics and for limitations to photosynthesis *in vivo*. *Plant Physiol.* **130**, 1992–1998. (doi:10.1104/pp.008250)
73. Bernacchi CJ, Singsaas EL, Pimentel C, Portis Jr AR, Long SP. 2001 Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant Cell Environ.* **24**, 253–259. (doi:10.1111/j.1365-3040.2001.00668.x)
74. Harley PC, Tenhunen JD. 1991 Modeling the photosynthetic response of C3 leaves to environmental factors. In *Modeling crop photosynthesis – from biochemistry to canopy* (eds KJ Boote, RS Loomis), pp. 1–16. Madison, WI: CSSA. (doi:10.2135/cssaspecpub19.c2)
75. Cen YP, Sage RF. 2005 The regulation of Rubisco activity in response to variation in temperature and atmospheric CO₂ partial pressure in sweet potato. *Plant Physiol.* **139**, 979–990. (doi:10.1104/pp.105.066233)
76. Johnson JE, Berry JA. 2021 The role of cytochrome b6f in the control of steady-state photosynthesis: a conceptual and quantitative model. *Photosynth. Res.* **148**, 101–136. (doi:10.1007/s11120-021-00840-4)
77. Walker B, Ariza LS, Kaines S, Badger MR, Cousins AB. 2013 Temperature response of *in vivo* Rubisco kinetics and mesophyll conductance in *Arabidopsis thaliana*: comparisons to *Nicotiana tabacum*. *Plant Cell Environ.* **36**, 2108–2119. (doi:10.1111/pce.12166)
78. Diaz-espejo A. 2013 New challenges in modelling photosynthesis: temperature dependencies of Rubisco kinetics. *Plant Cell Environ.* **36**, 2104–2107. (doi:10.1111/pce.12192)
79. Sargent D, Amthor JS, Stinziano JR, Evans JR, Whitney SM, Bange MP, Tissue DT, Conaty WC, Sharwood RE. 2024 The importance of species-specific and temperature-sensitive parameterisation of A/C_i models: a case study using cotton (*Gossypium hirsutum* L.) and the automated 'OptiFitACI' R-package. *Plant Cell Environ.* **47**, 1701–1715. (doi:10.1111/pce.14800)
80. Gregory LM, Scott KF, Sharpe LA, Roze LV, Schmieg SC, Hammer JM, Way DA, Walker BJ. 2024 Rubisco activity and activation state dictate photorespiratory plasticity in *Betula papyrifera* acclimated to future climate conditions. *Sci. Rep.* **14**, 26340. (doi:10.1038/s41598-024-77049-7)
81. Badger MR, Collatz GJ. 1977 Studies on the kinetic mechanism of ribulose 1,5-bisphosphate carboxylase and oxygenase reactions, with particular reference to the effect of temperature on kinetic parameters. *Carnegie Inst. Wash. Yearb.* **76**, 355–361.
82. Jordan DB, Ogren WL. 1984 The CO₂/O₂ specificity of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Planta* **161**, 308–313. (doi:10.1007/bf00398720)
83. Lochocki EB, Rohde S, Jaiswal D, Matthews ML, Miguez F, Long SP, McGrath JM. 2022 BioCro II: a software package for modular crop growth simulations. *In Silico Plants* **4**, diac003. (doi:10.1093/insilicoplants/diac003)

84. Matthews ML, Marshall-Colón A, McGrath JM, Lochocki EB, Long SP. 2021 Soybean-BioCro: a semi-mechanistic model of soybean growth. *In Silico Plants* **4**, diab032. (doi:10.1093/insilicoplants/diab032)
85. Miguez FE, Zhu X, Humphries S, Bollero GA, Long SP. 2009 A semimechanistic model predicting the growth and production of the bioenergy crop *Miscanthus × giganteus*: description, parameterization and validation. *GCB Bioenergy* **1**, 282–296. (doi:10.1111/j.1757-1707.2009.01019.x)
86. Ruiz-Vera UM, Siebers M, Gray SB, Drag DW, Rosenthal DM, Kimball BA, Ort DR, Bernacchi CJ. 2013 Global warming can negate the expected CO₂ stimulation in photosynthesis and productivity for soybean grown in the Midwestern United States. *Plant Physiol.* **162**, 410–423. (doi:10.1104/pp.112.211938)
87. Zhu XG, de Sturler E, Long SP. 2007 Optimizing the distribution of resources between enzymes of carbon metabolism can dramatically increase photosynthetic rate: a numerical simulation using an evolutionary algorithm. *Plant Physiol.* **145**, 513–526. (doi:10.1104/pp.107.103713)
88. Leakey ADB, Ferguson JN, Pignon CP, Wu A, Jin Z, Hammer GL, Lobell DB. 2019 Water use efficiency as a constraint and target for improving the resilience and productivity of C3 and C4 crops. *Annu. Rev. Plant Biol.* **70**, 781–808. (doi:10.1146/annurev-arplant-042817-040305)
89. Monteith JL. 1965 Light distribution and photosynthesis in field crops. *Ann. Bot.* **29**, 17–37. (doi:10.1093/oxfordjournals.aob.a083934)
90. He Y, Matthews ML. 2023 Seasonal climate conditions impact the effectiveness of improving photosynthesis to increase soybean yield. *Field Crop. Res.* **296**, 108907. (doi:10.1016/j.fcr.2023.108907)
91. Zhang N, Portis AR. 1999 Mechanism of light regulation of Rubisco: a specific role for the larger Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proc. Natl Acad. Sci. USA* **96**, 9438–9443. (doi:10.1073/pnas.96.16.9438)
92. Hazra S, Henderson JN, Liles K, Hilton MT, Wachter RM. 2015 Regulation of ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) activase: product inhibition, cooperativity, and magnesium activation. *J. Biol. Chem.* **290**, 24222–24236. (doi:10.1074/jbc.M115.651745)
93. Long SP, Taylor SH, Burgess SJ, Carmo-Silva E, Lawson T, De Souza AP, Leonelli L, Wang Y. 2022 Into the shadows and back into sunlight: photosynthesis in fluctuating light. *Annu. Rev. Plant Biol.* **73**, 617–648. (doi:10.1146/annurev-arplant-070221-024745)
94. Zhang N, Kallis RP, Ewy RG, Portis AR. 2002 Light modulation of Rubisco in *Arabidopsis* requires a capacity for redox regulation of the larger Rubisco activase isoform. *Proc. Natl Acad. Sci. USA* **99**, 3330–3334. (doi:10.1073/pnas.042529999)
95. Carmo-Silva AE, Salvucci ME. 2013 The regulatory properties of Rubisco activase differ among species and affect photosynthetic induction during light transitions. *Plant Physiol.* **161**, 1645–1655. (doi:10.1104/pp.112.213348)
96. Suganami M, Suzuki Y, Tazoe Y, Yamori W, Makino A. 2020 Co-overproducing Rubisco and Rubisco activase enhances photosynthesis in the optimal temperature range in rice. *Plant Physiol.* **185**, 108–119. (doi:10.1093/plphys/kiaa026)
97. Ainsworth EA, Ort DR. 2010 How do we improve crop production in a warming world? *Plant Physiol.* **154**, 526–530. (doi:10.1104/pp.110.161349)
98. Dusenke ME, Duarte AG, Way DA. 2019 Plant carbon metabolism and climate change: elevated CO₂ and temperature impacts on photosynthesis, photorespiration and respiration. *New Phytol.* **221**, 32–49. (doi:10.1111/nph.15283)
99. Slattery RA, Ort DR. 2019 Carbon assimilation in crops at high temperatures. *Plant Cell Environ.* **42**, 2750–2758. (doi:10.1111/pce.13572)
100. Yamori W, Hikosaka K, Way DA. 2014 Temperature response of photosynthesis in C3, C4, and CAM plants: temperature acclimation and temperature adaptation. *Photosynth. Res.* **119**, 101–117. (doi:10.1007/s11120-013-9874-6)
101. Cavanagh AP, Ort DR. 2023 Transgenic strategies to improve the thermotolerance of photosynthesis. *Photosynth. Res.* **158**, 109–120. (doi:10.1007/s11120-023-01024-y)
102. NOAA. *Global Monitoring Laboratory*. See https://gml.noaa.gov/aftp/data/radiation/surfrad/Bondville_IL/ (accessed 2025).
103. Soybean-BioCro. *The code in this repository reproduces the results in Matthews et al. Soybean-BioCro: a semi-mechanistic model of soybean growth in silico plants*. GitHub. See <https://github.com/cropsinsilico/Soybean-BioCro> (accessed 2025).